

| Key Indicators for Goa<br>from NFHS-3  | Residence           |        |        | Education                    |                                   |                                    |  | NFHS-2<br>(1998-99) | NFHS-1<br>(1992-93) |
|--|---------------------|--------|--------|------------------------------|-----------------------------------|------------------------------------|--|---------------------|---------------------|
|  | NFHS-3<br>(2005-06) | Urban  | Rural  | No<br>education <sup>4</sup> | <5 years<br>complete <sup>4</sup> | 5-9 years<br>complete <sup>4</sup> | 10 years<br>complete<br>and above <sup>4</sup> |                     |                     |
| <b>Marriage and Fertility</b>  |                     |        |        |                              |                                   |                                    |  |                     |                     |
| 1. Women age 20-24 married by age 18 (%)   | 12.1                | 15.0   | 8.2    | (62.8)                       | *                                 | 12.1                               | 4.8  | 10.1                | 7.2                 |
| 2. Men age 25-29 married by age 21 (%)   | 7.2                 | 7.9    | 6.5    | *                            | *                                 | 8.0                                | 1.5  | na                  | na                  |
| 3. Total fertility rate (children per woman)   | 1.8                 | 1.8    | 1.8    | 2.9                          | 1.6                               | 1.9                                | 1.8  | 1.8                 | 1.9                 |
| 4. Women age 15-19 who were already mothers or pregnant at the time of the survey (%)  | 3.6                 | 5.4    | 1.2    | *                            | *                                 | 4.2                                | 1.2  | na                  | na                  |
| 5. Median age at first birth for women age 25-49   | ≥25                 | ≥25    | ≥25    | 20.7                         | 23.7                              | ≥25                                | ≥25  | 24.1                | 23.7                |
| 6. Married women with 2 living children wanting no more children <sup>1</sup> (%)  | 83.0                | 83.5   | 82.3   | 77.5                         | 78.5                              | 83.0                               | 84.8   | 75.3                | 70.3                |
| 6a. Two sons   | 86.3                | 84.2   | 89.1   | (79.5)                       | *                                 | 83.7                               | 89.1   | 77.3                | 76.0                |
| 6b. One son, one daughter  | 84.8                | 84.9   | 84.7   | (79.5)                       | (73.2)                            | 86.7                               | 86.3   | 86.4                | 76.4                |
| 6c. Two daughters  | 71.7                | 77.4   | 63.8   | *                            | *                                 | (71.6)                             | 74.6   | 52.1                | 48.9                |
| <b>Family Planning (currently married women, age 15-49)</b>  |                     |        |        |                              |                                   |                                    |  |                     |                     |
| <b>Current use</b>   |                     |        |        |                              |                                   |                                    |  |                     |                     |
| 7. Any method (%)  | 48.2                | 51.3   | 44.1   | 56.4                         | 50.9                              | 42.9                               | 48.2   | 47.5                | 47.8                |
| 8. Any modern method (%)   | 37.2                | 38.6   | 35.2   | 52.3                         | 45.4                              | 35.1                               | 31.3   | 35.9                | 37.9                |
| 8a. Female sterilization (%)   | 25.8                | 24.5   | 27.4   | 49.7                         | 42.5                              | 26.8                               | 12.7   | 27.8                | 29.5                |
| 8b. Male sterilization (%)   | 0.1                 | 0.1    | 0.0    | 0.0                          | 0.0                               | 0.0                                | 0.1  | 0.4                 | 1.0                 |
| 8c. IUD (%)  | 2.2                 | 2.6    | 1.8    | 0.5                          | 1.0                               | 2.1                                | 3.2  | 1.9                 | 2.7                 |
| 8d. Pill (%)   | 1.5                 | 1.9    | 1.0    | 0.6                          | 0.0                               | 1.1                                | 2.5  | 0.9                 | 0.7                 |
| 8e. Condom (%)   | 7.5                 | 9.3    | 5.0    | 1.5                          | 1.3                               | 5.1                                | 12.6   | 4.9                 | 3.9                 |
| <b>Unmet need for family planning</b>  |                     |        |        |                              |                                   |                                    |  |                     |                     |
| 9. Total unmet need (%)  | 13.1                | 12.0   | 14.6   | 5.2                          | 10.2                              | 14.9                               | 15.4   | 17.1                | 15.7                |
| 9a. For spacing (%)  | 7.4                 | 6.7    | 8.3    | 3.6                          | 4.1                               | 9.1                                | 8.3  | 7.3                 | 7.8                 |
| 9b. For limiting (%)   | 5.7                 | 5.3    | 6.2    | 1.6                          | 6.1                               | 5.8                                | 7.1  | 9.8                 | 7.9                 |
| <b>Maternal and Child Health</b>   |                     |        |        |                              |                                   |                                    |  |                     |                     |
| <b>Maternity care (for births in the last 3 years)</b>   |                     |        |        |                              |                                   |                                    |  |                     |                     |
| 10. Mothers who had at least 3 antenatal care visits for their last birth (%)  | 95.0                | 96.4   | 93.2   | 79.3                         | (86.0)                            | 96.1                               | 99.4   | 96.3                | 89.8                |
| 11. Mothers who consumed IFA for 90 days or more when they were pregnant with their last child (%)   | 72.0                | 78.9   | 63.3   | 48.0                         | (62.2)                            | 71.2                               | 80.0   | na                  | na                  |
| 12. Births assisted by a doctor/nurse/LHV/ANM/other health personnel (%) <sup>2</sup>  | 94.3                | 94.6   | 93.8   | 77.2                         | (78.9)                            | 95.5                               | 100.0  | 90.8                | 89.2                |
| 13. Institutional births (%) <sup>2</sup>  | 92.6                | 92.6   | 92.4   | 67.5                         | (78.9)                            | 94.9                               | 99.7   | 90.9                | 87.8                |
| 14. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/other health personnel within 2 days of delivery for their last birth (%) <sup>2</sup> | 78.4                | 80.4   | 75.8   | 69.8                         | (68.7)                            | 80.6                               | 80.2   | na                  | na                  |
| <b>Child immunization and vitamin A supplementation<sup>2</sup></b>  |                     |        |        |                              |                                   |                                    |  |                     |                     |
| 15a. Children 12-23 months fully immunized (BCG, measles, and 3 doses each of polio/DPT) (%)   | 78.6                | 83.5   | 73.0   | (52.4)                       | *                                 | 76.9                               | 87.8   | 82.6                | 74.9                |
| 15b. Children 12-23 months who have received BCG (%)   | 96.8                | 97.9   | 95.5   | (89.9)                       | *                                 | 97.2                               | 99.1   | 99.2                | 93.5                |
| 15c. Children 12-23 months who have received 3 doses of polio vaccine (%)  | 87.2                | 88.7   | 85.6   | (70.9)                       | *                                 | 87.5                               | 92.2   | 95.8                | 87.8                |
| 15d. Children 12-23 months who have received 3 doses of DPT vaccine (%)  | 87.5                | 90.7   | 83.8   | (72.7)                       | *                                 | 87.8                               | 91.9   | 93.4                | 86.7                |
| 15e. Children 12-23 months who have received measles vaccine (%)   | 91.2                | 93.8   | 88.3   | (66.6)                       | *                                 | 91.5                               | 99.1   | 84.3                | 77.8                |
| 16. Children age 12-35 months who received a vitamin A dose in last 6 months (%)   | 41.9                | 39.3   | 45.2   | 37.6                         | *                                 | 34.9                               | 49.2   | na                  | na                  |
| <b>Treatment of childhood diseases (children under 3 years)<sup>2</sup></b>  |                     |        |        |                              |                                   |                                    |  |                     |                     |
| 17. Children with diarrhoea in the last 2 weeks who received ORS (%)   | 51.6                | *      | (62.1) | *                            | *                                 | *                                  | *  | 55.6                | 26.9                |
| 18. Children with diarrhoea in the last 2 weeks taken to a health facility (%)   | 71.5                | *      | (79.3) | *                            | *                                 | *                                  | *  | 65.4                | 67.9                |
| 19. Children with acute respiratory infection or fever in the last 2 weeks taken to a health facility (%)  | 83.7                | 88.5   | 79.0   | *                            | *                                 | 88.5                               | 81.4   | na                  | na                  |
| <b>Child Feeding Practices and Nutritional Status of Children<sup>2,3</sup></b>  |                     |        |        |                              |                                   |                                    |  |                     |                     |
| 20. Children under 3 years breastfed within one hour of birth (%)  | 59.7                | 64.7   | 53.4   | 65.9                         | (74.3)                            | 58.9                               | 57.2   | 34.4                | 29.1                |
| 21. Children age 0-5 months exclusively breastfed (%)  | 17.7                | (19.0) | (15.8) | *                            | *                                 | *                                  | (13.9)   | na                  | na                  |
| 22. Children age 6-9 months receiving solid or semi-solid food and breastmilk (%)  | 69.8                | (62.2) | (78.6) | *                            | *                                 | (77.5)                             | (60.3)   | na                  | na                  |
| 23. Children under 3 years who are stunted (%)   | 25.9                | 22.8   | 29.3   | 47.7                         | *                                 | 27.4                               | 16.4   | 21.7                | 34.9                |
| 24. Children under 3 years who are wasted (%)  | 12.8                | 9.6    | 17.2   | 21.0                         | *                                 | 14.1                               | 10.1   | 16.5                | 17.5                |
| 25. Children under 3 years who are underweight (%)   | 21.3                | 15.1   | 28.9   | 38.5                         | *                                 | 25.3                               | 12.2   | 21.3                | 29.3                |
| <b>Nutritional Status of Ever-Married Adults (age 15-49)</b>   |                     |        |        |                              |                                   |                                    |  |                     |                     |
| 26. Women whose Body Mass Index is below normal (%)  | 20.5                | 16.1   | 26.1   | 31.6                         | 25.2                              | 22.5                               | 12.9   | 27.1                | na                  |
| 27. Men whose Body Mass Index is below normal (%)  | 16.8                | 14.5   | 20.0   | 24.5                         | 33.3                              | 19.3                               | 8.1  | na                  | na                  |
| 28. Women who are overweight or obese (%)  | 27.0                | 32.1   | 20.4   | 17.2                         | 15.1                              | 26.1                               | 35.0   | 21.2                | na                  |
| 29. Men who are overweight or obese (%)  | 20.8                | 23.2   | 17.3   | 9.8                          | 3.0                               | 19.9                               | 28.7   | na                  | na                  |
| <b>Anaemia among Children and Adults</b>   |                     |        |        |                              |                                   |                                    |  |                     |                     |
| 30. Children age 6-35 months who are anaemic (%)   | 49.2                | 49.4   | 49.2   | 57.8                         | *                                 | 53.7                               | 41.7   | 53.4                | na                  |
| 31. Ever-married women age 15-49 who are anaemic (%)   | 39.1                | 38.9   | 38.8   | 50.0                         | 40.3                              | 39.8                               | 32.9   | 36.4                | na                  |
| 32. Pregnant women age 15-49 who are anaemic (%)   | 36.9                | 43.3   | (26.1) | *                            | *                                 | (45.6)                             | 23.5   | 35.0                | na                  |
| 33. Ever-married men age 15-49 who are anaemic (%)   | 12.9                | 13.9   | 11.4   | 26.6                         | 18.4                              | 12.4                               | 8.3  | na                  | na                  |
| <b>Knowledge of HIV/AIDS among Ever-Married Adults (age 15-49)</b>   |                     |        |        |                              |                                   |                                    |  |                     |                     |
| 34. Women who have heard of AIDS (%)   | 83.3                | 85.0   | 81.0   | 47.8                         | 70.7                              | 88.1                               | 97.9   | 76.3                | 41.7                |
| 35. Men who have heard of AIDS (%)   | 92.2                | 93.9   | 89.7   | 62.8                         | 82.4                              | 95.9                               | 98.2   | na                  | na                  |
| 36. Women who know that consistent condom use can reduce the chances of getting HIV/AIDS (%)   | 48.8                | 51.6   | 45.1   | 18.8                         | 29.7                              | 47.7                               | 66.7   | na                  | na                  |
| 37. Men who know that consistent condom use can reduce the chances of getting HIV/AIDS (%)   | 57.3                | 63.6   | 48.3   | 19.1                         | 39.0                              | 59.1                               | 69.7   | na                  | na                  |
| <b>Women's Empowerment</b>   |                     |        |        |                              |                                   |                                    |  |                     |                     |
| 38. Currently married women who usually participate in household decisions (%)   | 47.0                | 43.9   | 51.3   | 38.4                         | 42.7                              | 40.8                               | 55.4   | na                  | na                  |
| 39. Ever-married women who have ever experienced spousal violence (%)  | 16.8                | 16.4   | 17.2   | 35.9                         | 20.1                              | 15.8                               | 8.8  | na                  | na                  |

na : Not available

nc : Not calculated because there are no cases

( ) Based on 25-49 unweighted cases

\* Based on fewer than 25 unweighted cases

<sup>1</sup> Excludes pregnant women.

<sup>2</sup> Based on the last 2 births in the 3 years before the survey to ever-married women.

<sup>3</sup> Based on WHO standard.

<sup>4</sup> For children education refers to mother's education. Children with missing information on the mother's education are not included in the education columns.