

NATIONAL FAMILY HEALTH SURVEY (NFHS-4)

INDIA

2015-16

TELANGANA

FEBRUARY 2018

Suggested citation: International Institute for Population Sciences (IIPS) and ICF. 2018.
National Family Health Survey (NFHS-4), India, 2015-16: Telangana. Mumbai: IIPS.

For additional information about the 2015-16 National Family Health Survey (NFHS-4), please contact:

International Institute for Population Sciences, Govandi Station Road, Deonar, Mumbai-400088

Telephone: 022-4237 2442

Fax: 022-2556 3257

E-mail: nfhs42013@gmail.com, director@iips.net

For related information, visit <http://www.rchiips.org/nfhs> or <http://www.iipsindia.org>

CONTRIBUTORS

Sarang Pedgaonkar

Dhananjay Bansod

H. Lhungdim

Sunita Kishor

RESEARCH STAFF

Y. Vaidehi

Poonam V. Kamble

Dnyaneswar D. Kale

CONTENTS

	Page
KEY FINDINGS	
Introduction.....	1
Household Characteristics	3
Education	4
Fertility	5
Family Planning.....	7
Infant and Child Mortality	9
Maternal Health.....	11
Child Health.....	14
Breastfeeding, Nutrition, and Anaemia	17
Adult Health and Health Care.....	20
HIV/AIDS.....	22
Sexual Behaviour	23
Women's Empowerment.....	24
Domestic Violence	26

TABLES

Table 1	Results of the household and individual interviews.....	29
Table 2	Results of the household and individual interviews by district.....	30
Table 3	Household population by age, schooling, residence, and sex	31
Table 4	Household and housing characteristics.....	32
Table 5	Household possessions and land ownership	35
Table 6	School attendance	36
Table 7	Children's living arrangements and orphanhood.....	37
Table 8	Birth registration of children under age five	38
Table 9	Birth registration of children under age five by district.....	39
Table 10	Background characteristics of respondents	40
Table 11	Current fertility	42
Table 12	Fertility by background characteristics.....	43
Table 13	Teenage pregnancy and motherhood	44
Table 14	Birth order	45
Table 15	Birth intervals	46
Table 16	Fertility preferences by number of living children	47
Table 17	Desire not to have any more children.....	48
Table 18	Ideal number of children	49
Table 19	Indicators of sex preference.....	50
Table 20	Knowledge of contraceptive methods	52
Table 21	Current use of contraception by background characteristics.....	54
Table 22	Current use of contraceptive methods by district.....	56
Table 23	Hysterectomy	57

	Page
Table 24	Contraceptive use by men at last sexual intercourse.....58
Table 25	Source of modern contraceptive methods.....60
Table 26	Informed choice.....61
Table 27	Twelve-month contraceptive discontinuation rates *
Table 28	Men's contraception-related perceptions and knowledge.....62
Table 29	Need and demand for family planning among currently married women... ..63
Table 30	Unmet need for family planning by district64
Table 31	Pregnancy outcome65
Table 32	Characteristics of abortions66
Table 33	Age at first marriage.....67
Table 34	Early childhood mortality rates68
Table 35	Early childhood mortality rates by background characteristics69
Table 36	High-risk fertility behaviour70
Table 37	Antenatal care.....71
Table 38	Antenatal care services and information received.....72
Table 39	Antenatal care indicators73
Table 40	Antenatal care indicators by district74
Table 41	Advice received during pregnancy75
Table 42	Pregnancies for which an ultrasound test was done76
Table 43	Pregnancy registration and Mother and Child Protection Card.....78
Table 44	Delivery and postnatal care.....79
Table 45	Delivery and postnatal care by background characteristics.....81
Table 46	Delivery and postnatal care by district.....82
Table 47	Delivery costs and financial assistance.....83
Table 48	Birth order and delivery characteristics by district.....84
Table 49	Timing of first health check after birth for the newborn.....85
Table 50	Trends in maternal care indicators..... *
Table 51	Male involvement in maternal care: Men's report86
Table 52	Vaccinations by background characteristics87
Table 53	Selected vaccinations by district88
Table 54	Prevalence and treatment of symptoms of ARI and fever.....89
Table 55	Prevalence and treatment of diarrhoea.....90
Table 56	Feeding practices during diarrhoea92
Table 57	Knowledge of ORS packets93
Table 58	ICDS coverage and utilization of ICDS services94
Table 59	Utilization of ICDS services during pregnancy and while breastfeeding96
Table 60	Nutritional status of children.....97
Table 61	Initial breastfeeding.....100
Table 62	Breastfeeding status by age101
Table 63	Median duration of breastfeeding and infant and young child feeding (IYCF) practices102
Table 64	Child feeding practices and nutritional status of children by district104
Table 65	Prevalence of anaemia in children.....105

	Page
Table 66	Micronutrient intake among children.....107
Table 67	Presence of iodized salt in household.....109
Table 68	Presence of iodized salt in household by district.....110
Table 69	Women's and men's food consumption111
Table 70	Nutritional status of adults.....112
Table 71	Prevalence of anaemia in adults113
Table 72	Nutritional status and anaemia among children and women by district114
Table 73	Knowledge and prevention of HIV/AIDS.....115
Table 74	Accepting attitudes toward those living with HIV/AIDS.....117
Table 75	Sexual behaviour, HIV testing, blood transfusion, and injections119
Table 76	Knowledge of HIV/AIDS and sexual behaviour among youth.....120
Table 77	Prevalence of tuberculosis121
Table 78	Knowledge and attitudes toward tuberculosis122
Table 79	Health insurance coverage among women and men124
Table 80	Source of health care and health insurance coverage among households125
Table 81	Health problems.....126
Table 82	Health examinations.....128
Table 83.1	Blood pressure status: Women129
Table 83.2	Blood pressure status: Men130
Table 84.1	Random blood glucose levels: Women.....131
Table 84.2	Random blood glucose levels: Men.....132
Table 85	Tobacco and alcohol use by women and men133
Table 86	Methods of menstrual protection134
Table 87	Employment and cash earnings of women and men135
Table 88	Control over and magnitude of women's cash earnings136
Table 89	Decision making.....137
Table 90	Decision making by background characteristics.....138
Table 91	Women's access to money and credit140
Table 92	Ownership of assets.....142
Table 93	Gender role attitudes.....143
Table 94	Gender role attitudes by background characteristics144
Table 95	Experience of physical and sexual violence146
Table 96	Experience of violence during pregnancy.....147
Table 97	Forms of spousal violence148
Table 98	Spousal violence by background characteristics.....149
Table 99	Spousal violence by husband's characteristics and empowerment indicators151
Table 100	Injuries to women due to spousal violence.....153
Table 101	Help seeking154

* Table 27 is not shown because of small number of cases and Table 50 is not shown as trend data for Telangana is not available.

APPENDIX

Estimates of sampling errors.....	155
-----------------------------------	-----

INTRODUCTION

The 2015-16 National Family Health Survey (NFHS-4), the fourth in the NFHS series, provides information on population, health, and nutrition for India as a whole and for each state and union territory. For the first time, NFHS-4 provides district-level estimates for many important indicators. All four NFHS surveys have been conducted under the stewardship of the Ministry of Health and Family Welfare (MoHFW), Government of India. MoHFW designated the International Institute for Population Sciences (IIPS), Mumbai, as the nodal agency for the surveys. Funding for NFHS-4 was provided by the United States Agency for International Development (USAID), the United Kingdom Department for International Development (DFID), the Bill and Melinda Gates Foundation (BMGF), UNICEF, UNFPA, the MacArthur Foundation, and the Government of India. Technical assistance for NFHS-4 was provided by ICF, Maryland, USA. Assistance for the HIV component of the survey was provided by the National AIDS Control Organization (NACO) and the National AIDS Research Institute (NARI), Pune.

Four survey questionnaires—household, woman’s, man’s, and biomarker—were used to collect information in 19 languages using Computer Assisted Personal Interviewing (CAPI). All women age 15-49 and men age 15-54 in the selected sample households were eligible for interview. In the household questionnaire, basic information was collected on all usual members of the household and visitors who stayed in the household the previous night, as well as on socioeconomic characteristics of the household, water and sanitation, health insurance, and number of deaths in the household in the three years preceding the survey. Two versions of the woman’s questionnaire were used in NFHS-4. The first version (district module), which collected information on women’s characteristics, marriage, fertility, contraception, reproductive health, children’s immunizations, and treatment of childhood illnesses, was fielded in the entire sample of NFHS-4 households. Information on these topics is available at the district, state, and national levels. In the second version of the questionnaire (state module), four additional topics, namely, sexual behaviour, HIV/AIDS, husband’s background and women’s work, and domestic violence, were also included. This version was fielded in a subsample of NFHS-4 households designed to provide information only at the state and national levels. The man’s questionnaire covered the man’s characteristics, marriage, number of children, contraception, fertility preferences, nutrition, sexual behaviour, attitudes towards gender roles, HIV/AIDS, and lifestyle. The biomarker questionnaire covered measurements of height, weight, and haemoglobin levels for children; height, weight, haemoglobin, blood pressure, and random blood glucose for women age 15-49 years and men age 15-54 years, and the collection of finger-stick blood for HIV testing in a laboratory. Questionnaire information and biomarkers were collected only with informed consent from the respondents.

The NFHS-4 sample was designed to provide estimates of all key indicators at the national and state levels, as well as estimates for most key indicators at the district level (for all 640 districts in India, as of the 2011 Census). The total sample size of approximately 572,000 households for India was based on the size needed to produce reliable indicator estimates for

each district and for urban and rural areas in districts in which the urban population accounted for 30-70 percent of the total district population. The rural sample was selected through a two-stage sample design with villages as the Primary Sampling Units (PSUs) at the first stage (selected with probability proportional to size), followed by a random selection of 22 households in each PSU at the second stage. In urban areas, there was also a two-stage sample design with Census Enumeration Blocks (CEB) selected at the first stage and a random selection of 22 households in each CEB at the second stage. At the second stage in both urban and rural areas, households were selected after conducting a complete mapping and household listing operation in the selected first-stage units.

The figures of NFHS-4 and earlier rounds may not be strictly comparable due to differences in sample size, and NFHS-4 will be a benchmark for future surveys. NFHS-4 fieldwork for Telangana was conducted in all 10 districts of the state from 6 April to 19 September, 2015 by GFK Mode Private Limited, and collected information from 7,786 households, 7,567 women age 15-49 (including 1,302 women interviewed in PSUs in the state module), and 1,133 men age 15-54. Survey response rates were 95 percent for households, 92 percent for women, and 83 percent for men.

This report presents the key findings of the NFHS-4 survey in Telangana, followed by detailed tables and an appendix on sampling errors. No trend data are available for Telangana, because at the time of NFHS-3 and earlier, Telangana was still a part of the state of Andhra Pradesh. At the time of finalization of this report, wealth quintiles for the country as a whole were not ready. Therefore, on finalization of the national report, breakup of key indicators by wealth quintiles for all states will be provided as an additional document and uploaded on the official website of MoHFW and IIPS.

HOUSEHOLD CHARACTERISTICS

Household composition

In Telangana, 45 percent of surveyed households are in urban areas. On average, households are comprised of about 3.9 members. Fifteen percent of households are headed by women, with 12 percent of the population living in female-headed households.

The vast majority of households in Telangana have household heads who are Hindu (87%). Nine percent of households have household heads who are Muslim and 3 percent of households have Christian household heads.

More than half (57%) of households in Telangana have household heads who belong to an other backward class (OBC), 19 percent belong to a scheduled caste, and 8 percent belong to a schedule tribe. Sixteen percent of Telangana's household heads do not belong to a scheduled caste, scheduled tribe, or other backward class.

One fourth of Telangana's population is under age 15, and 8 percent is age 65 and over. The overall sex ratio of the population is 1,007 females per 1,000 males, and the sex ratio of the population under seven years of age is 907 females per 1,000 males. Nearly all (95%) persons have an *Aadhaar* card.

Among children below 18 years of age, 6 percent have experienced the death of one or both parents. In all, 89 percent of children below 18 years of age live with both parents, 9 percent live with one parent (mostly with their mother), and the remaining 3 percent live with neither parent. Births of 83 percent of children under five years of age were registered with the civil authorities, and 72 percent have a birth certificate.

Housing characteristics

Three-fourths of households in Telangana live in a *pucca* house, and almost every household (98%) has electricity. Half of households have an improved sanitation facility, and 31 percent do not use a sanitation facility, which means that household members practice open defecation. Open defecation is much more common for rural households (51%) than for urban households (7%).

About two-fifths (41%) of households in Telangana have water piped into their dwelling, yard, or plot.

Over three-fourths (78%) of households use an improved source of drinking water, but only 41 percent have water piped into their dwelling, yard, or plot. Urban households (60%) are more likely than rural households (25%) to have water piped into their dwelling, yard, or plot. Twenty-six percent of households treat their drinking water to make it potable (mostly by

using a ceramic, sand, or other water filter or by straining it through a cloth). Almost one-third (31%) of all households use solid fuel for cooking (mainly wood), including 51 percent of rural households. About two-thirds (67%) of households use a clean fuel for cooking.

Selected household possessions

Almost all urban households (96%) and most rural households (86%) in Telangana have a mobile phone. In 92 percent of households, at least one usual member has a bank or post office account, and in 66 percent of households at least one usual member is covered by a health scheme/health insurance. Eighty-two percent of households in Telangana have a BPL card, including 92 percent of rural households and 69 percent of urban households. Fifty-seven percent of households own a house. Agricultural land is owned by 56 percent of rural households and 13 percent of urban households. Overall, about 37 percent of all households in Telangana own agricultural land. Less than one fourth (24%) of the households own farm animals, including 5 percent of urban households.


EDUCATION

School attendance among children

Ninety percent of children age 6-17 years in Telangana attend school (91% in urban areas and 88% in rural areas). School attendance is almost universal (96%) in the age group 6-14 years, but it drops to 78 percent in the age group 15-17 years. There is no gender disparity in school attendance in the age group 6-14 years; however, in the age group 15-17 years, a higher proportion of boys (81%) than girls (76%) are attending school.

Are there gender differentials in children's school attendance?

Percentage of children attending school by age



Literacy, educational attainment, and media exposure

In NFHS-4, literate persons are those who have either completed at least standard six or passed a simple literacy test conducted as part of the survey. According to this measure, 65 percent of women age 15-49 and 83 percent of men age 15-49 are literate.

In Telangana, 33 percent of women and 17 percent of men age 15-49 have never been to school and only 24 percent of women and 31 percent of men age 15-49 have completed 12 or more years of schooling.

Only 24 percent of women and 31 percent of men age 15-49 in Telangana have completed 12 or more years of schooling.

Media exposure is quite high among women and men in Telangana. Eighty-nine percent of women and 92 percent of men watch television at least once a week. Men (61%) are much more likely than women (33%) to read a newspaper or magazine at least once a week. Overall, only 9 percent of women and 4 percent of men are not regularly exposed to any of the forms of media asked about.

FERTILITY

Age at first marriage

The median age at first marriage is 18.8 years among women age 20-49 years and 24.3 years among men age 25-49 years. More than one-fourth (26%) of women age 20-24 years got married before the legal minimum age of 18 and almost one-fifth (18%) of men age 25-29 years got married before the legal minimum age of 21. These data show that early marriage is still quite common in Telangana.


Fertility levels

The total fertility rate (TFR) in Telangana is 1.8 children per woman, implying that the state is well below the replacement level fertility of 2.1 children. Fertility in urban areas, at 1.7 children per woman, is lower than in rural areas by 0.2 children per woman. In keeping with the low fertility, only 4 percent of births in the three years preceding the survey were of birth order 4 or higher and 12 percent were of birth order 3.

Fertility in Telangana does not vary greatly by religion and caste/tribe, but does vary by schooling. Women with no schooling have 2.2 children per woman, almost one-half of a child more than women with 12 or more years of schooling (1.8 children per woman).

How does fertility vary with schooling?

Total fertility rate (children per woman)



Pregnancy outcome

Ninety-two percent of last pregnancies in the five years preceding the survey ended in a live birth, and the remaining 8 percent terminated in foetal wastage, largely due to miscarriage and abortion.

Teenage pregnancy

Among women age 15-19 in Telangana, 11 percent have already begun childbearing, that is, they have already had a live birth or are pregnant with their first child. Less than 1 percent of women age 15-16 years have started childbearing, and this proportion increases sharply to 14 percent among women who are age 18 years old and to 26 percent among women age 19 years. Teenage childbearing is twice as common among rural teenagers (14%) as among urban teenagers (7%).

Birth intervals

The median interval between births in the five years before the survey in Telangana is 29.1 months. Almost two-thirds (64%) of births occur within three years of the previous birth including 14 percent of births that occur within 18 months of the previous birth and 18 percent that occur within 18-23 months. Research shows that waiting at least three years between children reduces the risk of infant mortality.


Almost two-thirds (64%) of births in Telangana occur within three years of the previous birth.

Fertility preferences

Sixty-eight percent of currently married women and 72 percent of currently married men want no more children, are already sterilized, or have a spouse who is sterilized. Among those who want another child, 25 percent of women and 19 percent of men would like to wait at least two years before the next birth. Eighty-four percent of women and 82 percent of men consider the ideal family size to be two or fewer children.

How does son preference affect women's desire for children?

Percentage of currently married women with two children who want no more children



In Telangana, there is some indication of a preference for sons among both women and men. Eleven percent of women and 14 percent of men want more sons than daughters, compared with only 4 percent of both women and men who want more daughters than sons. Despite son preference, however, most women (80%) and men (86%) would like to have at least one daughter.

Women's desire for more children is less affected by their current number of living sons than is men's desire for more children. For example, among women with two children, the desire for no more children varies from 87 percent among those with two sons to 78 percent among those with no son, a range of 9 percentage points. Among men with two children, by comparison, the corresponding range is almost twice as large, at 16 percentage points, with 96 percent of men with 2 sons not wanting any more children, compared with 80 percent of men with no sons.

In Telangana, at least some pregnancies remain unplanned. If all women were to have only the number of children they wanted, the total fertility rate would have been even lower, at 1.6 children per woman, than the current level of 1.8 children per woman.

FAMILY PLANNING


Contraceptive knowledge and use

Knowledge of contraception is almost universal (98%) in Telangana. However, some methods are still less well known. Only 30 percent of currently married women know about IUDs/PPIUDs, 16 percent know about female condoms, and 6 percent know about the lactational amenorrhoea method (LAM). Among all women, 18 percent know about emergency contraception.

The contraceptive prevalence rate (CPR) and the modern contraceptive prevalence rate among currently married women age 15-49 in Telangana are both 57 percent. Female sterilization, at 54 percent, accounts for 95 percent of all contraceptive use in the state. Contraceptive use increases sharply with age from 5 percent for women age 15-19 to 73 percent for women age 40-49.


How many women use family planning?

Percentage of currently married women



What contraceptive methods do women use?

Currently married women



Contraceptive use varies inversely with education. Sixty-six percent of currently married women with no schooling use a contraceptive method, compared with 42 percent of women with 12 or more years of schooling. Contraceptive use increases with number of children from 2 percent among women with no children to 17 percent among women with one child, 72 percent among women with two children, and 76-77 percent among women with three or more children.

In Telangana, the use of modern spacing methods is negligible; no modern spacing method is used by more than 1 percent of currently married women. This is true even in urban areas, where use of modern spacing methods is slightly more common than in rural areas.

The contraceptive prevalence rate among currently married women age 15-49 is 57 percent. The share of female sterilization in contraceptive use is 95 percent.

Contraceptive prevalence does vary widely across the state, ranging from 39 percent in Karimnagar district and 47 percent in Nizamabad district to 69 percent in both Rangareddy and Khammam districts. Male sterilization which is only 2 percent in the state as a whole, is most common in the districts of Warangal (8%) and Karimnagar (4%). Notably, female sterilization is lower in these two districts than in any other district.

Sixty-seven percent of sterilized women and 89 percent of sterilized men had their sterilization operations in the public sector, mainly in a government or municipal hospital (32% and 21%, respectively). Also, 22 percent of men had their sterilization operation in a camp.

Informed choice


Women who know about all available contraceptive methods and their side effects can make better choices about what method to use. Thirty percent of users of selected modern contraceptive methods were ever told by a health or family planning worker about other methods they could use. Even fewer, 24 percent, were told about the possible side effects or problems with their method, and 19 percent were told what to do if they experienced any side effects.

Men's attitudes

About half (47%) of men age 15-49 in Telangana agree that contraception is women's business and a man should not have to worry about it. However, 40 percent think that women using contraception may become promiscuous. Only 53 percent of men know that a condom, if used correctly, protects against pregnancy most of the time.

Contraceptive Prevalence Rate by District

Percentage of currently married women



Unmet need


Unmet need for family planning is defined as the percentage of currently married women who either want to space their next birth or stop childbearing entirely, but are not using contraception. According to this definition, in Telangana, only 7 percent of currently married women have an unmet need for family planning. Currently, 89 percent of the demand for family planning is being satisfied and 88 percent of the demand is being satisfied by modern methods. Across the 10 districts of the state, unmet need for family planning is highest in Hyderabad, at 11 percent, followed by Karimnagar, at 10 percent, and is lowest in Nalgonda, at 4 percent.

INFANT AND CHILD MORTALITY

The infant mortality rate in Telangana is estimated at 30 deaths before the age of one year per 1,000 live births and the under-five mortality rate is estimated at 34 deaths before the age of five years per 1,000 live births. Rural-urban differentials in mortality in Telangana are substantial. For example, the infant mortality rate in rural areas, at 38 deaths per 1,000 live births, is nearly twice as high as in urban areas (20 deaths per 1,000 live births).

Infant Mortality


Deaths per 1,000 live births



Infant and child mortality rates vary greatly by mother's schooling. Among mothers with no schooling, the infant mortality rate, at 69 deaths per 1,000 live births, is more than four times as high as the infant mortality rate among mothers who have completed 10 or more years of schooling, at 15 deaths per 1,000 live births.

High-risk births have higher mortality rates

Deaths in the first year of life per 1,000 live births



Children whose mothers have no schooling are more than four times as likely to die before their first birthday as children whose mothers have completed 10 or more years of schooling.

Note: Categories "Mother's age at birth 30-39 and 40-49" and "birth order 4 or more" are not shown since there are fewer than 250 unweighted person-years of exposure to the risk of death.


MATERNAL HEALTH

Antenatal care

Among mothers who gave birth in the five years preceding the survey, 95 percent received antenatal care (ANC) for their last birth from a skilled provider (86% from a doctor and 9% from an auxiliary nurse midwife (ANM), lady health visitor (LHV), nurse, or midwife). Three percent did not receive any antenatal care. Additionally, 92 percent of mothers registered the pregnancy for their most recent live birth in the five years preceding the survey. Among the registered pregnancies, 89 percent received a Mother and Child Protection Card (MCP Card).

Just over four-fifths (83%) of women received antenatal care during the first trimester of pregnancy, as is recommended. Three-quarters (75%) of mothers had four or more antenatal care visits. For 91 percent of their last births, women received iron and folic acid (IFA) supplements, but only 53 percent consumed them for the recommended 100 days or more. Almost nine out of 10 (89%) last births were protected against neonatal tetanus through tetanus toxoid vaccinations given to the mother. Twenty-three percent of mothers took an intestinal parasite drug during pregnancy.

Four or More Antenatal Care Visits by District
Percentage of last births in the past five years




Seventy-five percent of mothers in Telangana had at least four antenatal care visits for their last birth.

Among women with a live birth in the five years preceding the survey who met with a community health worker in the last three months of pregnancy for their most recent live birth, about two-thirds or more received advice on each of five different areas (93% received advice on breastfeeding, 87% on keeping the baby warm, 80% on the importance of institutional delivery, 68% on cord care, and 64% on family planning).

Even when women receive antenatal care, sometimes they do not receive all the services needed to monitor their pregnancy. In Telangana, however, nearly all women who received antenatal care for their last birth received each of the services needed to monitor their pregnancy: 99 percent each had their weight taken, blood pressure measured, and urine and blood samples taken and 97 percent had their abdomen examined.

How does appropriate antenatal care vary with schooling?

Percentage of last births in the past five years




An ultrasound test was performed during 91 percent of pregnancies in the five years preceding the survey. Pregnancies of women with at least 12 years of schooling were slightly more likely to have an ultrasound test (94%) than pregnancies of women with no schooling (88%). Pregnancies of women with no children at the time of pregnancy were more likely (93%) to have an ultrasound than pregnancies of women with three living children (80%) at the time of pregnancy.

Delivery care

More than nine out of 10 (92%) births take place in a health facility (mostly a private facility) and 8 percent take place at home. Institutional births are more common among women who have received an antenatal check, women in urban areas, women who are having their first birth, women with 12 or more years of schooling, and Muslim and Hindu women. More than nine out of 10 births in the past five years were delivered in a health facility in all districts of Telangana except Adilabad (71%) and Mahbubnagar (78%).

Are babies being delivered safely?

Percentage of births in the past five years



Ninety-one percent of births during the past five years took place with assistance from a skilled provider, and another 3 percent were delivered by a traditional birth attendant.


Fifty-eight percent of births during the five years preceding the survey were delivered by caesarean section. Two-fifths of caesarean sections (23% of all births) were emergency caesarean sections.

For 97 percent of home births, a clean blade was used to cut the cord, as is recommended. A disposable delivery kit was used for 42 percent of home deliveries. For 80 percent of home births the recommendation that the baby be immediately wiped dry and then wrapped without being bathed first was followed.

Among women who had a live birth in the five years preceding the survey that was delivered in a health facility, 12 percent received financial assistance under the *Janani Suraksha Yojana* (JSY). Rural women were more likely than urban women to receive financial assistance under JSY (14% versus 10%). Scheduled caste women were more likely than any other caste/tribe group of women to receive financial assistance under JSY.

Institutional Delivery by District

Percentage of births in the past five years



More than nine out of 10 (92%) children in Telangana who were born in the past five years were born in a health facility.

Postnatal care

Early postnatal care for a mother helps safeguard her health and can reduce maternal mortality. In Telangana, 87 percent of mothers had a postnatal check after their last birth and 83 percent of mothers had a postnatal check within two days of the birth, as is recommended. Postnatal care is most common following births in a health facility; however, almost 1 in 10 births in both public and private health facilities were not followed by a postnatal check for the mother. One-third of home births were followed by a postnatal check for the mother within two days of birth.

Only 26 percent of last births in the five years preceding the survey received a health check in the first two days after birth.

Male involvement in maternal care

Ninety-five percent of men with a child under three years of age said that the youngest child's mother received antenatal care. More than two-thirds (69%) of men with a child under three years said they were present during at least one antenatal check-up received by the child's mother (72% in urban areas and 67% in rural areas), but only 33 percent of men were told by a health provider or health worker what to do if the mother had a pregnancy complication. Only 24-42 percent of men were told about the signs of specific pregnancy complications (convulsions, vaginal bleeding, prolonged labour, high blood pressure, and severe abdominal pain).

A majority of men with a child less than three years of age were given information about various aspects of maternal care by a health provider or worker: 80 percent were told about the importance of proper nutrition for the mother during pregnancy, 53 percent were told about the importance of delivering the baby in a health facility, and 52 percent were told about family planning or delaying the next child.


CHILD HEALTH

Vaccination of children

More than two-thirds (68%) of children age 12-23 months received all basic vaccinations against six major childhood illnesses (tuberculosis, diphtheria, pertussis, tetanus, polio, and measles) at any time before the survey. However, most children are at least partially vaccinated; only 1 percent have not received any vaccinations at all.

Vaccination Coverage

Percentage of children 12-23 months receiving vaccinations




Ninety-seven percent of children have received a BCG vaccination and at least three-fourths have received each of the other basic vaccinations (75% have received at least the recommended three doses of polio vaccine, 88% have received the three recommended doses of DPT vaccine, and 91% have been vaccinated against measles). There is considerable dropout between the first and third doses of the polio vaccine (from 96% to 75%) and of the DPT vaccine (from 96% to 88%).

Ninety-four percent of children age 12-23 months have received at least one dose of hepatitis B vaccine, but far fewer, 71 percent, have received all three recommended doses of hepatitis B vaccine.

Coverage with all basic vaccinations does not vary greatly by most background characteristics; however, coverage is much higher for male children (72%) than for female children (64%).

Coverage of All Basic Vaccinations by District

Percentage of children 12-23 months



Coverage with all basic vaccinations is higher for male children than for female children.

Childhood illnesses

In the two weeks before the survey, 2 percent of children under age five years had symptoms of an acute respiratory infection (cough accompanied by (1) short, rapid breathing that is chest related and/or (2) difficult breathing that is chest related). Of these children, 88 percent were taken to a health facility or health provider.

Seventeen percent of children under age five were reported to have had fever in the two weeks preceding the survey; 75 percent of these children were taken to a health facility or provider for advice or treatment.


Overall, 8 percent of children under age five years had diarrhoea in the two weeks preceding the survey. Seventy-four percent of these children were taken to a health facility or health provider. Almost 9 out of 10 mothers (88%) of young children born in the past five years have heard of oral rehydration salt (ORS) packets for the treatment of diarrhoea, but only 57 percent of children with diarrhoea were given ORS. Eighteen percent of children with diarrhoea were given gruel and about two-thirds (65%) were given some type of oral rehydration therapy (ORT). Thirty-two percent were given zinc supplements. Sixteen percent of children with diarrhoea did not receive any type of treatment at all.

To reduce dehydration and minimise the effects of diarrhoea on nutritional status, it is recommended that normal feeding of children be continued when they have diarrhoea and that the amount of fluids given should be increased. However, in Telangana only 5 percent of children with diarrhoea received more to drink than normal. Over one-third (34%) of children received the same amount to drink as usual. Of greater concern, 61 percent of children with diarrhoea were given less to drink or nothing at all to drink. Only 24 percent of children with diarrhoea were given the same amount of food or more food, as recommended. Overall, much less than half (43%) of children with diarrhoea were given ORT and continued feeding.

Integrated Child Development Services (ICDS)

The ICDS programme provides nutrition and health services for children under age six years and for pregnant and breastfeeding women, as well as early childhood care or preschool activities for children age 3-5 years. These services are provided through community-based *anganwadi* centres.

How many children received *anganwadi* centre services?
Percentage of age-eligible children receiving services



Fifty-seven percent of children under 6 years of age receive services of some kind from an *anganwadi* centre. The most common services that age-eligible children receive are supplementary food (56%), growth monitoring (51%), health check-ups (44%), and immunizations (43%). The service that is least likely to be accessed is early childhood care or preschool (40% of children age 3-5 years). Mothers of about two-thirds (64%) of the children who were weighed at an *anganwadi* centre received counselling from an *anganwadi* worker or an ANM. Mothers of 62 percent of children under age six years received at least one service from an *anganwadi* centre during pregnancy, and 59 percent of their mothers received any service while breastfeeding.

BREASTFEEDING, NUTRITION, AND ANAEMIA

Infant feeding

Although breastfeeding is nearly universal in Telangana, only 67 percent of children under six months are exclusively breastfed, as the World Health Organization (WHO) recommends. Seventy-four percent are put to the breast within the first day of life, but only 36 percent started breastfeeding in the first hour of life (as recommended). Thus, many infants are still deprived of the highly nutritious first milk (colostrum) and the antibodies it contains.

It is recommended that nothing be given to children other than breastmilk even in the first three days when the milk has not begun to flow regularly because prelacteal feeds limit the frequency of suckling by the infant and expose the baby to the risk of infection. However, over one-fourth of children (26%) who were ever breastfed are given something other than breastmilk during the first three days. Overall, 89 percent of children continue breastfeeding at 1 year and more than two-thirds (71%) continue breastfeeding at 2 years. The median duration of breastfeeding is 25.7 months, which is the age to which half of children are breastfed.

After the first 6 months, breastmilk is no longer enough to meet the nutritional needs of infants. Therefore, complementary foods should be added to the diet of the child. At age 6-8 months less than two-thirds of children (56%) in Telangana receive breastmilk and complementary foods.

WHO has several recommendations for infant and young child feeding (IYCF) practices for children age 6-23 months. The key IYCF indicators measure the adequacy of dietary diversity and meal frequency for breastfed and nonbreastfed children. Less than one-third (30%) of children age 6-23 months are fed the recommended minimum number of times per day and even fewer (27%) are fed from the appropriate number of food groups. Only 10 percent are fed according to all three recommended practices.

Micronutrient deficiency is a major contributor to childhood morbidity and mortality. Vitamin A is an essential nutrient for the immune system. Severe vitamin A deficiency (VAD) can cause eye damage and a higher risk of dying from measles and diarrhoeal disease. The Government of India recommends that children under five years of age receive vitamin A supplements every six months, starting at age 9 months. In Telangana, about three-fourths (76%) of children age 9-59 months were given a vitamin A supplement in the past six months, but less than half (48%) of children age 9-23 months ate vitamin A-rich foods during the day or night before the survey.

Iron deficiency is a primary cause of anaemia. Eating foods rich in iron and taking iron supplements can help prevent anaemia. Only 31 percent of children age 9-23 months ate iron-rich foods during the day or night before the survey, and 38 percent of children age 6-59 months were given iron supplements in the week before the survey.


Children's nutritional status

Twenty-eight percent of children under age five years are stunted, or too short for their age, which indicates that they have been undernourished for some time. Eighteen percent are wasted, or too thin for their height, which may result from inadequate recent food intake or a recent illness causing weight loss, and 5 percent are severely wasted. Twenty-nine percent are underweight, which takes into account both chronic and acute undernutrition. One percent of children are overweight. Even during the first six months of life, when almost all babies are breastfed, 12 percent of children are stunted, 20 percent are underweight, and 40 percent are wasted.

There are pronounced differences in children's nutritional status by background characteristics. All three indicators of undernutrition are much higher in rural than urban areas, decline sharply with mothers' schooling, and are lower for first births than births at other birth orders. A higher proportion of Christian children are undernourished than Muslim children or Hindu children, and a higher proportion of scheduled caste children and scheduled tribe children are undernourished than children not belonging to a scheduled caste, scheduled tribe, or other backward class.

Children's Nutritional Status

Percentage of children under five years



Note: Nutritional status estimates are based on the 2006 WHO International Reference Population

Adults' nutritional status


Over half (51%) of women and 46 percent of men in Telangana are malnourished, that is, they are either too thin or are overweight or obese. More men and women are overweight or obese than thin: 23 percent of women and 21 percent of men in Telangana are too thin, and 28 percent of women and 24 percent of men are overweight or obese.

Undernutrition among adults is particularly common in the younger age groups, in rural areas, and among scheduled castes and scheduled tribes. Overweight and obesity are most prevalent among older adults and in the urban population.

The use of iodized salt prevents iodine deficiency, which can lead to miscarriage, goitre, and mental retardation. Almost all (96%) of Telangana's households were using iodized salt at the time of the survey. Mahbubnagar is the only district where less than 9 out of 10 households use iodized salt.

How many women are at a healthy weight for their height?

Percent distribution of women




Anaemia

Anaemia is a condition that is marked by low levels of haemoglobin in the blood. Iron deficiency is estimated to be responsible for about half of all anaemia globally, but anaemia can also be caused by malaria, hookworm and other helminths, other nutritional deficiencies, chronic infections, and genetic conditions. Anaemia can result in maternal mortality, weakness, diminished physical and mental capacity, increased morbidity from infectious diseases, perinatal mortality, premature delivery, low birth weight, and (in children) impaired cognitive performance, motor development, and scholastic achievement. Anaemia is a major health problem in Telangana, especially among women and children.

A majority of children age 6-59 months (61%) are anaemic. This includes 25 percent who are mildly anaemic, 34 percent who are moderately anaemic, and 3 percent who suffer from severe anaemia. Boys (62%) are slightly more likely than girls (59%) to have anaemia. Children whose mothers are anaemic have a much higher prevalence of anaemia (65-73%) than children whose mothers are not anaemic (54%). Although anaemia levels vary according to most background characteristics, at least half the children in every subgroup are anaemic, except children age 48-59 months (44%).

Anaemia among women, men, and children



Fifty-seven percent of women in Telangana have anaemia, including 37 percent with mild anaemia, 18 percent with moderate anaemia, and 3 percent with severe anaemia. Anaemia is particularly high for scheduled tribe women and Christian women, but half of women or more are anaemic in every subgroup of women except those not belonging to a scheduled caste, scheduled tribe or other backward class. However, even among women not belonging to a scheduled caste, scheduled tribe, and other backward class, 47 percent are anaemic. About one-sixth of men (15%) are anaemic. Men belonging to a scheduled tribe and men with no schooling are particularly likely to be anaemic.

ADULT HEALTH AND HEALTH CARE

Tuberculosis

In Telangana, 293 persons per 100,000 are estimated to have medically treated tuberculosis, based on reports from household respondents. The prevalence of medically treated tuberculosis is higher among men (412) than among women (173) and is higher in rural areas (413) than in urban areas (151).

The majority of respondents have heard of tuberculosis (72% of women and 73% of men), but even among those who have heard of tuberculosis, only 74 percent of women and 64 percent of men know that it is spread through the air by coughing or sneezing. Less than half of women and men have misconceptions about how tuberculosis is spread. Overall, more than 80 percent of women and men know that tuberculosis can be cured; however, 42 percent of women and 59 percent of men say that if a family member had tuberculosis, they would want to keep it a secret.

Diabetes, asthma, goitre, heart disease, and cancer

According to self-reports, 2,172 women age 15-49 and 2,137 men age 15-49 per 100,000 have diabetes. The prevalence of diabetes is particularly high among older women and men. Overall, 3,455 women and 1,519 men per 100,000 suffer from asthma. The prevalence of asthma among women is highest for the oldest age group (age 35-49) but among men it is highest for the youngest age group (age 15-19). Goitre or any other thyroid disorder is much more common among women, at 4,448 per 100,000, than among men, at 655 per 100,000. The prevalence of any heart disease is quite similar among women (1,648 per 100,000) and men (1,473 per 100,000). Among the five diseases, cancer is the least common, with 78 women per 100,000 and 75 men per 100,000 reportedly suffering from cancer.

Blood pressure (hypertension)

Based on blood pressure measurement during the survey, 13 percent of women age 15-49 in Telangana have hypertension, including 7 percent with Stage 1 hypertension, 2 percent with Stage 2 hypertension, and 1 percent with Stage 3 hypertension. Over two-thirds (67%) of women have normal blood pressure, including 2 percent of women with normal blood pressure who are taking medicine to lower their blood pressure. The prevalence of hypertension among men age 15-49 is somewhat higher than among women. Twenty percent of men in Telangana have hypertension, including 12 percent with Stage 1 hypertension, 4

percent with Stage 2 hypertension, and 3 percent with Stage 3 hypertension. Almost half (48%) of men have normal blood pressure, including 1 percent of men with normal blood pressure who are taking medicine to lower their blood pressure. For both women and men, hypertension tends to increase sharply with age. About one-third (32%) of women age 45-49 have hypertension and more than one-third (36-38%) of men age 35-49 have hypertension.

Blood glucose

NFHS-4 included measurement of random blood glucose among women age 15-49 and men age 15-54. Three percent of women and 2 percent of men age 15-49 in Telangana have high blood glucose levels, and an additional 4 percent each of women and men have very high blood glucose levels. Prevalence of high blood glucose generally increases with age and is higher in urban than rural areas for both women and men.

Health examinations

In Telangana, 32 percent of women have ever undergone an examination of the cervix, 10 percent have ever undergone a breast examination, and 11 percent have ever undergone an examination of the oral cavity.

Tobacco and alcohol use

Twenty-eight percent of men, but only 3 percent of women, age 15-49 use some form of tobacco. Men are much more likely to smoke cigarettes (14%) or *bidis* (7%) than to use other types of tobacco. Among men, the use of any form of tobacco is higher in rural areas than in urban areas. In urban areas, cigarettes are more popular than *bidis*, but in rural areas cigarettes and *bidis* are almost equally popular. Among men who smoke cigarettes or *bidis*, 37 percent smoked 10 or more cigarettes or *bidis* in the past 24 hours. Men are more likely to drink alcohol (54%) than to use tobacco, and 9 percent of women say that they drink alcohol. Among women and men who drink alcohol, almost three-fifths drink alcohol at least once a week.

Source of health care

The private health sector is the main source of health care for 72 percent of urban households and 64 percent of rural households. Household members are much more likely to go to private hospitals (42%) than to private doctors or clinics (18%).

Health insurance

Two-thirds of households in Telangana have some kind of health insurance that covers at least one usual member of the household. Health insurance coverage is more common in rural areas (77%) than in urban areas (54%). In Telangana, the most common health insurance programme is the state health insurance scheme. Other health insurance programmes used by at least 1 percent of households are the Employees' State Insurance Scheme (ESIS), the Central Government Health Scheme (CGHS), and the *Rashtriya Swasthya Bima Yojana* (RSBY).

Sixty-one percent of women and 71 percent of men age 15-49 years in Telangana are covered by any health scheme or health insurance. More women and men in rural areas than in urban areas are covered by any health scheme or health insurance. Health scheme or health

insurance coverage is higher among women and men with little or no schooling than among those with more years of schooling.

HIV/AIDS


Awareness of HIV or AIDS

Ninety percent of women in Telangana have heard of HIV or AIDS. In urban areas, 95 percent have heard about HIV or AIDS, compared with 86 percent in rural areas. Knowledge of HIV or AIDS declines from a high of 95 percent for women age 15-24 years to 83 percent for women age 40-49. Women with no schooling (78%) and women belonging to scheduled tribes (82%) are less likely than other women to have heard of HIV or AIDS.

Men are much more likely than women to know about HIV or AIDS. Almost all men (96%) in Telangana have heard of HIV or AIDS in both urban and rural areas.

Knowledge of prevention and transmission

Men are much more likely than women to know how HIV is transmitted and how to keep from getting it. For example, only 59 percent of women know that consistent condom use can help prevent HIV/AIDS, compared with 81 percent of men; and 73 percent of women know that having just one uninfected partner who has no other partners can reduce the chance of getting HIV/AIDS, compared with 80 percent of men.



HIV-related stigma

About two-thirds of women (65%) and men (67%) in Telangana would be willing to take care of a relative with HIV/AIDS in their home and similar proportions of women (62%) and men (65%) say that they would buy fresh vegetables from a shopkeeper with HIV/AIDS. A somewhat higher proportion of women and men (71% each) say that a female teacher who has

HIV/AIDS but is not sick should be allowed to continue teaching. Thirty-seven percent of women and 29 percent of men say that if a family member got infected with HIV/AIDS, they would not want to keep it a secret. Overall, only 15 percent of women and 11 percent of men express accepting attitudes on all four of these indicators.

HIV testing prior to NFHS-4, blood transfusions, and safe injections

Only 32 percent of women and 13 percent of men age 15-49 had ever been tested for HIV prior to NFHS-4. Women in urban areas are somewhat more likely to have ever been tested for HIV prior to NFHS-4 than rural women. Forty percent of women who had a live birth in the past five years and received ANC during pregnancy were tested for HIV during ANC. Urban women are more likely than rural women to have been tested for HIV during ANC (47% versus 34%).

Forty percent of women who had a live birth in the past five years and received ANC during pregnancy were tested for HIV during ANC.

In Telangana, similar proportions of men (12%) and women (10%) have ever had a blood transfusion. Women are more likely than men to have received an injection in the past 12 months (61% of women, compared with 54% of men).

For two-thirds of adults who received an injection in the past 12 months, a disposable syringe was used.

SEXUAL BEHAVIOUR

NFHS-4 included questions on respondents' sexual behaviour. Respondents were asked about their age at first sex, their current and previous sexual partners, higher-risk intercourse, and condom use. In addition, men were asked whether they had paid for sex in the past year. These questions are sensitive and subject to reporting bias, so the results should be interpreted with caution.

Higher-risk behaviour

Higher-risk sex is sexual intercourse with someone who is neither a spouse nor a cohabiting partner. Among those who had sex in the past 12 months, only 1 percent of women and 6 percent of men reported having had higher-risk sex during the past 12 months. The percentage of men who had higher-risk sex in the past 12 months is slightly higher in rural areas than in urban areas. Similarly low percentages of respondents said that they had multiple sex partners in the past 12 months (1% of women and 4% of men).

Less than one-fourth (24%) of men who had higher-risk sex in the past 12 months reported using a condom the last time they had higher-risk sex.

One percent of men said they had paid for sex in the past year.

WOMEN'S EMPOWERMENT

Women's hygiene

Using a hygienic method of menstrual protection is important for women's health and personal hygiene. In NFHS-4, young women age 15-24 were asked what method they use for menstrual protection, if anything. In Telangana, 60 percent of women age 15-24 use sanitary napkins, 33 percent use cloth, 18 percent use locally prepared napkins, and 3 percent use tampons. Overall, 76 percent of women age 15-24 use a hygienic method of menstrual protection. Women with at least 10 years of schooling are more than twice as likely to be using a hygienic method as women with no schooling or less than five years of schooling.

Employment and earnings

Only 47 percent of all women age 15-49 were employed in the 12 months preceding the survey; in the same period, 81 percent of all men age 15-49 were employed. Among employed women, 95 percent earned cash, including 2 percent whose earnings were in both cash and kind, and 3 percent were not paid at all. Similarly, most men (96%) who were employed earned cash and 3 percent were not paid at all. Forty-six percent of employed women worked in a non-agricultural occupation, compared with 68 percent of employed men.

Only 47 percent of all women age 15-49 were employed in the 12 months preceding the survey.

Among currently married women who work and are paid in cash, 74 percent decide how their earnings will be used, either alone or jointly with their husbands. Over two-fifths (41%) of women who work for cash say that they earn more or about the same as their husbands. By contrast, 86 percent of currently married men whose wives have cash earnings report that they alone or jointly with their wives decide how her earnings are used and 52 percent of men who have cash earnings and whose wives have cash earnings say that their wife earns more or about the same as them.

Decision making

Currently married women were asked who makes decisions about their own health care, major household purchases, and visits to their own family or relatives. Women are about equally likely to participate alone or jointly with their husband in all of the three decisions: 68 percent participate in decisions about their own health care and 71-72 percent participate in decisions about major household purchases and visits to their own family and relatives.

Overall, 57 percent of currently married women participate in making all three of these decisions, and 19 percent do not participate in making any of the three decisions. Participation in all three decisions varies most by age, increasing sharply from 26 percent among women age 15-19 to 67 percent among women age 40-49.

Currently married men were also asked who makes decisions about their own health care and major household purchases. Men were much more likely than women to report that they alone or jointly with their wife participated in these decisions: 88 percent participated in making decisions about their own health care, 93 percent in making decisions about major household purchases, and 84 percent in making both these decisions.

Other indicators of women's empowerment

Only 24 percent of women have some money that they can decide how to use. The proportion of women with money which they control is higher among urban than rural women, increases sharply with age, and is higher for women who are employed for cash (31%) than for women who are not employed or employed but not for cash.

Three out of five women have a bank or savings account that they themselves use.

Three out of five women have a bank or savings account that they themselves use. This percentage is highest, at 68 percent, among women who are employed for cash. Sixty percent of women know of a microcredit programme in the area and 23 percent have ever taken a loan from a microcredit programme.

Only 45 percent of women are allowed to go by themselves to all three of the following places: the market, a health facility, and places outside the village/community. The groups of women in which higher proportions are allowed to go to all three places alone are women age 40-49, women with 3-4 children, and employed women who earn cash.

Ownership of assets

Forty-seven percent of women and 71 percent of men own a house alone or jointly with someone else, and 31 percent of women and 49 percent of men own land alone or jointly with someone else. Ownership of both a house and land is more common in rural than urban areas for both women and men.

Forty-seven percent of women own a house alone or jointly with someone else, and 31 percent of women own land alone or jointly with someone else.

Less than half (48%) of women have a mobile phone that they themselves use, and among women who have a mobile phone that they themselves use, two in three can read SMS messages. Sixty-three percent of urban women have a mobile phone they themselves use, compared with 34 percent of rural women. Having a mobile phone that women themselves use increases sharply with education. Only 26 percent of scheduled tribe women and 38 percent of scheduled caste women have a mobile phone that they themselves use, compared with 73 percent of women not belonging to a scheduled caste, scheduled tribe or other backward class.

Less than half (48%) of women have a mobile phone that they themselves use.

Gender-role attitudes

Eighty-four percent of women believe it is justifiable for a husband to beat his wife under some circumstances. Women are most likely to believe that wife beating is justified if a woman neglects the house or children (70%), followed by if she shows disrespect for her in-laws (66%) and if she argues with her husband (46%). Men are only somewhat less likely to agree: 75 percent say that wife beating is justified in some circumstances, especially if the wife neglects the house or children (57%) or the wife shows disrespect for in-laws (51%). Even among women and men who have completed at least 12 years of schooling, more than three in four (78% of women and 75% of men) say that a husband is justified in beating his wife for one or more of the specified reasons.

Sixty-two percent of women and 58 percent of men believe that a woman is justified in refusing to have sex with her husband for all of the following three reasons: if she knows he has a sexually transmitted disease, if she knows he has intercourse with other women, and if she is tired or not in the mood. However, about one in six (17%) women and one in eight (12%) men agree that a wife is not justified in refusing sex with her husband for any of the three reasons. A minority of men, 43 percent, agree that if a wife refuses to have sex with her husband he does not have the right to do even one of the following: get angry and reprimand her, refuse to give her financial support, use force to have sex even if she doesn't want to, or have sex with another woman.

DOMESTIC VIOLENCE

Among women age 15-49, 44 percent have ever experienced physical violence and 6 percent have ever experienced sexual violence. In all, 45 percent of women in Telangana have experienced physical or sexual violence.

Forty-five percent of women age 15-49 in Telangana have experienced physical or sexual violence.

Violence during pregnancy


Six percent of women who have ever been pregnant have ever experienced physical violence during one or more of their pregnancies. Widowed/divorced/separated/deserted women (23%), women belonging to the scheduled castes (14%), and those with no schooling (9%), are more likely than women in any other group to have experienced violence during pregnancy.

Spousal violence

Thirty-six percent of ever-married women report having been slapped by their husband, 21 percent report having been pushed, shaken, or having something thrown at them, 11-16 percent report being kicked, dragged, or beaten up; having their arm twisted or hair pulled; or being punched; and 1-2 percent each report being choked or burned on purpose or being threatened or attacked with a knife, gun, or any other weapon. Five percent report that their husbands have physically forced them to have sex even when they did not want to and 3 percent report that their husband forced them with threats or other ways to perform sexual acts they did not want to perform. Overall, 43 percent of ever-married women have experienced spousal physical or sexual violence from their current husband or, if not currently married, from their most recent husband, and 35 percent have experienced such violence in the past 12 months. Twenty percent report spousal emotional violence. Few ever-married women (5%) have ever initiated violence against their husband.

Is alcohol use related to spousal physical or sexual violence?

Percentage of ever-married women experiencing violence



Note: Category "Husband drinks, but never gets drunk" not shown since there are fewer than 25 unweighted cases.

Although the prevalence of spousal violence is lower among more educated women, more than 1 in 4 (27%) women who have at least 12 years of schooling have experienced physical or sexual spousal violence. The contextual and intergenerational aspects of spousal violence are clear from the fact that women whose mothers were beaten by their fathers are much more likely than women whose fathers did not beat their mothers to be in abusive marriages themselves. Women whose husbands consume alcohol are much more likely than women whose husbands do not consume alcohol to experience spousal violence, especially if

the husband often gets drunk. However, 29 percent of even women whose husbands do not drink alcohol have experienced physical or sexual spousal violence.

About one-fourth (23%) of women who have experienced spousal physical or sexual violence have suffered injuries as a result of the violence. The most common type of injuries are cuts, bruises, or aches.

Help seeking

Only 16 percent of women who have ever experienced physical or sexual violence by anyone have sought help. About three-fourths (74%) of women have neither sought help nor told anyone about the violence. Abused women who have sought help most often seek help from their own families. Only 2 percent of abused women who sought help for the violence sought help from the police.

Table 1 Results of the household and individual interviews

Number of households, number of interviews with women and men, and response rates, according to residence, Telangana, 2015-16

Result	Residence		Total
	Urban	Rural	
Household interviews			
Households selected	3,026	5,386	8,412
Households occupied	2,962	5,273	8,235
Households interviewed	2,739	5,047	7,786
Household response rate ¹	92.5	95.7	94.5
Interviews with women age 15-49			
Number of eligible women	3,168	5,071	8,239
Number of eligible women interviewed	2,775	4,792	7,567
Eligible women response rate ²	87.6	94.5	91.8
Interviews with men age 15-54			
Number of eligible men	557	814	1,371
Number of eligible men interviewed	410	723	1,133
Eligible men response rate ²	73.6	88.8	82.6

Note: Eligible women and men are women age 15-49 and men age 15-54 who stayed in the household the night before the household interview (including both usual residents and visitors). This table is based on the unweighted sample.

¹ Households interviewed/households occupied

² Respondents interviewed/eligible respondents

Table 2 Results of the household and individual interviews by district

Number of households, number of women and men interviewed, and response rates by residence and district, Telangana, 2015-16

District	Households interviewed			Household response rate ¹	Number of eligible women interviewed			Eligible women response rate ²	Number of eligible men interviewed			Eligible men response rate ²
	Urban	Rural	Total		Urban	Rural	Total		Urban	Rural	Total	
Adilabad	223	559	782	95.0	225	539	764	92.7	33	84	117	83.6
Hyderabad	766	0	766	92.0	830	0	830	88.2	118	0	118	73.8
Karimnagar	202	574	776	95.8	223	505	728	92.7	24	84	108	89.3
Khammam	187	619	806	96.2	169	609	778	93.8	29	86	115	85.8
Mahbubnagar	126	665	791	96.1	120	637	757	94.5	19	93	112	88.2
Medak	171	589	760	93.4	174	569	743	89.4	25	86	111	79.9
Nalgonda	147	654	801	96.5	136	587	723	95.1	21	97	118	83.7
Nizamabad	172	591	763	94.8	174	578	752	90.2	22	71	93	80.9
Rangareddy	527	230	757	91.3	520	242	762	89.1	79	44	123	75.5
Warangal	218	566	784	94.5	204	526	730	93.7	40	78	118	90.1
Telangana	2,739	5,047	7,786	94.5	2,775	4,792	7,567	91.8	410	723	1,133	82.6

Note: This table is based on the unweighted sample.

¹ Households interviewed/households occupied

² Respondents interviewed/eligible respondents

Table 3 Household population by age, schooling, residence, and sex

Percent distribution of the *de facto* household population by residence and sex, according to age and schooling; and the percentage of individuals who have an *Aadhaar* card, Telangana, 2015-16

Background characteristic	Urban			Rural			Total		
	Male	Female	Total	Male	Female	Total	Male	Female	Total
Age									
0-4	9.2	8.3	8.8	9.3	7.8	8.6	9.2	8.0	8.6
5-9	7.8	7.9	7.8	8.5	7.8	8.1	8.1	7.8	8.0
10-14	8.5	8.2	8.4	8.9	7.9	8.4	8.7	8.0	8.4
15-19	7.5	7.3	7.4	8.7	7.8	8.2	8.2	7.6	7.9
20-24	9.9	11.0	10.4	9.0	9.5	9.2	9.4	10.2	9.8
25-29	10.5	11.2	10.9	9.1	9.8	9.4	9.8	10.4	10.1
30-34	8.7	8.1	8.4	7.4	6.8	7.1	8.0	7.4	7.7
35-39	8.4	7.9	8.2	6.8	7.1	7.0	7.6	7.5	7.5
40-44	6.5	6.7	6.6	5.4	5.3	5.3	5.9	5.9	5.9
45-49	6.3	6.3	6.3	5.3	6.7	6.0	5.8	6.5	6.2
50-54	4.4	4.5	4.5	4.1	4.7	4.4	4.3	4.6	4.4
55-59	3.6	4.3	4.0	3.7	4.6	4.1	3.7	4.5	4.1
60-64	2.8	3.1	2.9	4.0	5.0	4.5	3.4	4.1	3.8
65-69	2.8	2.7	2.8	4.4	4.3	4.3	3.6	3.6	3.6
70-74	1.6	1.1	1.4	2.7	2.4	2.6	2.2	1.8	2.0
75-79	0.7	0.5	0.6	1.3	1.1	1.2	1.0	0.9	1.0
80 +	0.7	0.8	0.7	1.3	1.5	1.4	1.0	1.2	1.1
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Aadhaar card									
Percentage with an <i>Aadhaar</i> card	93.6	94.3	94.0	96.3	96.5	96.4	95.0	95.5	95.3
Number	6,942	6,776	13,719	7,895	8,169	16,064	14,837	14,945	29,782
Sex ratio, all ages ¹	na	na	976	na	na	1,035	na	na	1,007
Sex ratio, age 0-6 years ¹	na	na	923	na	na	894	na	na	907
Schooling²									
No schooling	11.5	23.3	17.4	28.3	49.5	39.2	20.5	37.7	29.2
<5 years complete	12.3	12.0	12.2	17.1	13.4	15.2	14.8	12.8	13.8
5-9 years complete	22.8	23.7	23.3	24.6	18.9	21.7	23.7	21.1	22.4
10-11 years complete	17.7	15.7	16.7	14.1	9.9	11.9	15.8	12.5	14.1
12 or more years complete	35.4	25.2	30.3	15.8	8.2	11.9	25.0	15.9	20.4
Don't know	0.3	0.1	0.2	0.1	0.1	0.1	0.2	0.1	0.1
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Number	6,186	6,083	12,269	7,022	7,398	14,420	13,207	13,481	26,688
Median number of years of schooling completed	9.2	6.8	8.5	4.7	0	2.7	6.9	3.8	5.5

na = Not applicable

¹ Females per 1,000 males

² Population age 6 and above

Table 4 Household and housing characteristics

Percent distribution of urban, rural, and total households and *de jure* population by household and housing characteristics, Telangana, 2015-16

Household and housing characteristic	Urban	Rural	Total	<i>De jure</i> population
Household headship				
Male	87.1	82.9	84.8	88.4
Female	12.9	17.1	15.2	11.6
Total	100.0	100.0	100.0	100.0
Mean household size	4.0	3.8	3.9	na
Household structure¹				
Nuclear	69.4	65.0	66.9	56.6
Non-nuclear	30.6	35.0	33.1	43.4
Total	100.0	100.0	100.0	100.0
Religion of household head				
Hindu	80.7	92.8	87.4	85.6
Muslim	16.0	3.9	9.3	11.2
Christian	3.1	3.1	3.1	3.1
Other	0.2	0.1	0.2	0.1
Total	100.0	100.0	100.0	100.0
Caste/tribe of household head				
Scheduled caste	13.1	23.2	18.7	18.6
Scheduled tribe	3.2	11.4	7.7	7.8
Other backward class	58.0	55.6	56.7	57.1
Other	24.7	9.0	16.0	15.8
Don't know	0.9	0.8	0.9	0.7
Total	100.0	100.0	100.0	100.0
Electricity				
Yes	99.5	97.2	98.2	98.9
No	0.5	2.8	1.8	1.1
Total	100.0	100.0	100.0	100.0
Source of drinking water				
Improved source	80.1	75.6	77.6	78.1
Piped water into dwelling/yard/plot	60.0	24.9	40.5	41.7
Public tap/standpipe	17.7	28.7	23.8	23.4
Tube well or borehole	1.5	18.7	11.1	11.0
Other improved ²	0.8	3.3	2.2	2.1
Unimproved source ³	19.8	24.3	22.3	21.9
Other source	0.1	0.1	0.1	0.0
Total	100.0	100.0	100.0	100.0
Time to obtain drinking water (round trip)				
Water on premises/delivered to dwelling	78.5	41.7	58.1	58.8
Less than 30 minutes	17.8	45.9	33.4	32.8
Thirty minutes or longer	3.4	12.4	8.4	8.3
Don't know	0.3	0.0	0.1	0.1
Total	100.0	100.0	100.0	100.0

Continued...

Table 4 Household and housing characteristics—Continued

Percent distribution of urban, rural and total households and *de jure* population by household and housing characteristics, Telangana, 2015-16

Household and housing characteristic	Urban	Rural	Total	<i>De jure</i> population
Water treatment prior to drinking⁴				
Boil	4.9	1.2	2.9	2.8
Strain through cloth	4.7	8.4	6.8	6.7
Use ceramic, sand, or other water filter	20.5	1.3	9.8	10.1
Electronic purifier	6.3	0.6	3.1	3.2
Other treatment	4.4	3.7	4.0	4.0
No treatment	60.7	85.3	74.4	74.3
Percentage using an appropriate treatment method ⁵	31.0	3.5	15.7	16.0
Sanitation facility				
Improved, not shared facility	64.4	38.9	50.2	51.7
Flush/pour flush to piped sewer system, septic tank, or pit latrine	63.5	37.4	49.0	50.3
Pit latrine with slab	0.9	0.8	0.8	0.9
Other ⁶	0.1	0.7	0.4	0.5
Shared facility ⁷	13.1	8.2	10.4	9.6
Flush/pour flush to piped sewer system, septic tank, or pit latrine	12.9	8.1	10.2	9.4
Pit latrine with slab	0.2	0.0	0.1	0.1
Other ⁶	0.0	0.1	0.1	0.1
Unimproved	22.5	52.9	39.4	38.7
Flush/pour flush not to piped sewer system, septic tank, or pit latrine	15.1	0.8	7.2	7.3
Pit latrine without slab/open pit	0.2	0.9	0.6	0.6
Other unimproved facility ⁸	0.1	0.1	0.1	0.1
No facility/open space/field	6.9	51.0	31.4	30.6
Other	0.2	0.1	0.2	0.2
Total	100.0	100.0	100.0	100.0
Type of house⁹				
<i>Kachha</i>	0.4	4.2	2.5	2.2
<i>Semi-pucca</i>	7.2	34.7	22.5	21.9
<i>Pucca</i>	92.1	60.8	74.7	75.6
Missing	0.2	0.3	0.3	0.3
Total	100.0	100.0	100.0	100.0
Cooking fuel				
Electricity	0.5	0.1	0.3	0.3
LPG/natural gas	89.5	47.6	66.3	67.4
Biogas	0.1	0.4	0.3	0.3
Kerosene	2.5	0.7	1.5	1.4
Coal/lignite	0.1	0.1	0.1	0.1
Charcoal	0.2	0.9	0.6	0.5
Wood	6.8	48.4	29.9	29.2
Straw/shrubs/grass	0.0	0.7	0.4	0.4
Agricultural crop waste	0.0	0.4	0.2	0.2
Dung cakes	0.0	0.2	0.1	0.1
No food cooked in the household	0.2	0.4	0.3	0.1
Other	0.1	0.0	0.0	0.0
Total	100.0	100.0	100.0	100.0
Percentage using clean fuel for cooking ¹⁰	90.1	48.2	66.8	68.0
Percentage using solid fuel for cooking ¹¹	7.1	50.7	31.3	30.5

Continued...

Table 4 Household and housing characteristics—Continued

Percent distribution of urban, rural and total households and *de jure* population by household and housing characteristics, Telangana, 2015-16

Household and housing characteristic	Urban	Rural	Total	<i>De jure</i> population
Place for cooking				
In the house, separate room	53.9	30.8	41.1	43.2
In the house, no separate room	35.3	38.9	37.3	36.1
In a separate building	5.5	4.8	5.1	5.2
Outdoors	5.1	25.1	16.2	15.4
No food cooked in household	0.2	0.4	0.3	0.1
Total	100.0	100.0	100.0	100.0
Number	3,465	4,321	7,786	30,340
Type of fire/stove among households using solid fuels¹¹				
Stove	0.9	1.3	1.2	1.2
<i>Chullah</i>	91.5	85.2	85.8	86.6
Open fire	7.6	13.5	12.9	12.2
Other	0.0	0.1	0.1	0.0
Total	100.0	100.0	100.0	100.0
Number using solid fuel	247	2,190	2,437	9,242
Frequency of smoking in the house¹²				
Daily	14.3	30.1	23.1	24.7
Weekly	7.9	7.4	7.6	7.7
Monthly	1.2	1.1	1.2	1.1
Less than monthly	1.9	1.7	1.8	1.7
Never	74.7	59.8	66.4	64.7
Total	100.0	100.0	100.0	100.0
Number	3,465	4,321	7,786	30,340

na = Not applicable

¹ Nuclear households are households comprised of a married couple or a man or a woman living alone or with unmarried children (biological, adopted, or fostered) with or without unrelated individuals.

² Protected dug well, protected spring, rainwater, community RO plant

³ Surface water, unprotected dug well, unprotected spring, cart with small tank, tanker truck, bottled water

⁴ Total may add to more than 100.0 because households may use more than one method of purification

⁵ Appropriate water treatment methods include boiling, bleaching, filtering, and electronic purifying

⁶ Includes ventilated improved pit (VIP)/biogas latrine and twin pit/composting toilet

⁷ Facilities that would be considered improved if they were not shared by two or more households

⁸ Includes dry toilet

⁹ Houses made from mud, thatch, or other low-quality materials are called *kachha* houses, houses that use partly low-quality and partly high-quality materials are called semi-*pucca* houses, and houses made with high quality materials throughout, including the floor, roof, and exterior walls, are called *pucca* houses.

¹⁰ Electricity, LPG/natural gas, or biogas

¹¹ Includes coal/lignite, charcoal, wood, straw/shrubs/grass, agricultural crop waste, and dung cakes

¹² Frequency of smoking by anyone inside the house

Table 5 Household possessions and land ownership

Percentage of urban, rural, and total households and *de jure* population possessing various household goods, means of transport, agricultural land, a house and farm animals and having a bank/post office account, health scheme/health insurance, a BPL card, and a long-lasting insecticide-treated (LLIN) mosquito net, Telangana, 2015-16

Household possessions	Urban	Rural	Total	<i>De jure</i> population
Household goods				
Mattress	78.0	44.6	59.5	61.8
Pressure cooker	67.9	22.9	42.9	45.2
Chair	94.0	81.7	87.2	89.4
Cot or bed	87.0	80.3	83.3	84.0
Table	69.3	43.3	54.9	57.1
Electric fan	97.2	89.1	92.7	94.4
Radio or transistor	5.3	1.5	3.2	3.3
Television (black and white)	2.0	2.8	2.4	2.4
Television (colour)	89.4	68.1	77.6	81.8
Any television	90.4	70.5	79.4	83.5
Sewing machine	22.5	10.6	15.9	17.8
Mobile telephone	96.1	85.5	90.2	93.8
Landline telephone	8.0	1.0	4.1	4.3
Internet	12.9	0.7	6.1	6.4
Computer	15.9	1.3	7.8	8.3
Refrigerator	49.4	9.6	27.3	29.1
Air conditioner/cooler	38.6	9.1	22.2	23.6
Washing machine	24.8	1.3	11.8	12.4
Watch or clock	85.2	62.4	72.6	75.6
Water pump	41.0	27.4	33.4	33.8
Thresher	1.1	0.9	1.0	1.0
Tractor	0.7	1.6	1.2	1.5
None of the above	0.3	1.2	0.8	0.4
Means of transport				
Bicycle	23.5	30.0	27.2	30.0
Motorcycle or scooter	54.5	26.5	38.9	43.0
Animal-drawn cart	0.5	7.6	4.4	5.0
Car	9.9	1.2	5.1	5.6
None of the above	34.3	48.6	42.3	37.4
Agricultural land				
No agricultural land	87.3	44.5	63.5	62.4
Agricultural land	12.7	55.5	36.5	37.6
Irrigated land only	5.5	15.7	11.2	11.3
Non-irrigated land only	5.2	27.7	17.7	18.1
Both irrigated and non-irrigated land	1.3	11.5	7.0	7.6
Irrigation not determined	0.7	0.6	0.6	0.6
Total	100.0	100.0	100.0	100.0
Percentage owning a house	47.7	63.6	56.5	56.7
Percentage owning farm animals ¹	4.5	39.2	23.8	26.1
Percentage having a bank account/post office account ²	91.3	91.7	91.5	92.4
Percentage covered by a health scheme/health insurance ³	53.5	76.7	66.4	66.8
Percentage having a BPL card	68.8	92.1	81.7	82.4
Percentage with an LLIN mosquito net	0.1	0.3	0.2	0.2
Number	3,465	4,321	7,786	30,340

BPL = Below poverty line

¹ Cows, bulls, buffaloes, camels, horses, donkeys, mules, goats, sheep, chickens, or ducks

² Percentage of households in which any usual member of the household has a bank account/post office account

³ Percentage of households in which any usual member of the household is covered by a health scheme/health insurance

Table 6 School attendance

Percentage of *de facto* household population age 6-17 years attending school in the 2014-15 school year by sex and residence, according to selected background characteristics, Telangana, 2015-16

Background characteristic	Male			Female			Total		
	Urban	Rural	Total	Urban	Rural	Total	Urban	Rural	Total
Age									
6-10 (Primary)	97.7	98.1	98.0	99.3	98.0	98.6	98.5	98.1	98.2
6-13 (Elementary)	96.1	97.7	97.0	97.7	96.0	96.8	96.9	96.8	96.9
11-13 (Upper primary)	93.3	96.9	95.3	95.2	92.9	94.0	94.3	94.9	94.6
14-15 (Secondary)	83.5	88.1	86.2	89.3	80.0	84.2	86.3	84.5	85.3
16-17 (Higher secondary)	76.7	65.8	71.1	72.2	54.3	61.4	74.8	59.8	66.4
11-14 years	94.6	97.0	95.9	96.8	94.2	95.4	95.7	95.7	95.7
15-17 years	82.3	79.6	80.7	82.7	71.4	76.0	82.5	75.5	78.4
6-14 years	95.2	97.0	96.2	96.8	95.0	95.8	96.0	96.1	96.0
6-17 years	90.6	90.9	90.7	92.4	85.6	88.6	91.4	88.3	89.7
Religion									
Hindu	92.6	91.1	91.7	93.9	86.1	89.1	93.2	88.7	90.5
Muslim	84.0	87.8	84.8	87.5	72.5	84.6	85.7	80.8	84.7
Christian	(93.3)	90.3	91.6	(93.8)	88.6	90.9	93.6	89.5	91.2
Caste/tribe									
Scheduled caste	89.8	90.8	90.5	94.4	85.3	88.0	92.1	88.2	89.3
Scheduled tribe	(86.7)	86.2	86.3	(82.0)	81.2	81.3	84.3	83.9	84.0
Other backward class	91.8	92.1	92.0	93.6	86.4	89.8	92.7	89.4	90.9
Other	89.7	90.4	89.9	89.6	88.8	89.4	89.7	89.6	89.6
Don't know	*	*	*	*	*	*	*	(87.8)	(79.3)

Note: In this table, children's age refers to their age at the start of the 2014-15 school year (assumed here to be April 2014).

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

Table 7 Children's living arrangements and orphanhood

Percent distribution of *de jure* children under age 18 by their living arrangements, and percentage of children with one or both biological parents dead, according to background characteristics, Telangana, 2015-16

Background characteristic	Living with both parents	Living with mother but not with father	Living with father but not with mother	Not living with either parent	Total	Percentage with one or both parents dead ¹	Number of children
Age							
<5 years	94.4	3.9	0.6	1.1	100.0	1.4	2,577
5-9 years	90.5	6.2	0.8	2.4	100.0	5.0	2,411
10-14 years	85.2	9.5	2.1	3.2	100.0	9.5	2,540
15-17 years	81.5	10.7	2.2	5.5	100.0	10.3	1,345
Residence							
Urban	90.9	6.2	0.7	2.2	100.0	4.9	4,042
Rural	87.0	8.0	1.9	3.1	100.0	7.0	4,833
Sex							
Male	89.2	7.0	1.4	2.4	100.0	5.9	4,633
Female	88.4	7.3	1.2	3.1	100.0	6.2	4,242
Total age <15 years	90.1	6.5	1.2	2.2	100.0	5.3	7,529
Total age <18 years	88.8	7.2	1.3	2.7	100.0	6.0	8,874

¹ Includes children with father dead, mother dead, both parents dead, and one parent dead but missing information on survival status of the other parent

Table 8 Birth registration of children under age five

Percentage of *de jure* children under age five years whose birth was registered with the civil authorities, according to background characteristics, Telangana, 2015-16

Background characteristic	Percentage of children whose birth was registered		Total registered	<i>De jure</i> children
	Registered, has a birth certificate	Registered, does not have a birth certificate		
Age				
<2 years	70.8	13.5	84.3	982
2-4 years	72.5	9.5	82.0	1,596
Sex				
Male	70.2	11.3	81.5	1,362
Female	73.7	10.7	84.4	1,215
Residence				
Urban	81.0	8.9	89.9	1,221
Rural	63.6	12.9	76.5	1,356
Total	71.9	11.0	82.9	2,577

Table 9 Birth registration of children under age five by district

Percentage of *de jure* children under age five years whose birth was registered with the civil authorities, by district, Telangana, 2015-16

District	Percentage of children whose birth was registered		Total registered	<i>De jure</i> children
	Registered, has a birth certificate	Registered, does not have a birth certificate		
Adilabad	54.6	8.0	62.5	190
Hyderabad	83.1	9.8	92.9	417
Karimnagar	80.2	11.7	91.9	230
Khammam	70.2	10.4	80.6	187
Mahbubnagar	53.3	16.0	69.3	284
Medak	77.1	13.9	91.0	201
Nalgonda	69.1	12.7	81.8	239
Nizamabad	73.0	13.4	86.4	172
Rangareddy	77.4	7.8	85.2	450
Warangal	67.6	9.4	77.0	207
Telangana	71.9	11.0	82.9	2,577

Table 10 Background characteristics of respondents

Percent distribution of women and men age 15-49 by selected background characteristics and percentage exposed to various media, Telangana, 2015-16

Background characteristic	Weighted percent		Number of women		Number of men	
	Women	Men	Weighted	Unweighted	Weighted	Unweighted
Age						
15-19	13.2	16.5	997	1,053	174	180
20-24	17.9	14.6	1,355	1,361	155	155
25-29	18.9	17.4	1,428	1,391	184	188
30-34	13.8	14.7	1,042	1,025	155	148
35-39	13.5	13.3	1,019	1,013	140	135
40-44	11.1	11.0	836	827	116	125
45-49	11.8	12.5	890	897	132	123
Residence						
Urban	47.8	47.8	3,619	2,775	506	380
Rural	52.2	52.2	3,948	4,792	552	674
Schooling						
No schooling	32.5	17.0	2,458	2,757	179	198
<5 years complete	4.5	4.2	337	371	44	53
5-9 years complete	19.7	24.3	1,493	1,512	257	266
10-11 years complete	19.0	23.2	1,435	1,370	245	243
12 or more years complete	24.4	31.4	1,844	1,557	333	294
Literacy						
Literate ¹	65.2	83.4	4,936	4,628	882	853
Not literate	33.6	16.3	2,542	2,836	173	196
Not measured	1.2	0.3	89	103	3	5
Percentage exposed to various media						
Reads a newspaper/magazine at least once a week	33.1	61.2	2,503	2,168	647	623
Watches television at least once a week	88.7	92.2	6,712	6,601	975	961
Listens to the radio at least once a week	10.6	23.0	800	774	244	220
Visits the cinema/theatre at least once a month	16.3	49.6	1,232	992	525	499
Not regularly exposed to any media	9.0	4.1	684	794	44	51
Marital status						
Never married	18.4	35.8	1,390	1,386	379	370
Currently married	74.3	62.9	5,619	5,603	665	670
Married, <i>gauna</i> not performed	0.0	0.1	2	2	1	1
Widowed	5.7	0.4	428	437	4	5
Divorced/separated/deserted	1.7	0.8	128	139	9	8
Religion						
Hindu	86.3	86.0	6,529	6,493	910	906
Muslim	10.2	11.4	774	818	120	118
Christian	3.3	2.6	248	238	27	30
Other	0.2	0.0	15	18	0	0

Continued...

Table 10 Background characteristics of respondents—Continued

Percent distribution of women and men age 15-49 by selected background characteristics and percentage exposed to various media, Telangana, 2015-16

Background characteristic	Weighted percent		Number of women		Number of men	
	Women	Men	Weighted	Unweighted	Weighted	Unweighted
Caste/tribe						
Scheduled caste	18.4	15.1	1,395	1,483	160	174
Scheduled tribe	7.9	8.2	597	717	87	106
Other backward class	58.1	63.0	4,393	4,324	666	647
Other	15.0	13.7	1,137	988	145	127
Don't know	0.6	0.0	44	55	0	0
Total age 15-49	100.0	100.0	7,567	7,567	1,058	1,054
Age 50-54	na	6.6	na	na	75	79
Total age 15-54	na	100	na	na	1,133	1,133
Employment status (past 12 months)						
Employed at any time	47.4	81.4	610	667	861	863
In agricultural occupation	25.8	26.0	332	401	275	333
In non-agricultural occupation	21.1	51.5	272	258	545	490
Don't know	0.5	3.9	6	8	41	40
Not employed	52.6	18.6	675	635	197	191
Total age 15-49	100.0	100.0	1,285	1,302	1,058	1,054
Age 50-54	na	6.6	na	na	75	79
Total age 15-54	na	100	na	na	1,133	1,133

na = Not applicable

¹ Refers to women/men who can read a whole sentence or part of a sentence and women/men who completed standard 6 or higher (who are assumed to be literate)

Table 11 Current fertility

Age-specific and total fertility rates and crude birth rates, by residence, Telangana 2015-16

Age	Urban	Rural	Total
15-19	0.045	0.088	0.068
20-24	0.173	0.180	0.177
25-29	0.094	0.083	0.088
30-34	0.016	0.018	0.017
35-39	0.005	0.004	0.004
40-44	0.002	0.001	0.002
45-49	0.000	0.003	0.002
TFR (15-49)	1.67	1.88	1.79
CBR	17.1	17.2	17.1

Note: Rates are for the period 1-36 months preceding the survey (approximately 2013-15). Age-specific fertility rates are expressed per woman. Rates for the age group 45-49 might be slightly biased due to truncation.

TFR = Total fertility rate, expressed per woman

CBR = Crude birth rate, expressed per 1,000 population

Table 12 Fertility by background characteristics

Total fertility rate for the three years preceding the survey, percentage of women age 15-49 currently pregnant, mean number of children ever born to women age 40-49, and total wanted fertility rate, by background characteristics, Telangana, 2015-16

Background characteristic	Total fertility rate	Percentage of women age 15-49 currently pregnant	Mean number of children ever born to women age 40-49	Total wanted fertility rate
Residence				
Urban	1.67	3.3	2.7	1.53
Rural	1.88	3.1	2.9	1.64
Schooling				
No schooling	2.23	1.4	3.0	1.85
<5 years complete	2.05	1.7	2.9	1.64
5-9 years complete	1.90	3.5	2.6	1.70
10-11 years complete	1.87	4.7	2.4	1.67
12 or more years complete	1.83	4.4	2.0	1.70
Religion				
Hindu	1.79	3.1	2.8	1.61
Muslim	1.87	4.0	3.4	1.52
Christian	1.65	3.5	3.1	1.52
Caste/tribe				
Scheduled caste	1.72	3.0	2.9	1.51
Scheduled tribe	1.79	3.8	3.4	1.51
Other backward class	1.77	3.3	2.8	1.58
Other	1.93	2.7	2.6	1.80
Don't know	(2.44)	0.0	*	(2.11)
Total	1.79	3.2	2.8	1.59

() Based on 125-249 unweighted woman-years of exposure

* Based on fewer than 25 unweighted cases

Table 13 Teenage pregnancy and motherhood

Percentage of women age 15-19 who have had a live birth or who are pregnant with their first child, and percentage who have begun childbearing, by background characteristics, Telangana, 2015-16

Background characteristic	Percentage of women age 15-19 who:		Percentage of women age 15-19 who have begun childbearing	Number of women
	Have had a live birth	Are pregnant with first child		
Age				
15	0.6	0.0	0.6	156
16	0.0	0.9	0.9	179
17	2.0	2.8	4.8	192
18	10.1	4.1	14.2	247
19	21.0	5.3	26.3	223
Residence				
Urban	5.4	1.1	6.5	442
Rural	9.5	4.3	13.8	555
Schooling				
No schooling	(20.5)	(12.1)	(32.6)	36
5-9 years complete	9.9	3.1	13.0	263
10-11 years complete	6.5	2.4	8.9	382
12 or more years complete	5.1	2.4	7.5	303
Marital status				
Never married	0.0	0.0	0.0	816
Currently married	42.0	16.0	58.0	180
Religion				
Hindu	7.8	2.9	10.7	832
Muslim	5.4	2.5	7.9	138
Christian	(14.3)	(7.3)	(21.6)	22
Caste/tribe				
Scheduled caste	6.9	3.4	10.3	185
Scheduled tribe	15.0	4.1	19.1	85
Other backward class	7.9	3.0	10.9	588
Other	2.7	1.1	3.8	138
Total	7.7	2.9	10.6	997

Note: Total includes women with less than 5 years of schooling, widowed/divorced/separated/deserted women, women belonging to "other" religions, and women who don't know their caste/ tribe, who are not shown separately.

() Based on 25-49 unweighted cases

Table 14 Birth order

Percent distribution of births to all women during the three years preceding the survey by birth order, according to background characteristics, Telangana, 2015-16

Background characteristic	Birth order				Total	Number of births
	1	2	3	4+		
Mother's current age						
15-19	86.8	13.2	0.0	0.0	100.0	85
20-29	42.3	42.9	12.0	2.8	100.0	1,174
30-39	29.6	42.0	16.2	12.3	100.0	129
Residence						
Urban	43.9	44.5	7.7	3.9	100.0	635
Rural	43.7	38.0	15.0	3.3	100.0	761
Mother's schooling						
No schooling	27.3	39.7	21.8	11.2	100.0	252
<5 years complete	26.0	48.3	16.2	9.4	100.0	50
5-9 years complete	36.8	45.6	14.8	2.8	100.0	298
10-11 years complete	48.5	38.8	10.2	2.4	100.0	333
12 or more years complete	55.7	39.4	4.6	0.3	100.0	463
Religion						
Hindu	44.9	41.6	11.1	2.4	100.0	1,198
Muslim	35.0	36.1	16.0	13.0	100.0	159
Christian	(41.4)	(45.8)	(10.2)	(2.5)	100.0	37
Caste/tribe						
Scheduled caste	44.0	41.4	11.9	2.8	100.0	255
Scheduled tribe	41.9	32.0	20.4	5.8	100.0	118
Other backward class	43.8	42.0	10.4	3.8	100.0	810
Other	44.8	42.2	10.5	2.6	100.0	204
Total	43.8	41.0	11.6	3.6	100.0	1,396

Note: Total includes information on births to women age 40-49, women belonging to "other" religions, and women who don't know their caste/tribe, which is not shown separately.

() Based on 25-49 unweighted cases

Table 15 Birth intervals

Percent distribution of births during the five years preceding the survey by interval since the preceding birth, and median number of months since the preceding birth, according to background characteristics, Telangana, 2015-16

Background characteristic	Months since preceding birth						Total	Number of non-first order births	Median number of months since preceding birth
	7-17	18-23	24-35	36-47	48-59	60+			
Mother's current age									
20-29	15.0	20.4	34.8	14.3	7.2	8.4	100.0	1,077	27.5
30-39	8.9	10.6	20.4	16.8	10.2	33.1	100.0	259	41.0
Residence									
Urban	16.3	16.9	31.5	15.1	6.3	13.9	100.0	621	28.3
Rural	12.1	19.3	31.8	14.5	8.7	13.6	100.0	741	29.9
Mother's schooling									
No schooling	11.3	15.9	26.1	17.2	8.2	21.4	100.0	346	34.3
<5 years complete	8.1	17.9	36.7	9.4	14.9	13.0	100.0	75	31.4
5-9 years complete	16.9	21.3	32.4	12.5	7.0	9.9	100.0	329	27.3
10-11 years complete	15.7	17.6	36.8	14.6	7.4	7.9	100.0	310	27.6
12 or more years complete	13.7	18.0	30.8	16.2	6.1	15.2	100.0	303	29.4
Religion									
Hindu	12.6	17.5	30.9	15.5	8.2	15.3	100.0	1,126	30.6
Muslim	22.0	24.6	34.4	8.7	4.0	6.4	100.0	194	24.4
Christian	(14.8)	(5.3)	(41.2)	(23.5)	(9.6)	(5.4)	100.0	42	(28.7)
Caste/tribe									
Scheduled caste	16.7	14.1	29.9	15.2	10.5	13.7	100.0	255	29.9
Scheduled tribe	13.7	20.2	28.1	15.8	9.2	12.9	100.0	118	31.8
Other backward class	12.9	18.5	34.7	14.3	7.0	12.6	100.0	772	28.4
Other	15.5	19.8	25.2	15.5	5.6	18.5	100.0	209	30.3
Birth order									
2-3	14.0	18.0	31.6	14.9	7.7	13.8	100.0	1,279	29.3
4-6	14.7	21.4	33.1	13.0	6.8	11.0	100.0	80	26.9
Sex of preceding birth									
Male	14.9	19.8	31.4	14.5	4.8	14.6	100.0	677	28.2
Female	13.1	16.6	31.9	15.1	10.4	12.9	100.0	685	30.0
Survival of preceding birth									
Living	12.8	18.5	32.3	15.1	7.6	13.7	100.0	1,305	29.3
Dead	40.8	11.6	17.7	7.2	7.7	15.1	100.0	57	22.9
Total	14.0	18.2	31.7	14.8	7.6	13.7	100.0	1,362	29.1

Note: First-order births are excluded from the table. The interval for multiple births is the number of months since the preceding pregnancy that ended in a live birth. Total includes information on births to women age 15-19 or 40-49, births whose caste/tribe is not known, and births of birth order 7 or more, which is not shown separately.

() Based on 25-49 unweighted cases

Table 16 Fertility preferences by number of living children

Percent distribution of currently married women and men age 15-49 by desire for children, according to number of living children, Telangana, 2015-16

Desire for children	Number of living children ¹							Total
	0	1	2	3	4	5	6+	
WOMEN								
Want another soon ²	72.6	39.5	4.2	2.1	1.8	3.0	(4.6)	15.1
Want another later ³	6.2	22.7	1.3	1.2	0.8	0.0	(2.1)	5.4
Want another, undecided when	2.2	4.0	0.5	0.2	0.2	0.0	(2.9)	1.2
Undecided	3.7	2.9	1.0	0.5	2.4	0.0	(0.0)	1.5
Want no more	2.8	12.5	14.4	10.7	8.1	12.3	(14.9)	12.1
Sterilized ⁴	2.0	11.7	69.7	75.3	76.8	73.6	(58.6)	55.8
Declared infecund	10.3	6.5	8.9	10.0	9.8	11.2	(16.9)	8.9
Missing	0.2	0.2	0.0	0.0	0.0	0.0	(0.0)	0.1
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Number	425	1,002	2,631	1,165	291	80	25	5,619
MEN								
Want another soon ²	35.3	48.4	5.2	4.8	(1.8)	*	*	17.4
Want another later ³	4.1	14.0	1.5	2.3	(2.2)	*	*	4.5
Want another, undecided when	1.1	4.9	0.8	0.0	(0.0)	*	*	1.6
Undecided	10.2	1.1	0.5	0.6	(3.6)	*	*	2.2
Want no more	22.2	22.1	72.8	66.5	(84.4)	*	*	55.8
Sterilized ⁴	18.8	8.3	17.5	25.0	(8.0)	*	*	16.4
Declared infecund	8.3	1.3	1.8	0.8	(0.0)	*	*	2.2
Missing	0.0	0.0	0.0	0.0	(0.0)	*	*	0.0
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Number	86	132	277	120	40	6	5	665

¹ Includes current pregnancy of woman/wife

² Want next birth within 2 years

³ Want to delay next birth for 2 or more years

⁴ For women: Includes both female and male sterilization and women who have had a hysterectomy;

For men: Includes male sterilization and men who mention in response to the question about desire for children that their wife has been sterilized

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

Table 17 Desire not to have any more children

Percentage of currently married women and men age 15-49 who want no more children by number of living children, according to background characteristics, Telangana, 2015-16

Background characteristic	Women: Number of living children ¹						Men: Number of living children ¹					
	0	1	2	3	4+	Total	0	1	2	3	4+	Total
Age												
15-24	0.7	6.7	74.6	77.3	*	33.0	*	*	*	*	*	(26.4)
25-34	2.0	19.0	84.7	85.4	78.3	68.9	(34.0)	10.6	80.9	(82.3)	*	55.8
35-49	23.9	66.0	86.9	87.1	85.8	83.0	(60.9)	59.4	97.5	94.2	(93.8)	87.5
Residence												
Urban	2.5	27.0	86.6	88.2	84.3	69.3	(41.8)	(31.3)	92.5	(86.7)	*	69.8
Rural	6.8	21.4	81.7	84.5	84.5	66.6	40.2	29.4	88.6	94.4	(88.2)	74.1
Schooling												
No schooling	11.4	42.2	80.2	85.4	85.2	75.4	*	(56.0)	91.0	90.9	*	84.9
<5 years complete	*	(37.3)	81.4	85.8	*	75.5	*	*	*	*	*	(73.9)
5-9 years complete	5.6	24.6	87.4	87.3	78.2	70.8	*	(35.2)	91.1	(87.5)	*	75.8
10-11 years complete	0.0	20.1	84.7	81.8	(88.6)	64.4	*	(29.9)	91.8	*	*	70.0
12 or more years complete	2.4	13.8	86.5	93.9	*	51.9	(28.7)	(17.2)	88.4	*	*	58.3
Religion												
Hindu	5.4	24.5	84.8	87.0	85.1	68.4	39.8	30.7	91.1	95.6	(87.8)	72.2
Muslim	(0.0)	18.0	72.1	78.2	81.3	62.5	*	*	*	*	*	63.4
Christian	*	(27.1)	90.5	(88.9)	*	69.1	*	*	*	*	*	*
Caste/tribe												
Scheduled caste	7.5	25.7	80.7	86.7	85.0	65.8	*	*	(89.7)	*	*	77.2
Scheduled tribe	(0.0)	24.3	79.4	87.5	78.5	67.5	*	*	(84.3)	*	*	68.9
Other backward class	4.6	22.2	84.7	87.2	84.6	68.1	36.2	27.4	92.2	89.8	(89.4)	71.2
Other	4.3	30.4	87.2	76.9	87.7	68.9	*	*	(86.9)	*	*	73.3
Don't know	*	*	(88.3)	*	*	(78.1)	*	*	*	*	*	*
Number of living sons²												
0	4.8	20.7	77.9	80.7	(78.8)	39.2	41.0	29.8	80.2	*	*	51.0
1	na	27.9	84.7	87.3	84.5	74.7	na	31.0	91.3	92.2	*	78.3
2	na	na	86.9	86.4	85.0	86.6	na	na	96.2	(92.0)	*	94.1
3	na	na	na	85.1	81.1	83.6	na	na	na	*	*	*
4+	na	na	na	na	na	(93.2)	na	na	na	na	na	*
Total	4.8	24.2	84.1	86.0	84.4	67.8	41.0	30.4	90.3	91.5	88.8	72.2

Note: Women who have been sterilized or whose husband has been sterilized are considered to want no more children. Men who are sterilized or who mention in response to the question about desire for children that their wife has been sterilized are considered to want no more children.

na = Not applicable

¹ Includes current pregnancy of woman/wife

² Excludes pregnant women and men with pregnant wives

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

Table 18 Ideal number of children

Percent distribution of women and men age 15-49 by ideal number of children and mean ideal number of children, by number of living children, Telangana, 2015-16

Ideal number of children	Number of living children ¹							Total
	0	1	2	3	4	5	6+	
WOMEN								
0	16.2	3.0	4.1	9.7	13.2	11.5	(29.9)	8.5
1	3.9	8.3	1.2	0.4	0.0	0.0	(0.0)	2.7
2	75.0	82.6	85.3	48.5	31.3	15.9	(12.0)	72.7
3	3.7	4.4	7.1	34.5	19.8	25.4	(7.7)	11.3
4	0.5	1.1	1.8	5.7	33.5	30.9	(35.4)	3.8
5	0.2	0.0	0.0	0.7	1.3	15.1	(0.0)	0.4
6+	0.0	0.0	0.0	0.2	0.7	1.2	(15.0)	0.1
Non-numeric responses	0.4	0.6	0.4	0.5	0.2	0.0	(0.0)	0.4
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Number	1,881	1,111	2,840	1,293	324	87	30	7,567
Mean ideal number of children for²:								
All respondents	1.7	1.9	2.0	2.3	2.7	3.1	(2.8)	2.0
Number of all respondents	1,873	1,105	2,828	1,287	324	87	30	7,534
Currently married respondents	1.9	1.9	2.0	2.3	2.7	3.1	(2.7)	2.1
Number of currently married respondents	422	1,000	2,619	1,159	291	80	25	5,597
MEN								
0	3.5	0.4	1.2	4.6	(0.0)	*	*	2.4
1	4.1	12.2	5.8	2.5	(2.4)	*	*	5.4
2	77.9	74.2	80.4	52.9	(53.2)	*	*	73.8
3	11.1	10.1	9.7	23.8	(30.4)	*	*	13.0
4	2.5	3.1	1.9	15.4	(14.0)	*	*	4.4
5	0.0	0.0	0.0	0.0	(0.0)	*	*	0.1
6+	0.0	0.0	0.3	0.0	(0.0)	*	*	0.2
Non-numeric responses	1.0	0.0	0.6	0.8	(0.0)	*	*	0.8
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Number	466	142	277	120	41	6	5	1,058
Mean ideal number of children for²:								
All respondents	2.1	2.0	2.1	2.4	(2.6)	*	*	2.1
Number of all respondents	461	142	276	119	41	6	4	1,049
Currently married respondents	2.1	2.0	2.1	2.4	(2.6)	*	*	2.2
Number of currently married respondents	86	132	276	119	40	6	4	661

¹ Includes current pregnancy of woman/wife

² Means are calculated excluding respondents who gave non-numeric responses

() Based on 25-49 unweighted cases

* Based on fewer than 25 unweighted cases

Table 19 Indicators of sex preference

Percentage of women and men age 15-49 who want more sons than daughters, percentage who want more daughters than sons, percentage who want at least one son, and percentage who want at least one daughter, according to background characteristics, Telangana, 2015-16

Background characteristic	Women				Men				
	Percentage who want more sons than daughters	Percentage who want more daughters than sons	Percentage who want at least one son	Percentage who want at least one daughter	Percentage who want more sons than daughters	Percentage who want more daughters than sons	Percentage who want at least one son	Percentage who want at least one daughter	Number of men
Age									
15-19	3.8	2.4	73.5	72.9	10.5	4.1	85.3	84.2	170
20-29	8.3	3.6	84.1	82.4	11.6	3.1	89.1	87.9	337
30-39	12.1	4.2	84.1	82.0	16.2	3.7	88.5	85.6	293
40-49	16.3	6.0	77.6	76.2	17.3	5.8	86.8	84.5	246
Residence									
Urban	8.8	4.2	82.4	80.9	15.4	3.8	84.5	82.1	500
Rural	12.1	4.1	80.1	78.4	12.9	4.4	90.7	89.3	547
Schooling									
No schooling	15.1	5.2	80.0	78.3	16.4	3.5	90.7	88.2	178
<5 years complete	16.9	6.3	81.0	79.9	14.7	4.1	87.0	89.8	44
5-9 years complete	9.5	4.0	81.5	80.4	14.5	5.9	86.6	82.6	254
10-11 years complete	7.2	4.2	80.6	79.5	13.8	3.6	90.4	88.1	241
12 or more years complete	6.8	2.5	83.0	80.8	12.6	3.4	85.2	84.9	330
Marital status									
Never married	4.7	2.7	70.9	69.8	12.1	3.5	87.3	86.0	374
Currently married	11.8	4.5	84.5	82.8	15.3	4.3	87.8	85.5	660
Widowed/divorced/separated/deserted	12.7	4.1	73.5	71.7	*	*	*	*	13
Religion									
Hindu	10.2	3.9	81.6	80.1	12.6	3.9	87.3	85.7	900
Muslim	12.8	6.7	75.8	73.9	25.8	5.4	89.2	86.3	120
Christian	11.4	3.3	86.8	84.7	(12.8)	(3.2)	(97.7)	(87.5)	27

Continued...

Table 19 Indicators of sex preference—Continued

Percentage of women and men age 15-49 who want more sons than daughters, percentage who want more daughters than sons, percentage who want at least one son, and percentage who want at least one daughter, according to background characteristics, Telangana, 2015-16

Background characteristic	Women				Men				Number of men
	Percentage who want more sons than daughters	Percentage who want more daughters than sons	Percentage who want at least one son	Percentage who want at least one daughter	Percentage who want more sons than daughters	Percentage who want more daughters than sons	Percentage who want at least one son	Percentage who want at least one daughter	
Caste/tribe									
Scheduled caste	11.0	3.0	80.5	78.6	14.8	6.4	89.9	86.6	159
Scheduled tribe	11.5	3.6	82.5	80.3	7.0	2.7	96.0	95.1	87
Other backward class	10.5	4.6	81.7	80.6	13.5	3.6	86.0	84.4	659
Other	9.1	4.3	79.5	77.1	20.0	4.4	88.7	86.0	143
Don't know	17.1	3.2	74.4	69.9	nc	nc	nc	nc	0
Total	10.5	4.1	81.2	79.6	14.1	4.1	87.8	85.9	1,047

Note: Table excludes women and men who gave non-numeric responses to the questions on ideal number of children or ideal number of sons or daughters. Total includes women/men belonging to "other" religions, who are not shown separately.

nc = No cases

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

Table 20 Knowledge of contraceptive methods

Percentage of all women and men, currently married women and men, and never married women and men who know any contraceptive method by specific method and residence, Telangana, 2015-16

Method	Women			Men		
	All women	Currently married women	Never married women	All men	Currently married men	Never married men
URBAN						
Any method	97.9	98.9	94.7	98.5	99.6	96.9
Any modern method	97.9	98.9	94.7	98.5	99.6	96.9
Female sterilization	97.2	98.3	93.3	93.7	96.8	88.7
Male sterilization	86.9	91.4	70.7	84.8	87.3	80.7
IUD or PPIUD	35.5	40.3	22.0	13.0	15.5	8.6
Injectables	59.2	61.0	55.7	59.8	60.0	60.6
Pill	69.7	71.0	68.2	73.6	73.4	73.1
Condom/ <i>Nirodh</i>	58.3	58.6	61.6	94.1	95.5	91.8
Female condom	19.5	20.2	19.0	40.3	41.6	38.8
Lactational amenorrhoea method (LAM)	5.7	6.6	3.1	4.9	5.9	3.1
Emergency contraception	23.5	25.4	19.5	31.8	31.4	32.9
Other modern method	0.3	0.4	0.0	0.0	0.0	0.0
Pill, IUD/PPIUD, and condom/ <i>Nirodh</i> ¹	29.3	33.1	18.6	12.0	13.8	8.6
Any traditional method	16.0	19.3	6.9	39.2	44.9	29.8
Rhythm	14.5	17.3	6.8	22.5	26.9	16.1
Withdrawal	8.7	10.7	3.3	34.7	40.5	24.9
Other	0.3	0.4	0.2	3.8	1.9	6.8
Mean number of methods known by respondents age 15-49	4.8	5.0	4.2	5.6	5.8	5.3
Number of respondents age 15-49	3,619	2,637	754	506	302	199
RURAL						
Any method	97.3	98.7	90.4	98.3	98.6	98.3
Any modern method	97.3	98.7	90.3	98.2	98.4	98.3
Female sterilization	96.5	98.1	89.0	95.3	97.2	91.6
Male sterilization	78.3	81.7	62.9	77.9	80.9	71.3
IUD or PPIUD	20.1	21.4	17.4	8.1	7.6	9.2
Injectables	45.0	45.4	48.8	46.9	43.9	52.3
Pill	53.2	53.5	56.9	61.9	58.9	68.2
Condom/ <i>Nirodh</i>	36.0	34.3	49.8	82.1	77.5	92.7
Female condom	13.5	13.0	19.2	27.1	24.2	33.6
Lactational amenorrhoea method (LAM)	4.5	4.6	4.5	1.9	1.9	1.8
Emergency contraception	13.4	13.7	14.6	26.3	26.1	26.1
Other modern method	0.1	0.1	0.0	0.0	0.0	0.0
Pill, IUD/PPIUD, and condom/ <i>Nirodh</i> ¹	13.0	13.5	13.5	7.6	7.1	8.6
Any traditional method	13.1	14.6	8.2	36.8	39.4	32.8
Rhythm	11.6	12.9	7.5	19.5	21.3	16.4
Withdrawal	7.1	8.1	3.5	31.3	33.4	28.1
Other	0.6	0.7	0.6	5.7	6.8	3.6
Mean number of methods known by respondents age 15-49	3.8	3.9	3.7	4.8	4.8	5.0
Number of respondents age 15-49	3,948	2,983	638	552	363	181

Continued...

Table 20 Knowledge of contraceptive methods—Continued

Percentage of all women and men, currently married women and men, and never married women and men who know any contraceptive method by specific method and residence, Telangana, 2015-16

Method	Women			Men		
	All women	Currently married women	Never married women	All men	Currently married men	Never married men
TOTAL						
Any method	97.6	98.8	92.7	98.4	99.0	97.6
Any modern method	97.6	98.8	92.7	98.4	98.9	97.6
Female sterilization	96.8	98.2	91.3	94.5	97.0	90.1
Male sterilization	82.4	86.2	67.1	81.2	83.8	76.2
IUD or PPIUD	27.5	30.2	19.9	10.5	11.2	8.9
Injectables	51.8	52.7	52.6	53.0	51.2	56.7
Pill	61.1	61.7	63.0	67.5	65.5	70.8
Condom/ <i>Nirodh</i>	46.7	45.7	56.2	87.9	85.7	92.2
Female condom	16.4	16.4	19.1	33.4	32.1	36.3
Lactational amenorrhoea method (LAM)	5.1	5.6	3.7	3.3	3.7	2.5
Emergency contraception	18.2	19.2	17.3	28.9	28.5	29.7
Other modern method	0.2	0.2	0.0	0.0	0.0	0.0
Pill, IUD/PPIUD, and condom/ <i>Nirodh</i> ¹	20.8	22.7	16.3	9.7	10.1	8.6
Any traditional method	14.5	16.8	7.5	38.0	41.9	31.2
Rhythm	13.0	14.9	7.1	21.0	23.9	16.2
Withdrawal	7.9	9.3	3.4	32.9	36.6	26.4
Other	0.5	0.6	0.4	4.8	4.6	5.3
Mean number of methods known by respondents age 15-49	4.3	4.4	4.0	5.2	5.2	5.1
Number of respondents age 15-49	7,567	5,619	1,392	1,058	665	380

IUD = Intrauterine device ; PPIUD = Postpartum intrauterine device

¹ All three methods

Table 21 Current use of contraception by background characteristics

Background characteristic	Percent distribution of currently married women by contraceptive method currently used, according to background characteristics, Telangana, 2015-16											Number of women				
	Modern method					Any traditional method		Traditional method		Not currently using	Total					
	Any method	Any modern method	Female sterilization	Male sterilization	Pill	IUD or PPIUD	Injectables	Condom/Nirodha	LAM				Other modern method	Any traditional method	Rhythm	Withdrawal
Age																
15-19	5.1	5.1	5.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	94.9	100.0	180
20-24	27.0	26.8	24.3	0.4	0.4	0.8	0.0	0.4	0.1	0.2	0.2	0.1	0.1	73.0	100.0	895
25-29	50.8	50.8	48.3	1.0	0.4	0.2	0.1	0.7	0.0	0.0	0.1	0.0	0.0	49.2	100.0	1,288
30-39	69.5	69.0	65.6	2.0	0.3	0.4	0.0	0.6	0.0	0.0	0.5	0.0	0.0	30.5	100.0	1,852
40-49	72.7	72.6	69.8	2.6	0.1	0.1	0.0	0.1	0.0	0.0	0.0	0.0	0.0	27.3	100.0	1,404
Residence																
Urban	58.8	58.3	54.9	1.4	0.5	0.6	0.0	0.8	0.0	0.0	0.5	0.4	0.0	41.2	100.0	2,637
Rural	55.8	55.7	53.6	1.7	0.1	0.1	0.0	0.2	0.0	0.0	0.0	0.0	0.0	44.2	100.0	2,983
Schooling																
No schooling	66.0	66.0	63.9	1.9	0.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	34.0	100.0	2,068
<5 years complete	61.5	61.5	59.1	2.1	0.0	0.0	0.0	0.3	0.0	0.0	0.0	0.0	0.0	38.5	100.0	277
5-9 years complete	58.1	57.9	55.9	1.6	0.2	0.0	0.0	0.2	0.0	0.0	0.2	0.0	0.0	41.9	100.0	1,152
10-11 years complete	53.7	53.7	50.7	1.3	0.6	0.4	0.1	0.5	0.0	0.1	0.0	0.0	0.0	46.3	100.0	1,004
12 or more years complete	41.9	41.1	36.3	1.1	0.4	1.3	0.1	1.6	0.1	0.2	0.8	0.8	0.0	58.1	100.0	1,118
Religion																
Hindu	58.1	57.8	55.1	1.7	0.2	0.3	0.0	0.5	0.0	0.1	0.3	0.2	0.0	41.9	100.0	4,913
Muslim	49.6	49.6	46.3	0.3	1.3	0.7	0.2	0.7	0.0	0.0	0.0	0.0	0.0	50.4	100.0	523
Christian	57.3	57.3	54.3	2.2	0.2	0.0	0.7	0.0	0.0	0.0	0.0	0.0	0.0	42.7	100.0	173
Caste/tribe																
Scheduled caste	52.5	52.5	49.2	2.4	0.1	0.5	0.1	0.1	0.0	0.0	0.0	0.0	0.0	47.5	100.0	987
Scheduled tribe	56.6	56.6	54.7	1.5	0.2	0.0	0.0	0.2	0.0	0.0	0.0	0.0	0.0	43.4	100.0	450
Other backward class	57.5	57.3	54.6	1.6	0.3	0.1	0.0	0.5	0.0	0.1	0.2	0.2	0.0	42.5	100.0	3,285
Other	60.7	60.2	57.2	0.5	0.3	1.3	0.0	0.9	0.0	0.0	0.6	0.6	0.0	39.3	100.0	859
Don't know	(74.1)	(74.1)	(67.8)	(3.6)	(2.7)	(0.0)	(0.0)	(0.0)	(0.0)	(0.0)	(0.0)	(0.0)	(0.0)	(25.9)	100.0	38

Continued...

Table 21 Current use of contraception by background characteristics—Continued

Background characteristic	Modern method											Total	Number of women					
	Any method	Any modern method	Female sterilization	Male sterilization	Pill	IUD or PPIUD	Injectables	Condom/Nirodh	LAM	Other modern method	Any traditional method			Traditional method	Withdrawal	Not currently using		
Number of living children																		
No children	2.0	2.0	1.2	0.4	0.2	0.0	0.0	0.0	0.2	0.0	0.0	0.0	0.0	0.0	0.0	98.0	100.0	545
1 child	16.5	15.7	11.3	0.9	0.6	1.4	0.0	1.3	0.8	0.1	0.2	0.0	0.0	0.0	0.0	83.5	100.0	966
1 son	16.3	15.5	10.8	0.9	1.0	1.8	0.0	0.8	0.8	0.2	0.1	0.0	0.0	0.0	0.0	83.7	100.0	512
No sons	16.7	16.0	11.8	0.8	0.3	1.0	0.0	1.9	0.7	0.0	0.3	0.0	0.0	0.0	0.0	83.3	100.0	454
2 children	72.2	72.0	69.2	1.9	0.2	0.2	0.0	0.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	27.8	100.0	2,579
1 or more sons	74.4	74.3	71.5	1.9	0.2	0.2	0.0	0.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	25.6	100.0	2,138
No sons	61.3	60.7	58.1	1.6	0.4	0.1	0.0	0.4	0.0	0.0	0.2	0.0	0.0	0.0	0.0	38.7	100.0	441
3 children	77.4	77.4	75.1	1.9	0.1	0.1	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	22.6	100.0	1,138
1 or more sons	78.4	78.4	76.1	1.9	0.1	0.1	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	21.6	100.0	1,019
No sons	69.1	69.1	66.6	1.9	0.0	0.0	0.0	0.6	0.0	0.0	0.0	0.0	0.0	0.0	0.0	30.9	100.0	119
4+ children	76.4	76.4	73.6	2.3	0.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	23.6	100.0	392
1 or more sons	76.8	76.8	74.2	2.2	0.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	23.2	100.0	360
No sons	(72.7)	(72.7)	(66.8)	(3.3)	(2.6)	(0.0)	(0.0)	(0.0)	(0.0)	(0.0)	(0.0)	(0.0)	(0.0)	(0.0)	(0.0)	(27.3)	100.0	32
Total	57.2	56.9	54.2	1.6	0.3	0.3	0.0	0.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	42.8	100.0	5,619

Note: If more than one method is used, only the most effective method is considered in this table. Total includes women belonging to "other" religions, who are not shown separately.
 IUD = Intrauterine device; PPIUD = Postpartum intrauterine device; LAM = Lactational amenorrhoea method
 () Based on 25-49 unweighted cases

Table 22 Current use of contraceptive methods by district

Percentage of currently married women age 15-49 years using any contraceptive method, any modern method, female sterilization, male sterilization, any modern spacing method, and any traditional method, according to district, Telangana, 2015-16

District	Any method	Any modern method	Female sterilization	Male sterilization	Any modern spacing method ¹	Any traditional method ²	Number of women
Adilabad	48.6	48.1	45.3	1.6	1.2	0.6	415
Hyderabad	55.8	55.5	53.7	0.0	1.7	0.3	813
Karimnagar	38.6	38.6	34.2	4.2	0.1	0.0	599
Khammam	69.1	69.1	68.2	0.0	0.9	0.0	444
Mahbubnagar	64.3	64.3	64.0	0.0	0.3	0.0	502
Medak	49.6	49.1	46.5	0.8	1.8	0.6	453
Nalgonda	68.0	68.0	66.3	0.3	1.3	0.0	512
Nizamabad	47.0	47.0	46.7	0.0	0.3	0.0	358
Rangareddy	69.1	68.6	65.9	1.0	1.6	0.5	989
Warangal	50.8	50.8	41.2	7.9	1.7	0.0	534
Telangana	57.2	56.9	54.2	1.6	1.2	0.2	5,619

¹ Includes pill, IUD/PPIUD, injectables, male condom, female condom, and lactational amenorrhoea method (LAM)

² Includes rhythm, withdrawal, and other traditional methods

Table 23 Hysterectomy

Percentage of women age 15-49 who have had a hysterectomy, and among women with a hysterectomy, percent distribution by place the hysterectomy was performed, according to background characteristics, Telangana, 2015-16

Background characteristic	Percentage of women who have had a hysterectomy	Number of women	Place of hysterectomy (%)			Total	Number of women with a hysterectomy
			Public health sector	Private health sector ¹	Elsewhere or don't know place		
Age							
< 30	1.1	3,780	7.4	92.6	0.0	100.0	41
30-39	9.7	2,060	15.0	85.0	0.0	100.0	199
40-49	20.0	1,727	21.9	77.6	0.5	100.0	345
Residence							
Urban	5.0	3,619	29.0	70.6	0.4	100.0	180
Rural	10.3	3,948	13.8	85.9	0.2	100.0	405
Schooling							
No schooling	15.8	2,458	17.0	82.9	0.2	100.0	389
<5 years complete	15.1	337	(29.3)	(70.7)	(0.0)	100.0	51
5-9 years complete	6.3	1,493	16.5	83.5	0.0	100.0	95
10-11 years complete	1.7	1,435	*	*	*	100.0	24
12 or more years complete	1.4	1,844	*	*	*	100.0	26
Religion							
Hindu	8.2	6,529	18.0	81.7	0.3	100.0	537
Muslim	4.0	774	(20.7)	(79.3)	(0.0)	100.0	31
Christian	7.0	248	*	*	*	100.0	17
Caste/tribe							
Scheduled caste	7.5	1,395	24.0	76.0	0.0	100.0	105
Scheduled tribe	13.3	597	17.4	82.6	0.0	100.0	79
Other backward class	7.4	4,393	17.3	82.2	0.5	100.0	324
Other	6.4	1,137	18.1	81.9	0.0	100.0	73
Don't know	7.6	44	*	*	*	100.0	3
Total	7.7	7,567	18.5	81.2	0.3	100.0	585

Note: Total includes women belonging to "other" religions, who are not shown separately.

¹ Includes nongovernmental organizations or trust hospitals/clinics

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

Table 24 Contraceptive use by men at last sexual intercourse

Percent distribution of currently married men and sexually active unmarried men age 15-49 by contraceptive used the last time they had sex, according to type of partner and background characteristics, Telangana, 2015-16

Background characteristic	Any method	Modern method						Not used at last sex	Total	Number of men
		Any modern method	Female sterilization	Male sterilization	Pill	Condom/Nirodh	Other modern method			
Type of sexual partner										
Currently married	10.3	10.3	6.3	2.0	0.3	1.5	0.1	89.7	100.0	665
Wife	12.9	12.9	8.3	2.0	0.4	2.1	0.1	87.1	100.0	494
Other	2.7	2.7	0.5	2.2	0.0	0.0	0.0	97.3	100.0	171
Age										
20-24	(7.6)	(7.6)	(0.0)	(0.0)	(0.0)	(7.6)	(0.0)	(92.4)	100.0	35
25-29	5.2	5.2	3.9	0.0	0.0	1.3	0.0	94.8	100.0	124
30-34	11.9	11.9	6.7	2.0	1.6	1.6	0.0	88.1	100.0	137
35-39	13.8	13.8	7.4	2.1	0.0	3.8	0.5	86.2	100.0	132
40-44	10.0	10.0	9.3	0.7	0.0	0.0	0.0	90.0	100.0	112
45-49	11.4	11.4	5.9	5.5	0.0	0.0	0.0	88.6	100.0	130
Residence										
Urban	6.9	6.9	3.3	0.4	0.7	2.6	0.0	93.1	100.0	304
Rural	13.5	13.5	8.7	3.4	0.0	1.2	0.2	86.5	100.0	367
Schooling										
No schooling	9.7	9.7	6.3	3.4	0.0	0.0	0.0	90.3	100.0	158
<5 years complete	(21.6)	(21.6)	(21.6)	(0.0)	(0.0)	(0.0)	(0.0)	(78.4)	100.0	40
5-9 years complete	11.5	11.5	8.5	1.1	0.0	1.9	0.0	88.5	100.0	159
10-11 years complete	9.2	9.2	4.9	1.2	0.0	2.7	0.5	90.8	100.0	145
12 or more years complete	8.7	8.7	1.6	2.7	1.3	3.2	0.0	91.3	100.0	170

Continued...

Table 24 Contraceptive use by men at last sexual intercourse—Continued

Percent distribution of currently married men and sexually active unmarried men age 15–49 by contraceptive used the last time they had sex, according to type of partner and background characteristics, Telangana, 2015–16

Background characteristic	Modern method							Total	Number of men	
	Any method	Any modern method	Female sterilization	Male sterilization	Pill	Condom/Nirodh	Other modern method			Not used at last sex
Religion										
Hindu	10.7	10.7	6.5	2.3	0.4	1.6	0.0	89.3	100.0	591
Muslim	10.5	10.5	4.5	0.0	0.0	4.8	1.2	89.5	100.0	59
Caste/tribe										
Scheduled caste	7.9	7.9	6.1	1.8	0.0	0.0	0.0	92.1	100.0	100
Scheduled tribe	8.5	8.5	8.5	0.0	0.0	0.0	0.0	91.5	100.0	60
Other backward class	11.9	11.9	6.8	2.7	0.0	2.1	0.2	88.1	100.0	425
Other	8.0	8.0	1.8	0.0	2.5	3.7	0.0	92.0	100.0	87
Total age 15–49	10.5	10.5	6.2	2.0	0.3	1.8	0.1	89.5	100.0	672
Age 50–54	11.4	11.4	6.2	5.2	0.0	0.0	0.0	88.6	100.0	73
Total age 15–54	10.6	10.6	6.2	2.3	0.3	1.6	0.1	89.4	100.0	744

Note: Total includes men whose sexual partner is not currently married, men age 15–19, and Christian men, who are not shown separately.
 IUD = Intrauterine device; PPIUD = Postpartum intrauterine device; LAM = Lactational amenorrhoea method
 () Based on 25–49 unweighted cases

Table 25 Source of modern contraceptive methods

Most recent source of method	URBAN						RURAL						TOTAL					
	Female sterilization	Male sterilization	Condom/Nirodh	All modern methods ¹	Female sterilization	Male sterilization	Condom/Nirodh	All modern methods ¹	Female sterilization	Male sterilization	Condom/Nirodh	All modern methods ¹	Female sterilization	Male sterilization	Condom/Nirodh	All modern methods ¹		
Public health sector	58.8	(86.6)	*	57.7	74.7	90.3	*	74.9	67.2	88.8	(15.8)	66.7						
Government/municipal hospital	33.9	(39.0)	*	33.1	29.6	9.2	*	28.8	31.6	21.4	(1.5)	30.8						
Government dispensary	2.4	(8.7)	*	2.5	4.0	2.1	*	3.9	3.3	4.8	(0.0)	3.2						
UHC/UHP/UFWC	6.7	(7.2)	*	6.5	3.3	1.7	*	3.3	4.9	3.9	(0.0)	4.8						
CHC/rural hospital/Block PHC	4.2	(1.8)	*	4.0	11.0	23.7	*	11.3	7.8	14.7	(0.0)	7.9						
PHC/Additional PHC	5.8	(11.2)	*	5.8	18.4	26.5	*	18.6	12.4	20.3	(11.0)	12.5						
Sub-centre/ANM	0.2	(0.0)	*	0.1	0.4	0.0	*	0.4	0.3	0.0	(0.0)	0.3						
Government mobile clinic	0.0	(0.0)	*	0.0	0.1	0.0	*	0.1	0.1	0.0	(0.0)	0.1						
Camp	5.0	(14.9)	*	5.0	7.8	27.2	*	8.4	6.5	22.2	(0.0)	6.8						
ASHA	0.0	(0.0)	*	0.0	0.0	0.0	*	0.0	0.0	0.0	(3.4)	0.0						
Other public health sector	0.5	(3.7)	*	0.6	0.1	0.0	*	0.1	0.3	1.5	(0.0)	0.3						
NGO or trust hospital/clinic	0.8	(0.0)	*	0.8	0.1	0.0	*	0.1	0.4	0.0	(0.0)	0.4						
Private health sector	39.5	(13.4)	*	40.0	25.0	7.8	*	24.7	31.8	10.1	(49.6)	31.9						
Private hospital	37.7	(8.1)	*	37.4	23.4	7.8	*	23.0	30.2	7.9	(14.5)	29.8						
Private doctor/clinic	1.3	(3.4)	*	1.4	1.2	0.0	*	1.2	1.3	1.4	(0.0)	1.3						
Private mobile clinic	0.1	(0.0)	*	0.2	0.0	0.0	*	0.0	0.0	0.0	(5.8)	0.1						
Pharmacy/drugstore	0.0	(0.0)	*	0.5	0.0	0.0	*	0.1	0.0	0.0	(29.4)	0.3						
Other private health sector	0.4	(1.9)	*	0.5	0.3	0.0	*	0.3	0.4	0.8	(0.0)	0.4						
Other source	0.7	(0.0)	*	1.4	0.2	0.0	*	0.2	0.4	0.0	(34.6)	0.7						
Shop	0.0	(0.0)	*	0.6	0.0	0.0	*	0.0	0.0	0.0	(29.5)	0.3						
Husband	0.0	(0.0)	*	0.1	0.0	0.0	*	0.0	0.0	0.0	(5.1)	0.0						
Other	0.7	(0.0)	*	0.7	0.2	0.0	*	0.2	0.4	0.0	(0.0)	0.4						
Don't know	0.1	(0.0)	*	0.1	0.0	1.9	*	0.1	0.1	1.1	(0.0)	0.1						
Total	100.0	100.0	100.0	100.0	99.9	100.0	100.0	99.9	99.9	100.0	100.0	100.0						
Number of users	1,583	38	21	1,675	1,780	55	5	1,848	3,363	93	26	3,523						

Note: All information in this table is based on women's reports. Table includes all users of modern contraceptive methods regardless of their marital status. All methods includes users of pill, IUD or PPIUD, and injectables, which are not shown separately.
 UHC = Urban health centre; UHP = Urban health post; UFWC = Urban family welfare centre; CHC = Community health centre; PHC = Primary health centre; ANM = Auxiliary nurse midwife;
 ASHA = Accredited Social Health Activist; NGO = Nongovernmental organization
 IUD = Intrauterine device; PPIUD = Postpartum intrauterine device
¹ Excludes standard days method, lactational amenorrhoea method (LAM), and "other" modern methods
 () Based on 25-49 unweighted cases
 * Percentage not shown; based on fewer than 25 unweighted cases

Table 26 Informed choice

Among women who are current users of selected modern contraceptive methods who started the last episode of use within the five years preceding the survey, the percentage who were informed about possible side effects or problems of that method, the percentage who were informed about what to do if they experienced side effects, and the percentage who were ever told by a health or family planning worker about other methods of family planning they could use, by method, according to residence, Telangana, 2015-16

Method	Percentage who were informed about possible side effects or problems of method used	Percentage who were informed about what to do if they experienced side effects	Percentage who were ever told by a health or family planning worker of other methods that could be used	Number of women
URBAN				
Female sterilization ¹	25.6	20.5	30.0	390
Total	26.4	21.0	31.6	410
RURAL				
Female sterilization ¹	22.2	17.8	28.5	472
Total	22.3	18.0	28.7	476
TOTAL				
Female sterilization ¹	23.7	19.0	29.2	862
Total	24.2	19.4	30.1	886

Note: Table includes female sterilization, pill, IUD/PPIUD only and excludes users who obtained their method from friends/relatives or husband. Total includes selected modern contraceptive methods (pill and IUD/PPIUD), which are not shown separately.

IUD = Intrauterine device; PPIUD = Postpartum intrauterine device

¹ Among women who were sterilized in the five years preceding the survey

Table 28 Men's contraception-related perceptions and knowledge

Percentage of men age 15-49 who agree with two specific statements about women and contraception, and percent distribution of men according to their belief about the efficacy of condoms in preventing pregnancy, by background characteristics, Telangana, 2015-16

Background characteristic	Percentage of men who agree		Percentage of men who say that if a male condom is used correctly, it protects against pregnancy:				Total	Number of men
	Contraception is women's business and a man should not have to worry about it	Women who use contraception may become promiscuous	Most of the time	Some times	Not at all	Don't know/unsure ¹		
Age								
15-19	35.8	34.8	49.0	31.9	2.3	16.9	100.0	174
20-24	46.0	39.7	56.5	29.3	0.7	13.5	100.0	155
25-29	48.3	44.5	57.6	35.8	0.0	6.6	100.0	184
30-34	44.4	29.9	57.6	26.3	0.7	15.4	100.0	155
35-39	59.5	48.0	50.5	30.9	0.5	18.1	100.0	140
40-44	49.9	46.4	43.9	32.6	0.7	22.8	100.0	116
45-49	51.1	40.5	50.4	19.7	0.5	29.4	100.0	132
Residence								
Urban	43.5	40.2	56.4	32.9	0.8	10.0	100.0	506
Rural	50.9	40.2	49.3	27.0	0.8	22.9	100.0	552
Schooling								
No schooling	44.8	35.2	36.5	22.5	0.6	40.4	100.0	179
<5 years complete	55.4	46.8	54.1	15.5	0.0	30.4	100.0	44
5-9 years complete	39.8	38.8	52.6	25.9	1.0	20.5	100.0	257
10-11 years complete	47.6	39.6	55.6	33.3	0.7	10.4	100.0	245
12 or more years complete	53.2	43.7	59.1	36.0	0.8	4.0	100.0	333
Religion								
Hindu	50.0	42.1	52.8	28.5	0.8	17.8	100.0	910
Muslim	24.1	22.4	54.9	34.4	0.8	9.9	100.0	120
Christian	(59.7)	(57.1)	(36.9)	(50.6)	(0.0)	(12.4)	100.0	27
Caste/tribe								
Scheduled caste	56.3	49.2	46.0	29.5	0.0	24.6	100.0	160
Scheduled tribe	50.9	47.5	43.6	29.1	4.5	22.9	100.0	87
Other backward class	43.1	35.8	57.1	26.6	0.7	15.6	100.0	666
Other	54.9	46.6	45.0	45.3	0.0	9.7	100.0	145
Total age 15-49	47.3	40.2	52.7	29.8	0.8	16.8	100.0	1,058
Age 50-54	37.7	45.9	35.9	20.6	0.0	43.4	100.0	75
Total age 15-54	46.7	40.6	51.6	29.2	0.7	18.5	100.0	1,133

¹ Includes missing values and those who have never heard of male condoms

() Based on 25-49 unweighted cases

Table 29 Need and demand for family planning among currently married women

Percentage of currently married women age 15-49 with unmet need for family planning, percentage with met need for family planning, the total demand for family planning, and the percentage of the demand for contraception that is satisfied, by background characteristics, Telangana, 2015-16

Background characteristic	Unmet need for family planning			Met need for family planning (currently using)			Total demand for family planning ¹			Percentage of demand satisfied ²	Percentage of demand satisfied by modern methods ³	Number of women
	For spacing	For limiting	Total	For spacing	For limiting	Total	For spacing	For limiting	Total			
Age												
15-19	9.0	0.9	9.8	0.0	5.1	5.1	9.0	5.9	14.9	34.0	34.0	180
20-24	10.6	5.1	15.8	1.8	25.1	27.0	12.5	30.3	42.8	63.1	62.7	895
25-29	6.0	5.6	11.7	1.1	49.7	50.8	7.1	55.3	62.5	81.3	81.2	1,288
30-34	2.2	4.1	6.3	0.8	65.7	66.5	3.0	69.8	72.8	91.4	90.4	937
35-39	0.3	2.2	2.5	0.0	72.5	72.5	0.3	74.7	75.0	96.7	96.3	915
40-44	0.1	2.0	2.0	0.1	71.9	72.0	0.2	73.9	74.1	97.2	97.2	692
45-49	0.0	0.9	0.9	0.0	73.3	73.3	0.0	74.1	74.1	98.9	98.7	712
Residence												
Urban	4.2	4.6	8.8	1.3	57.5	58.8	5.5	62.1	67.6	87.0	86.3	2,637
Rural	3.4	2.6	6.0	0.2	55.5	55.8	3.6	58.2	61.7	90.3	90.3	2,983
Schooling												
No schooling	1.1	2.0	3.1	0.1	65.9	66.0	1.2	67.9	69.1	95.5	95.4	2,068
<5 years complete	1.9	4.0	5.8	0.0	61.5	61.5	1.9	65.4	67.3	91.3	91.3	277
5-9 years complete	3.0	4.1	7.2	0.2	57.9	58.1	3.2	62.1	65.3	89.0	88.6	1,152
10-11 years complete	5.3	4.5	9.8	0.8	52.9	53.7	6.2	57.4	63.6	84.5	84.5	1,004
12 or more years complete	8.5	4.7	13.3	2.5	39.4	41.9	11.0	44.1	55.2	76.0	74.4	1,118
Religion												
Hindu	3.6	3.4	7.0	0.6	57.4	58.1	4.2	60.8	65.0	89.3	88.9	4,913
Muslim	5.4	4.9	10.2	1.5	48.1	49.6	6.8	53.0	59.8	82.9	82.9	523
Christian	3.9	4.3	8.3	0.0	57.3	57.3	3.9	61.6	65.6	87.4	87.4	173
Caste/tribe												
Scheduled caste	3.4	4.6	8.0	0.5	52.0	52.5	4.0	56.5	60.5	86.8	86.8	987
Scheduled tribe	1.5	1.9	3.5	0.0	56.6	56.6	1.5	58.5	60.1	94.2	94.2	450
Other backward class	3.8	3.5	7.3	0.7	56.8	57.5	4.5	60.3	64.8	88.8	88.4	3,285
Other	5.3	3.2	8.5	1.2	59.6	60.7	6.5	62.8	69.3	87.7	86.9	859
Don't know	(4.9)	(4.7)	(9.6)	(2.7)	(71.4)	(74.1)	(7.6)	(76.1)	(83.7)	(88.5)	(88.5)	38
Total	3.8	3.5	7.3	0.7	56.5	57.2	4.5	60.0	64.5	88.7	88.3	5,619

Note: Numbers in this table correspond to the revised definition of unmet need described in Bradley et al., 2012, Revising Unmet Need for Family Planning, DHS Analytical Studies No. 25, ICF International, Calverton, Maryland, USA. Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are: at risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant; pregnant with a mistimed pregnancy; postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception. Women are considered to have unmet need for limiting if they are: at risk of becoming pregnant, not using contraception, and want no (more) children; pregnant with an unwanted pregnancy; postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception. Women who are classified as infecund have no unmet need because they are not at risk of becoming pregnant. Unmet need for family planning is the sum of unmet need for spacing and unmet need for limiting. Total includes women belonging to "other" religions, who are not shown separately.

¹ Total demand is the sum of unmet need and met need

² Percentage of demand satisfied is met need divided by total demand

³ Modern methods include female sterilization, male sterilization, pill, IUD/PPIUD, injectables, male condom, female condom, standard days method, diaphragm, foam/jelly, lactational amenorrhoea method (LAM), and other modern methods

() Based on 25-49 unweighted cases

Table 30 Unmet need for family planning by district

Percentage of currently married women age 15-49 with unmet need for family planning by district, Telangana, 2015-16

District	Total unmet need	Unmet need for spacing	Unmet need for limiting	Number of women
Adilabad	7.8	4.4	3.4	415
Hyderabad	10.7	6.0	4.7	813
Karimnagar	10.0	4.5	5.6	599
Khammam	4.5	2.0	2.5	444
Mahbubnagar	6.3	3.7	2.6	502
Medak	6.5	3.6	2.9	453
Nalgonda	3.5	2.3	1.2	512
Nizamabad	7.6	4.3	3.2	358
Rangareddy	5.7	3.3	2.3	989
Warangal	9.2	2.7	6.5	534
Telangana	7.3	3.8	3.5	5,619

Note: Numbers in this table correspond to the revised definition of unmet need described in Bradley et al., 2012, Revising Unmet Need for Family Planning, DHS Analytical Studies No. 25, ICF International Calverton, Maryland, USA. Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are: at risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant; pregnant with a mistimed pregnancy; postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception. Women are considered to have unmet need for limiting if they are: at risk of becoming pregnant, not using contraception, and want no (more) children; pregnant with an unwanted pregnancy; postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception. Women who are classified as infecund have no unmet need because they are not at risk of becoming pregnant. Unmet need for family planning is the sum of unmet need for spacing and unmet need for limiting.

Table 31 Pregnancy outcome

Percent distribution of last pregnancies among women age 15-49 during the five years preceding the survey by pregnancy outcome, according to background characteristics, Telangana, 2015-16

Background characteristic	Pregnancy outcome				Total	Number of pregnancies
	Live birth	Abortion	Miscarriage	Stillbirth		
Mother's current age						
15-19	94.6	0.7	4.7	0.0	100.0	80
20-29	93.9	2.8	2.9	0.5	100.0	1,458
30-39	86.5	6.4	6.4	0.6	100.0	329
Residence						
Urban	90.7	4.8	4.3	0.2	100.0	898
Rural	94.0	2.0	3.3	0.7	100.0	984
Mother's schooling						
No schooling	93.3	3.2	2.4	1.0	100.0	361
<5 years complete	95.1	1.9	2.2	0.7	100.0	84
5-9 years complete	92.8	3.8	3.1	0.3	100.0	416
10-11 years complete	93.0	1.7	4.9	0.4	100.0	440
12 or more years complete	90.8	4.6	4.4	0.2	100.0	581
Religion						
Hindu	92.6	3.2	3.8	0.4	100.0	1,625
Muslim	93.8	2.1	3.2	0.9	100.0	201
Christian	83.2	11.5	5.3	0.0	100.0	53
Caste/tribe						
Scheduled caste	92.1	3.8	3.4	0.7	100.0	354
Scheduled tribe	91.7	2.2	4.3	1.9	100.0	151
Other backward class	93.8	3.1	3.0	0.1	100.0	1,076
Other	88.0	4.4	7.0	0.6	100.0	292
Total	92.4	3.4	3.8	0.5	100.0	1,882

Note: Total includes pregnancies of women age 40-49, women who belong to "other" religions, and women who don't know their caste/tribe, which are not shown separately.

Table 32 Characteristics of abortions

Among women age 15-49 whose last pregnancy in the last five years ended in an abortion, percent distribution by place of abortion and person who performed the abortion and percentage who had complications from the abortion, according to residence, Telangana, 2015-16

Characteristic of abortion	Urban	Rural	Total
Place of abortion			
Public health sector	(21.6)	(15.0)	19.5
Private health sector ¹	(71.1)	(66.3)	69.6
At home	(7.3)	(18.7)	10.9
Total	100.0	100.0	100.0
Person who performed the abortion			
Doctor	(90.9)	(82.6)	88.3
Nurse/ANM/LHV	(0.0)	(7.6)	2.4
Family member/relative/friend	(1.5)	(0.0)	1.1
Self	(7.5)	(6.8)	7.3
Other	(0.0)	(3.0)	0.9
Total	100.0	100.0	100.0
Complications from the abortion			
Percentage of women who had complications from the abortion	(16.7)	(6.8)	13.5
Number of women with abortions	43	20	63

ANM = Auxiliary nurse midwife; LHV = Lady health visitor

¹ Includes nongovernmental organizations or trust hospitals/clinics

() Based on 25-49 unweighted cases

Table 33 Age at first marriage

Percentage of women and men age 15-49 who were first married by specific exact ages, percentage never married, and median age at first marriage, first cohabitation with spouse, and first sexual intercourse, according to current age, Telangana, 2015-16

Current age	Percentage first married by exact age:					Percentage never married	Number of respondents	Median age at first marriage	Median age at first cohabitation	Median age at first sex
	15	18	20	21	25					
WOMEN										
15-19	1.7	na	na	na	na	81.8	997	a	a	a
20-24	7.2	26.4	48.0	na	na	32.4	1,355	a	a	a
25-29	15.3	39.1	57.3	64.7	84.7	5.8	1,428	19.2	19.0	18.9
30-34	21.6	48.0	63.2	71.0	84.5	2.9	1,042	18.2	18.3	18.2
35-39	27.7	52.1	65.9	70.8	81.3	0.7	1,019	17.7	17.3	17.3
40-44	30.4	56.4	66.7	71.1	78.6	1.6	836	17.1	17.2	16.9
45-49	26.5	48.1	62.9	67.8	76.4	0.3	890	18.2	18.5	16.9
20-49	20.0	43.3	59.6	na	na	8.8	6,570	18.8	18.7	18.5
25-49	23.3	47.7	62.6	68.7	81.6	2.6	5,215	18.3	18.2	18.0
MEN										
15-19	0.0	na	na	na	na	100.0	174	a	a	a
20-24	0.6	1.4	5.7	na	na	79.1	155	a	a	a
25-29	1.2	3.7	12.2	18.1	55.1	31.6	184	24.3	24.6	23.7
30-34	2.4	6.0	10.7	18.2	52.0	10.7	155	24.7	23.9	24.9
35-39	1.1	9.8	21.9	27.4	56.6	4.5	140	24.4	24.1	22.9
40-44	1.8	11.2	28.0	34.8	57.3	1.5	116	24.0	22.6	21.4
45-49	1.4	8.0	20.9	30.6	55.9	0.0	132	23.9	23.5	22.4
20-49	1.4	6.3	15.7	na	na	23.3	883	a	a	a
25-49	1.6	7.3	17.8	24.8	55.2	11.4	728	24.3	23.8	23.0

na = Not applicable due to censoring

a = Omitted because less than 50 percent of the women or men were married, began living with their spouse, or had sexual intercourse for the first time before reaching the beginning of the age group.

Table 34 Early childhood mortality rates

Neonatal, postneonatal, infant, child, and under-five mortality rates for five-year periods preceding the survey, by residence, Telangana, 2015-16

Years preceding the survey	Neonatal mortality (NN)	Postneonatal mortality ¹ (PNN)	Infant mortality (${}_1q_0$)	Child mortality (${}_4q_1$)	Under-five mortality (${}_5q_0$)
URBAN					
0-4	12.2	8.1	20.3	4.6	24.9
5-9	28.9	12.1	41.1	3.9	44.8
10-14	22.2	10.7	32.9	3.3	36.1
RURAL					
0-4	30.6	7.6	38.2	3.2	41.3
5-9	40.1	12.4	52.5	8.5	60.6
10-14	52.7	18.0	70.7	11.6	81.5
TOTAL					
0-4	21.9	7.8	29.8	3.9	33.6
5-9	35.0	12.3	47.3	6.4	53.4
10-14	39.2	14.7	53.9	8.0	61.4

¹ Computed as the difference between the infant and neonatal mortality rates.

Table 35 Early childhood mortality rates by background characteristics

Neonatal, postneonatal, infant, child, and under-five mortality rates for the five-year period preceding the survey, by background characteristics, Telangana, 2015-16

Background characteristic	Neonatal mortality (NN)	Postneonatal mortality ¹ (PNN)	Infant mortality (₁ q ₀)	Child mortality (₄ q ₁)	Under-five mortality (₅ q ₀)
Residence					
Urban	12.2	8.1	20.3	4.6	24.9
Rural	30.6	7.6	38.2	3.2	41.3
Schooling					
No schooling	52.3	16.8	69.1	7.6	76.2
<10 years complete	21.9	6.8	28.7	5.2	33.7
10 or more years complete	10.1	4.7	14.7	0.6	15.3
Religion					
Hindu	21.7	8.7	30.4	3.8	34.1
Muslim	(25.5)	(4.1)	(29.7)	(5.6)	(35.1)
Caste/tribe					
Scheduled caste	(28.9)	(3.9)	(32.8)	(0.0)	(32.8)
Other backward class	20.1	10.7	30.8	5.2	35.8
Other ²	(9.5)	(2.1)	(11.5)	*	*
Child's sex					
Male	23.0	6.3	29.3	4.2	33.5
Female	20.7	9.6	30.3	3.6	33.8
Mother's age at birth					
<20	(21.2)	(8.9)	(30.0)	0.5	(30.5)
20-29	18.3	8.2	26.6	3.8	30.2
Birth order					
1	21.1	8.6	29.7	3.2	32.8
2-3	21.2	6.3	27.5	3.2	30.6
Previous birth interval³					
<2 years	(9.6)	(11.8)	(21.4)	(2.4)	(23.7)
2-3 years	24.6	5.6	30.3	5.0	35.2
4 years or more	(40.7)	(4.0)	(44.7)	*	*
Total	21.9	7.8	29.8	3.9	33.6

Note: Total includes children belonging to Christian or "other" religions, scheduled tribe children, children with mother's age at birth 30-39 or 40-49, and children of birth order 4 or more, who are not shown separately.

¹ Computed as the difference between the infant and neonatal mortality rates

² Not belonging to a scheduled caste, scheduled caste, or other backward class

³ Excludes first-order births

() Based on 250-499 unweighted person-years of exposure to the risk of death

* Rate not shown; based on fewer than 250 unweighted person-years of exposure to the risk of death

Table 36 High-risk fertility behaviour

Percent distribution of children born in the five years preceding the survey by category of elevated risk of mortality and the risk ratio, and percent distribution of currently married women by category of risk if they were to conceive a child at the time of the survey, Telangana, 2015-16

Risk category	Births in the 5 years preceding the survey		Percentage of currently married women ¹
	Percentage of births	Risk ratio	
Not in any high-risk category	34.9	1.0	69.3 ^a
Unavoidable risk category			
First order births between ages 18 and 34 years	40.5	1.1	7.9
Single high-risk category			
Mother's age <18	3.3	0.4	0.2
Mother's age >34	1.0	(7.4)	7.8
Birth interval <24 months	16.7	0.5	6.3
Birth order >3	2.1	2.2	2.2
Subtotal	23.1	0.9	16.5
Multiple high-risk category			
Mother's age <18 and birth interval <24 months ²	0.1	*	0.0
Mother's age >34 and birth interval <24 months	0.1	*	0.1
Mother's age >34 and birth order >3	0.1	*	5.5
Birth interval <24 months and birth order >3	1.2	(2.1)	0.6
Subtotal	1.5	(2.3)	6.3
In any avoidable high-risk category	24.6	1.0	22.8
Total	100.0	na	100.0
Number of births	2,421	na	5,619

Note: Risk ratio is the ratio of the proportion dead among births in a specific high-risk category to the proportion dead among births not in any high-risk category.

na = Not applicable

¹ Women are assigned to risk categories according to the status they would have at the birth of a child if they were to conceive at the time of the survey: current age less than 17 years and 3 months or greater than 34 years and 2 months, latest birth less than 15 months ago, or latest birth of order 3 or higher

² Includes the category age <18 and birth order >3

^a Includes sterilized women

() Based on 25-49 unweighted cases

* Based on fewer than 25 unweighted cases

Table 37 Antenatal care

Percent distribution of women who had a live birth in the five years preceding the survey by antenatal care (ANC) provider during pregnancy for the most recent live birth and percentage receiving ANC from a skilled provider, according to background characteristics, Telangana, 2015-16

Background characteristic	Doctor	ANM/ nurse/ midwife/ LHV	Community/ village health worker	Anganwadi/ ICDS worker	ASHA	Other	No one	Total	Number of women	Percentage receiving ANC from a skilled provider ¹
Mother's age at birth										
<20	84.8	11.2	0.0	2.7	0.4	0.0	0.9	100.0	236	95.9
20-34	86.1	8.5	0.1	2.0	0.2	0.2	2.9	100.0	1,536	94.6
Birth order										
1	86.9	8.2	0.0	2.5	0.6	0.2	1.6	100.0	609	95.1
2-3	85.6	8.9	0.1	2.0	0.1	0.2	3.2	100.0	1,114	94.5
4+	79.6	13.4	0.7	0.0	0.0	0.0	6.3	100.0	70	93.0
Residence										
Urban	88.9	5.7	0.1	2.1	0.1	0.3	2.8	100.0	841	94.6
Rural	83.1	11.6	0.1	2.1	0.4	0.1	2.7	100.0	952	94.7
Schooling										
No schooling	75.9	15.7	0.2	2.8	0.4	0.4	4.5	100.0	349	91.6
<5 years complete	83.5	13.1	0.0	1.4	0.0	0.0	2.0	100.0	82	96.6
5-9 years complete	85.8	8.3	0.1	2.2	0.2	0.4	3.1	100.0	391	94.1
10-11 years complete	88.1	5.9	0.0	2.1	0.5	0.1	3.3	100.0	419	94.0
12 or more years complete	90.8	6.4	0.0	1.7	0.0	0.0	1.1	100.0	552	97.2
Religion										
Hindu	85.4	9.0	0.1	2.0	0.2	0.2	3.0	100.0	1,549	94.5
Muslim	90.6	5.6	0.3	1.6	0.4	0.0	1.5	100.0	192	96.2
Christian	(79.9)	(15.3)	(0.0)	(3.8)	(0.0)	(0.0)	(1.0)	100.0	49	(95.1)
Caste/tribe										
Scheduled caste	83.8	10.6	0.3	3.0	0.2	0.0	2.1	100.0	334	94.5
Scheduled tribe	70.2	15.9	0.0	5.9	0.8	0.0	7.2	100.0	147	86.1
Other backward class	87.8	8.0	0.0	1.6	0.2	0.3	2.0	100.0	1,032	95.8
Other	89.2	5.8	0.0	0.7	0.0	0.0	4.2	100.0	271	95.0
Total	85.8	8.8	0.1	2.1	0.2	0.2	2.8	100.0	1,793	94.6

Note: If more than one source of ANC was mentioned, only the provider with the highest qualification is considered in this table. Total includes women with age at birth 35-49, women belonging to "other" religions, and women who don't know their caste/tribe, who are not shown separately.

ANM = Auxiliary nurse midwife; LHV = Lady health visitor; ICDS = Integrated Child Development Services; ASHA = Accredited Social Health Activist

¹ Skilled provider includes doctor, auxiliary nurse midwife, nurse, midwife, and lady health visitor

() Based on 25-49 unweighted cases

Table 38 Antenatal care services and information received

Percentage of women who had a live birth in the five years preceding the survey who received antenatal care (ANC) for the most recent live birth by services and information received, according to residence and source of antenatal care, Telangana, 2015-16

Services/information	Residence		Source of ANC			Total
	Urban	Rural	Public health sector	Private health sector ¹	Both public and private health sector ¹	
Percentage receiving selected services during antenatal care						
Weighed	100.0	98.6	99.1	99.6	100.0	99.3
Blood pressure measured	99.9	98.8	98.7	99.7	99.1	99.3
Urine sample taken	99.9	99.0	99.4	99.5	99.5	99.4
Blood sample taken	99.8	97.6	97.5	99.3	98.2	98.6
Abdomen examined	98.7	96.0	96.3	97.7	96.0	97.2
Percentage receiving information on specific pregnancy complications						
Vaginal bleeding	42.4	45.1	44.7	44.0	46.3	43.8
Convulsions	33.8	35.8	34.5	36.0	39.9	34.8
Prolonged labour	41.0	41.3	42.0	41.5	45.6	41.2
Severe abdominal pain	45.0	45.5	47.6	44.8	49.2	45.3
High blood pressure	45.7	45.0	45.2	46.6	51.0	45.3
Where to go if experienced pregnancy complications	56.1	58.3	59.3	58.8	73.2	57.3
Number of women	817	926	688	1,234	192	1,744

¹ Includes nongovernmental organizations or trust hospitals/clinics

Table 39 Antenatal care indicators

Among women with a live birth in the five years preceding the survey, percentage who received different types of antenatal care (ANC) during the pregnancy for their most recent live birth, by background characteristics, Telangana, 2015-16

Background characteristic	Percentage who had four or more ANC visits	Percentage with an ANC visit in the first trimester of pregnancy	Percentage who received two or more TT injections during the pregnancy	Percentage whose last live birth was protected against neonatal tetanus ¹	Percentage who were given or bought IFA	Percentage who took IFA for at least 100 days	Percentage who had full antenatal care ²	Percentage who took an intestinal parasite drug	Number of women
Age at birth									
<20	75.8	85.0	84.1	88.4	89.5	52.8	38.3	20.8	236
20-34	75.0	83.1	85.7	89.0	91.5	53.3	43.1	23.1	1,536
Birth order									
1	79.3	86.6	86.8	89.7	92.0	56.0	46.6	25.5	609
2-3	72.8	82.2	85.7	89.1	91.6	51.8	40.3	22.0	1,114
4+	72.3	66.5	74.7	82.2	76.2	40.9	32.6	20.3	70
Residence									
Urban	77.6	87.4	86.6	89.6	93.6	58.8	47.7	22.9	841
Rural	72.7	79.2	84.8	88.5	88.9	47.5	37.3	23.3	952
Schooling									
No schooling	64.1	73.5	78.2	83.3	86.9	39.1	26.5	22.0	349
<5 years complete	73.7	69.4	85.1	90.7	87.0	45.2	35.6	16.9	82
5-9 years complete	76.0	79.6	84.3	88.3	90.2	50.6	39.1	19.6	391
10-11 years complete	73.6	85.5	86.8	89.8	90.2	51.8	40.9	26.0	419
12 or more years complete	82.3	91.8	90.6	92.5	95.8	64.9	56.2	25.0	552
Religion									
Hindu	74.2	82.5	85.7	88.9	91.4	52.2	41.7	22.4	1,549
Muslim	83.7	87.5	88.8	93.6	87.2	55.0	46.6	27.2	192
Christian	(66.3)	(84.0)	(74.5)	(75.9)	(95.8)	(63.3)	(40.4)	(27.5)	49
Caste/tribe									
Scheduled caste	72.3	81.2	84.5	88.2	91.0	54.9	42.3	22.0	334
Scheduled tribe	68.6	73.1	77.6	81.5	86.2	42.0	32.2	19.4	147
Other backward class	75.7	84.3	85.9	89.4	91.4	51.9	40.6	22.5	1,032
Other	79.3	86.5	90.2	92.7	92.9	60.6	54.1	28.0	271
Total	75.0	83.1	85.7	89.0	91.1	52.8	42.2	23.1	1,793

Note: Total includes women with age at birth 35-49, women belonging to "other" religions, and women who don't know their caste/tribe, who are not shown separately.

TT = Tetanus toxoid; IFA = Iron and folic acid

¹ Includes mothers with two injections during the pregnancy for her last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth

² Full antenatal care includes having received at least four antenatal care visits, having received at least one TT injection, and having taken IFA tablets or syrup for 100 or more days

() Based on 25-49 unweighted cases

Table 40 Antenatal care indicators by district

Among women with a live birth in the five years preceding the survey, percentage who received different types of antenatal care (ANC) during the pregnancy for their most recent live birth, and among women with registered pregnancies, percentage who received a Mother and Child Protection Card (MCP Card), by district, Telangana, 2015-16

District	Percentage who had four or more ANC visits	Percentage with an ANC visit in the first trimester of pregnancy	Percentage who took IFA for at least 100 days	Percentage who received two or more TT injections during the pregnancy	Percentage who had full antenatal care ¹	Number of women	Percentage of women who received an MCP Card	Number of registered pregnancies
Adilabad	68.3	72.6	41.6	76.5	30.6	129	82.8	116
Hyderabad	85.8	90.4	73.1	92.3	62.6	269	82.3	222
Karimnagar	73.7	83.4	47.3	90.2	32.8	172	83.6	162
Khammam	79.2	83.9	56.4	91.6	45.1	131	93.3	130
Mahbubnagar	60.2	82.1	48.4	87.1	36.0	188	90.2	175
Medak	68.3	79.2	36.4	76.8	31.0	143	95.7	140
Nalgonda	72.8	77.8	36.5	81.5	29.3	168	91.5	162
Nizamabad	71.2	70.5	49.6	81.8	35.7	117	94.9	114
Rangareddy	77.2	88.9	53.8	84.5	43.0	335	91.2	288
Warangal	84.7	85.5	70.4	88.1	60.7	140	88.7	137
Telangana	75.0	83.1	52.8	85.7	42.2	1,793	89.2	1,646

TT = Tetanus toxoid; IFA = Iron and folic acid

¹ Full antenatal care includes having received at least four antenatal care visits, having received at least one TT injection, and having taken IFA tablets or syrup for 100 or more days

Table 41 Advice received during pregnancy

Among women with a live birth in the five years preceding the survey who met with a community health worker in the last three months of pregnancy for their most recent live birth, percentage who received different types of advice, by background characteristics, Telangana, 2015-16

Background characteristic	Percentage who received advice on:					Number of women who met with a community health worker in the last three months of pregnancy ¹
	Importance of institutional delivery	Cord care	Breastfeeding	Keeping the baby warm	Family planning	
Age at birth						
<20	81.3	58.5	90.5	85.6	60.4	151
20-34	79.6	69.5	92.9	87.4	65.0	798
Birth order						
1	79.8	67.8	92.5	87.3	58.1	340
2-3	79.5	67.9	92.5	87.0	67.2	587
4+	(90.0)	(66.9)	(94.7)	(90.1)	(77.7)	29
Residence						
Urban	77.5	70.6	93.1	85.7	69.2	302
Rural	81.0	66.5	92.3	87.9	62.0	654
Schooling						
No schooling	82.3	66.5	91.8	85.2	61.3	202
<5 years complete	80.0	69.2	92.0	85.7	56.9	41
5-9 years complete	76.7	67.2	91.5	86.7	63.0	231
10-11 years complete	79.9	64.7	90.6	83.9	61.8	256
12 or more years complete	81.0	73.0	96.6	93.5	72.5	225
Religion						
Hindu	79.8	68.8	92.9	87.6	64.3	862
Muslim	77.7	56.2	87.6	78.6	58.2	68
Christian	(86.6)	(62.1)	(94.7)	(94.7)	(77.9)	24
Caste/tribe						
Scheduled caste	83.8	71.9	93.1	87.4	68.0	197
Scheduled tribe	76.2	68.9	89.2	86.4	51.6	80
Other backward class	80.2	67.4	92.8	86.7	64.6	569
Other	74.9	62.6	92.7	89.4	65.2	105
Total	79.9	67.8	92.6	87.2	64.3	956

Note: Total includes women with age at birth 35-49, women belonging to "other" religions, and women who don't know their caste/tribe, who are not shown separately.

¹ Community health worker includes auxiliary nurse midwife (ANM), lady health visitor (LHV), Accredited Social Health Activist (ASHA), *anganwadi* worker, and other community health worker

() Based on 25-49 unweighted cases

Table 42 Pregnancies for which an ultrasound test was done

Percentage of all pregnancies in the five years preceding the survey for which an ultrasound test was done and percent distribution of pregnancies with an ultrasound test by pregnancy outcome, according to background characteristics, Telangana, 2015-16

Background characteristic	Percentage of pregnancies with an ultrasound test	Number of pregnancies	Pregnancy outcome ¹				Total percent	Number of pregnancies with an ultrasound test
			Son	Daughter	Termination	Still pregnant		
Mother's age at pregnancy								
<20	90.7	622	48.6	40.4	5.0	6.0	100.0	564
20-34	91.7	2,282	45.0	40.6	7.3	7.2	100.0	2,093
35-49	(80.0)	29	*	*	*	*	100.0	23
Residence								
Urban	92.0	1,413	44.4	40.1	8.3	7.2	100.0	1,300
Rural	90.8	1,520	46.8	41.0	5.6	6.6	100.0	1,381
Antenatal care visits²								
None	82.1	55	(57.5)	(32.9)	(5.5)	(4.1)	100.0	45
1-3	90.7	475	46.2	41.0	5.3	7.5	100.0	431
4+	92.6	1,635	45.5	41.1	7.5	6.0	100.0	1,513
Mother's schooling								
No schooling	87.9	551	46.3	42.5	6.3	4.9	100.0	484
<5 years complete	90.7	126	52.3	36.3	7.3	4.2	100.0	114
5-9 years complete	91.3	665	47.1	40.7	4.9	7.4	100.0	607
10-11 years complete	91.6	690	46.9	39.0	6.7	7.4	100.0	632
12 or more years complete	93.6	902	42.4	41.0	9.0	7.7	100.0	844
Religion								
Hindu	91.8	2,488	45.6	40.3	7.1	7.0	100.0	2,284
Muslim	90.9	348	48.5	39.9	4.8	6.8	100.0	316
Christian	81.9	94	33.4	50.0	12.0	4.6	100.0	77
Caste/tribe								
Scheduled caste	87.8	555	50.7	36.6	6.9	5.8	100.0	488
Scheduled tribe	85.9	242	46.5	40.3	5.1	8.1	100.0	208
Other backward class	92.8	1,652	44.4	42.5	5.4	7.7	100.0	1,532
Other	93.8	470	43.8	38.0	13.3	5.0	100.0	441

Continued...

Table 42 Pregnancies for which an ultrasound test was done—Continued

Percentage of all pregnancies in the five years preceding the survey for which an ultrasound test was done and percent distribution of pregnancies with an ultrasound test by pregnancy outcome, according to background characteristics, Telangana, 2015-16

Background characteristic	Percentage of pregnancies with an ultrasound test	Number of pregnancies	Pregnancy outcome ¹				Total percent	Number of pregnancies with an ultrasound test
			Son	Daughter	Termination	Still pregnant		
Mother's number of living children at time of pregnancy								
No children	92.9	1,325	45.9	39.1	7.5	7.6	100.0	1,230
1 child	91.3	1,224	45.2	42.6	6.0	6.1	100.0	1,118
0 sons	91.6	595	45.6	42.4	7.6	4.5	100.0	544
1 son	91.1	629	45.0	42.8	4.6	7.6	100.0	573
2 children	87.3	312	46.4	38.7	7.7	7.1	100.0	273
0 sons	89.1	131	49.7	33.3	8.7	8.3	100.0	117
1 or more sons	86.0	181	44.0	42.8	7.0	6.2	100.0	156
3 children	80.4	56	(44.0)	(45.2)	(4.8)	(6.1)	100.0	45
1 or more sons	(77.3)	34	(42.6)	(52.7)	(4.7)	(0.0)	100.0	26
Total	91.4	2,933	45.6	40.5	6.9	6.9	100.0	2,681

Note: Total includes pregnancies of women belonging to "other" religions, women who don't know the number of antenatal care visits, women who don't know their caste/tribe, women who have 3 children and no sons, and women who have 4 or more children, which are not shown separately.

¹ For multiple births, sex of pregnancy outcome is the sex of the first listed birth

² Includes only the most recent pregnancy in the five years preceding the survey

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

Table 43 Pregnancy registration and Mother and Child Protection Card

Percentage of women who had a live birth in the five years preceding the survey who registered the pregnancy for the most recent live birth; and among registered pregnancies, the percentage by the timing of the registration and the percentage who received a Mother and Child Protection Card (MCP Card), by background characteristics, Telangana, 2015-16

Background characteristic	Percentage of pregnancies that were registered	Number of pregnancies	Timing of pregnancy registration		Percentage of mothers given an MCP card	Number of registered pregnancies
			First trimester	Later		
Mother's age at birth						
<20	92.7	236	78.0	22.0	95.5	219
20-34	91.6	1,536	83.3	16.6	88.2	1,407
Birth order						
1	91.0	609	86.8	13.2	90.0	554
2-3	92.0	1,114	80.9	18.9	89.2	1,026
4+	93.9	70	70.0	30.0	82.0	66
Residence						
Urban	86.6	841	85.2	14.8	85.4	728
Rural	96.3	952	80.3	19.5	92.2	917
Schooling						
No schooling	93.8	349	77.8	22.0	92.3	327
<5 years complete	91.1	82	79.1	20.9	87.8	75
5-9 years complete	93.7	391	81.0	18.8	89.6	367
10-11 years complete	93.9	419	82.7	17.3	89.3	393
12 or more years complete	87.6	552	87.2	12.8	86.8	483
Religion						
Hindu	92.1	1,549	82.0	17.9	89.6	1,426
Muslim	88.3	192	87.6	12.4	86.2	170
Christian	(93.8)	49	(80.7)	(19.3)	(86.4)	46
Caste/tribe						
Scheduled caste	91.8	334	81.0	19.0	93.5	307
Scheduled tribe	93.3	147	83.8	15.7	85.7	137
Other backward class	93.1	1,032	82.2	17.7	88.7	961
Other	86.0	271	85.2	14.8	87.6	233
Total	91.8	1,793	82.5	17.4	89.2	1,646

Note: Total includes pregnancies of women with age at birth 35-49, women belonging to "other" religions, and women who don't know their caste/tribe, which are not shown separately.

() Based on 25-49 unweighted cases

Table 44 Delivery and postnatal care

Percent distribution of live births in the five years preceding the survey by place of delivery and assistance during delivery, percentage delivered by a skilled provider and by caesarean section, percentage of live births delivered at home that were delivered by a skilled provider and percentage for which the delivery protocol was followed, and percent distribution of women giving birth in the five years preceding the survey by timing and type of provider of the first postnatal check of the mother for their most recent live birth, by residence, Telangana, 2015-16

Delivery and postnatal care descriptors	Urban	Rural	Total
Place of delivery			
Health facility	96.3	87.3	91.5
Public sector	27.1	34.4	31.0
NGO/trust	0.5	0.4	0.4
Private sector	68.6	52.6	60.1
At home	3.6	11.9	8.0
Own home	1.8	7.2	4.7
Parent's home	1.7	4.5	3.1
Other home	0.1	0.2	0.1
Other ¹	0.1	0.8	0.5
Total	100.0	100.0	100.0
Assistance during delivery²			
Doctor	91.0	80.1	85.2
ANM/nurse/midwife/LHV	3.8	7.3	5.6
Other health personnel	0.0	1.1	0.6
Dai (TBA)	1.6	4.9	3.4
Friends/relatives	3.1	5.9	4.6
Other	0.4	0.6	0.5
No one	0.1	0.2	0.1
Total	100.0	100.0	100.0
Percentage of all births delivered by a skilled provider ³	94.7	88.4	91.4
Percentage of home births delivered by a skilled provider ³	1.7	4.0	2.9
Percentage delivered by caesarean section	63.2	53.4	58.0
Percentage delivered by emergency caesarean section ⁴	26.5	20.3	23.2
Number of births	1,129	1,282	2,411
For home deliveries⁵			
Disposable delivery kit (DDK) used	*	40.3	41.5
Clean blade used to cut the cord	*	95.5	96.6
Either of the above	*	96.2	97.1
Baby was immediately wiped dry and then wrapped without being bathed	*	80.9	80.0
Number of births delivered at home	30	100	130
Timing after delivery of mother's first postnatal check⁶			
Less than 4 hours	75.0	70.1	72.4
4-23 hours	2.1	3.8	3.0
1-2 days	8.1	6.5	7.3
3-41 days	2.4	4.8	3.7
Don't know ¹	0.9	0.5	0.7
No postnatal check	11.5	14.1	12.9
Total	100.0	100.0	100.0

Continued...

Table 44 Delivery and postnatal care—Continued

Percent distribution of live births in the five years preceding the survey by place of delivery and assistance during delivery, percentage delivered by a skilled provider and by caesarean section, percentage of live births delivered at home that were delivered by a skilled provider and percentage for which the delivery protocol was followed, and percent distribution of women giving birth in the five years preceding the survey by timing and type of provider of the first postnatal check of the mother for their most recent live birth, by residence, Telangana, 2015-16

Delivery and postnatal care descriptors	Urban	Rural	Total
Type of provider of mother's first postnatal check⁶			
Doctor	81.6	73.3	77.2
ANM/nurse/midwife/LHV	6.4	10.4	8.5
Other health personnel	0.1	0.3	0.2
<i>Dai</i> (TBA)	0.0	1.2	0.6
Other	0.4	0.6	0.5
No postnatal check	11.5	14.1	12.9
Total	100.0	100.0	100.0
Number of last live births	841	952	1,793

NGO = Nongovernmental organization; ANM = Auxiliary nurse midwife; LHV = Lady health visitor; TBA = Traditional birth attendant

¹ Includes missing

² If the respondent mentioned more than one person attending during delivery, only the most qualified person is considered in this table

³ Skilled provider includes doctor, auxiliary nurse midwife, nurse, midwife, lady health visitor, and other health personnel

⁴ A caesarean section for which the decision to have the operation was made after the onset of labour

⁵ Based on the last live birth in the five years preceding the survey

⁶ Based on the last live birth in the five years preceding the survey. Postnatal checks are checks on the woman's health within 42 days of the birth

* Percentage not shown; based on fewer than 25 unweighted cases

Table 45 Delivery and postnatal care by background characteristics

Percentage of live births in the five years preceding the survey delivered in a health facility and percentage delivered with assistance from health personnel, and percentage of women who had a live birth in the five years preceding the survey who received a postnatal check and who received a postnatal check within two days of birth for their most recent birth, by background characteristics, Telangana, 2015-16

Background characteristic	Percentage of births delivered in a public health facility	Percentage of births delivered in a private health facility	Percentage of births delivered in a health facility	Percentage of deliveries assisted by health personnel ¹	Number of births	Percentage of women with a postnatal check ^{2,3}	Percentage of women with a postnatal check within two days of birth ^{2,3}	Number of women
Mother's age at birth								
<20	39.5	52.7	92.2	89.5	384	87.5	84.3	236
20-34	29.6	61.9	91.5	91.8	2,003	87.1	82.5	1,536
35-49	(12.4)	(70.7)	(83.1)	(85.0)	23	*	*	22
Residence								
Urban	27.1	69.2	96.3	94.7	1,129	88.5	85.2	841
Rural	34.4	53.0	87.3	88.4	1,282	85.9	80.5	952
Birth order								
1	27.1	67.4	94.5	93.5	1,056	89.6	85.6	609
2-3	34.1	56.0	90.1	90.2	1,273	86.1	82.3	1,114
4+	32.5	43.7	76.2	82.5	81	80.9	64.0	70
Antenatal care visits²								
None	(20.6)	(48.6)	(69.2)	(73.8)	50	(62.4)	(55.4)	50
1-3	32.8	57.9	90.7	87.0	393	86.5	82.0	393
4+	30.8	63.4	94.2	93.9	1,344	88.4	84.2	1,344
Mother's schooling								
No schooling	40.3	35.4	75.7	80.8	473	81.3	73.3	349
<5 years complete	42.3	44.3	86.6	88.9	109	83.8	79.5	82
5-9 years complete	40.3	53.6	93.9	92.6	556	85.9	81.1	391
10-11 years complete	30.0	64.9	94.9	93.9	558	89.4	87.2	419
12 or more years complete	16.6	81.7	98.3	95.8	716	90.3	86.9	552
Religion								
Hindu	30.1	61.2	91.4	91.1	2,046	87.2	83.2	1,549
Muslim	35.5	58.1	93.6	92.8	294	84.2	79.5	192
Christian	34.9	52.6	87.5	91.5	67	(94.1)	(81.4)	49
Caste/tribe								
Scheduled caste	41.6	48.0	89.6	89.3	446	86.4	80.6	334
Scheduled tribe	32.7	46.9	79.7	85.3	200	84.6	75.7	147
Other backward class	28.4	64.6	92.9	91.8	1,382	89.4	85.2	1,032
Other	25.8	69.3	95.2	95.6	370	80.7	79.6	271
Place of delivery								
Public health facility	100.0	0.0	100.0	95.6	747	90.0	87.4	554
Private health facility ⁴	0.0	100.0	100.0	96.5	1,460	90.4	86.4	1,105
Home	na	na	na	36.2	193	47.1	33.0	130
Total	31.0	60.6	91.5	91.4	2,411	87.1	82.7	1,793

Note: Total includes information on births to women who don't know the number of antenatal care visits, births belonging to "other" religions, births whose caste/tribe is not known, and births delivered in an "other" place of delivery, which is not shown separately.

na = Not applicable

¹ Health personnel includes doctor, auxiliary nurse midwife, nurse, midwife, lady health visitor, and other health personnel. If the respondent mentioned more than one person attending during delivery, only the most qualified person is considered in this table.

² Based on the last live birth in the five years preceding the survey

³ Postnatal checks are checks on the woman's health within 42 days of the birth

⁴ Includes nongovernmental organizations or trust hospitals/clinics

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

Table 46 Delivery and postnatal care by district

Percentage of live births in the five years preceding the survey delivered in a health facility, percentage delivered with assistance from health personnel and percentage delivered by caesarean section, and percentage of women who had a live birth in the five years preceding the survey who received a postnatal check within two days of birth for their most recent birth, by district, Telangana, 2015-16

District	Percentage of births delivered in a public health facility	Percentage of births delivered in a private health facility	Percentage of births delivered in a health facility	Percentage of deliveries assisted by health personnel ¹	Percentage of births delivered by caesarean section	Number of births	Percentage of women with a postnatal check within two days of birth ²	Number of women
Adilabad	37.6	33.2	70.8	78.0	39.9	177	59.9	129
Hyderabad	23.9	73.3	97.2	99.9	62.2	374	79.0	269
Karimnagar	17.3	79.7	97.0	98.6	81.0	218	92.2	172
Khammam	32.4	61.8	94.2	96.5	65.6	183	86.0	131
Mahbubnagar	35.0	43.4	78.4	86.0	33.5	261	72.9	188
Medak	38.1	53.1	91.3	87.3	47.1	191	83.9	143
Nalgonda	25.8	69.0	94.8	93.4	69.0	226	92.7	168
Nizamabad	36.4	56.2	92.6	76.0	50.3	164	89.7	117
Rangareddy	35.7	59.5	95.3	91.6	58.7	433	81.7	335
Warangal	31.0	65.0	96.0	95.9	69.2	184	92.7	140
Telangana	31.0	60.6	91.5	91.4	58.0	2,411	82.7	1,793

¹ Health personnel includes doctor, auxiliary nurse midwife, nurse, midwife, lady health visitor, and other health personnel. If the respondent mentioned more than one person attending during delivery, only the most qualified person is considered in this table.

² Postnatal checks are checks on the woman's health within 42 days of the birth

Table 47 Delivery costs and financial assistance

The average out-of-pocket cost paid for delivery for the most recent live birth among women who had a live birth in the five years preceding the survey that was delivered in a health facility by type of facility, and among women who had a live birth in the five years preceding the survey for the most recent birth that was delivered in a health facility, the percentage who received financial assistance under *Janani Suraksha Yojana* (JSY), according to background characteristics, Telangana, 2015-16

Background characteristic	Average cost (Rs.) ¹			Percentage who received financial assistance under JSY	Number of births in a health facility
	Public health facility	Private health facility	Any health facility		
Mother's age at birth					
<20	4,861	14,621	10,174	15.3	308
20-34	4,018	16,192	12,391	11.5	1,333
Birth order					
1	4,436	16,183	12,575	12.4	994
2-3	4,024	16,019	11,514	11.7	633
4+	(3,359)	(10,423)	(6,796)	(11.1)	32
Residence					
Urban	4,145	15,876	12,437	10.2	811
Rural	4,289	16,211	11,706	14.0	848
Schooling					
No schooling	3,369	16,628	9,863	17.1	272
<5 years complete	2,446	22,496	12,191	17.1	72
5-9 years complete	3,587	14,935	9,883	16.3	374
10-11 years complete	4,001	15,709	11,757	10.8	398
12 or more years complete	8,298	16,057	14,902	7.1	543
Religion					
Hindu	4,276	16,293	12,326	12.0	1,432
Muslim	3,376	14,743	10,507	11.6	179
Christian	(6,361)	(12,959)	(10,255)	(17.4)	44
Caste/tribe					
Scheduled caste	4,975	15,722	10,476	21.2	299
Scheduled tribe	3,336	17,989	11,794	6.8	119
Other backward class	3,742	16,079	12,311	10.4	973
Other	5,803	15,621	13,378	10.0	259
Total	4,229	16,040	12,056	12.1	1,659

Note: Total includes information on births with mother's age at birth 35-49, births belonging to "other" religions, and births whose caste/tribe is not known, which is not shown separately.

¹ Excludes women who don't know the cost

() Based on 25-49 unweighted cases

Table 48 Birth order and delivery characteristics by district

Percentage of births during the three years preceding the survey of birth order 3 or more, percentage of women who had a live birth in the five years preceding the survey by antenatal care (ANC) provider during pregnancy for the most recent live birth and among women who had a live birth in the five years preceding the survey for the most recent birth that was delivered in a health facility, the percentage of women who received financial assistance under *Janani Suraksha Yojana* (JSY) by district, Telangana, 2015-16

District	Percentage of births of birth order 3 or more	Number of births	Percentage of last births receiving antenatal care from doctor	Percentage of last births receiving antenatal care from ANM/nurse/midwife/LHV	Number of last births	Percentage of births in a health facility receiving financial assistance under JSY	Number of births in a health facility
Adilabad	19.0	106	68.5	35.9	129	8.8	95
Hyderabad	16.7	227	95.1	13.7	269	9.3	263
Karimnagar	11.3	113	90.7	17.9	172	11.4	168
Khammam	7.9	109	86.0	33.0	131	16.1	125
Mahbubnagar	26.7	169	81.3	25.4	188	14.4	153
Medak	20.3	107	79.9	25.8	143	15.7	131
Nalgonda	14.3	127	84.7	23.0	168	10.1	161
Nizamabad	21.2	100	78.2	33.8	117	17.2	111
Rangareddy	6.6	237	87.3	18.5	335	12.2	316
Warangal	11.6	100	94.2	19.5	140	9.6	136
Telangana	15.3	1,396	85.8	22.8	1,793	12.1	1,659

ANM = Auxiliary nurse midwife; LHV = Lady health visitor

Table 49 Timing of first health check after birth for the newborn

Percent distribution of last births in the five years preceding the survey by time after birth of first health check, and the percentage of births with a health check in the first two days after birth, according to background characteristics, Telangana, 2015-16

Background characteristic	Time after birth of newborn's first health check							Total	Percentage of births with a health check in the first two days after birth	Number of births
	Less than 1 hour	1-3 hours	4-23 hours	1-2 days	3-6 days	Don't know	No health check ¹			
Mother's age at birth										
<20	8.2	16.2	0.8	3.7	1.5	0.2	69.5	100.0	28.8	333
20-34	6.4	15.7	1.0	2.9	0.9	0.2	72.9	100.0	26.0	1,439
Birth order										
1	8.0	18.1	0.8	3.3	0.7	0.3	68.8	100.0	30.2	609
2-3	6.0	14.5	1.1	2.8	1.2	0.1	74.4	100.0	24.4	1,114
4-5	7.9	14.3	0.0	4.3	2.2	1.9	69.3	100.0	26.6	66
Place of delivery										
Public health facility	7.2	15.2	0.7	1.9	1.0	0.1	73.8	100.0	25.0	554
Private health facility ²	6.8	16.4	0.9	2.8	0.8	0.2	72.2	100.0	26.8	1,105
Home	5.1	11.8	3.0	8.8	3.5	1.1	66.8	100.0	28.6	130
Residence										
Urban	6.8	11.9	0.3	2.7	0.4	0.0	77.9	100.0	21.7	841
Rural	6.8	19.0	1.5	3.3	1.6	0.4	67.4	100.0	30.6	952
Religion										
Hindu	6.9	16.2	0.9	3.0	1.2	0.2	71.5	100.0	27.1	1,549
Muslim	3.9	14.0	1.0	2.7	0.0	0.0	78.4	100.0	21.6	192
Christian	(12.9)	(6.7)	(2.7)	(3.3)	(0.8)	(0.0)	(73.6)	100.0	(25.6)	49
Caste/tribe										
Scheduled caste	5.9	15.8	1.3	2.4	1.4	0.6	72.7	100.0	25.4	334
Scheduled tribe	5.1	14.7	0.6	2.3	2.2	0.9	74.1	100.0	22.8	147
Other backward class	8.0	16.6	0.8	3.7	0.9	0.1	69.9	100.0	29.1	1,032
Other	4.1	13.0	1.3	1.6	0.5	0.0	79.6	100.0	19.9	271
Mother's schooling										
No schooling	5.8	14.7	1.4	4.0	1.6	0.2	72.3	100.0	25.9	349
<5 years complete	6.7	20.3	0.9	0.7	1.8	0.7	68.8	100.0	28.7	82
5-9 years complete	5.1	17.8	0.7	2.4	1.3	0.2	72.5	100.0	26.0	391
10-11 years complete	8.8	16.2	1.7	3.5	1.2	0.0	68.5	100.0	30.2	419
12 or more years complete	7.0	13.8	0.3	2.7	0.3	0.3	75.6	100.0	23.8	552
Total	6.8	15.7	1.0	3.0	1.0	0.2	72.3	100.0	26.4	1,793

Note: Total includes births of birth order 6 or more, births with mother's age at birth 35-49, births belonging to "other" religions, and births whose caste/tribe is not known, which are not shown separately.

¹ Includes newborns who received a health check after the first week

² Includes nongovernmental organizations or trust hospitals/clinics

() Based on 25-49 unweighted cases

Table 51 Male involvement in maternal care: Men's report

Among men age 15-49 whose youngest living child was age 0-35 months, percentage for whom the youngest child's mother received antenatal care, percentage who were present for at least one antenatal check-up, percentage who were told by a health provider or worker at any time during the pregnancy about specific signs of pregnancy complications, percentage to whom a health provider or worker spoke about specific aspects of maternal care at any time during the pregnancy, and percentage whose youngest child was delivered in a health facility, percentage who were given specific home delivery related information, by residence, Telangana, 2015-16

Antenatal and delivery care information	Urban	Rural	Total
Percentage of men for whom the youngest child's mother received antenatal care	95.6	94.2	94.8
Percentage of men who were present at any antenatal check-up	71.5	67.0	69.0
Percentage who were told by a health provider or worker about the following signs of pregnancy complications:			
Vaginal bleeding	41.5	20.9	30.3
Convulsions	30.9	18.6	24.2
Prolonged labour	37.1	25.9	31.0
Severe abdominal pain	54.2	31.2	41.6
High blood pressure	35.4	30.8	32.9
Percentage ever told what to do if the mother had any pregnancy complications	35.0	31.3	33.0
Percentage whose youngest child age 0-35 months was delivered in a health facility	100.0	87.8	93.4
Percentage to whom a health provider or worker spoke about the following aspects of maternal care:			
The importance of delivering in a health facility	60.3	47.7	53.4
The importance of proper nutrition for the mother during pregnancy	77.0	82.0	79.7
Family planning or delaying his next child	55.7	48.9	52.0
Number of men with a youngest child age 0-35 months	83	100	184

Table 52 Vaccinations by background characteristics

Percentage of children age 12-23 months who received specific vaccines at any time before the survey (according to a vaccination card or the mother's report) and percentage with a vaccination card seen by the interviewer, by background characteristics, Telangana, 2015-16

Background characteristic	Hepatitis B ¹			DPT			Polio ¹			Measles vaccinations ³	All basic vaccinations ²	No vaccination card seen	Percentage with a vaccination card seen	Number of children				
	0	1	2	3	1	2	3	0	1						2	3		
Sex																		
Male	89.1	94.8	88.7	75.2	98.1	96.2	95.0	87.3	95.5	97.1	92.6	78.7	93.1	72.0	1.0	71.6	261	
Female	87.3	92.8	85.2	65.3	96.6	94.9	92.9	88.6	95.0	94.6	89.3	71.6	87.8	63.6	1.3	65.4	234	
Birth order																		
1	89.6	93.1	85.5	70.8	96.0	95.5	93.0	86.9	95.5	94.8	89.6	74.7	88.8	67.5	1.5	68.3	218	
2-3	86.9	94.1	87.6	70.7	98.4	95.5	94.4	88.3	94.7	96.8	92.2	77.0	91.8	69.5	0.9	69.7	263	
Residence																		
Urban	92.3	95.1	86.3	69.7	97.6	94.6	92.9	89.5	97.9	98.0	93.6	74.4	91.8	67.8	0.0	67.9	246	
Rural	84.2	92.6	87.8	71.4	97.2	96.6	95.0	86.3	92.6	93.9	88.5	76.3	89.4	68.3	2.2	69.4	249	
Mother's schooling																		
No schooling	80.5	87.7	81.7	61.8	92.4	96.2	95.8	82.1	90.6	91.8	87.7	69.2	82.3	61.2	1.9	63.7	78	
5-9 years complete	79.4	92.8	88.9	74.1	97.8	96.0	93.5	89.1	94.3	96.5	91.9	77.0	91.8	70.3	1.5	66.3	102	
10-11 years complete	93.7	97.1	86.8	71.2	100.0	96.1	93.6	87.4	97.6	98.0	91.6	76.4	90.9	66.9	0.0	69.8	132	
12 or more years complete	92.1	94.4	87.5	70.3	97.2	94.1	92.9	89.7	95.6	95.3	90.6	74.6	94.0	69.9	1.6	70.8	164	
Religion																		
Hindu	88.9	94.9	88.2	71.4	98.1	95.4	94.1	88.0	95.3	95.9	91.6	76.2	91.9	69.3	1.1	70.1	430	
Muslim	82.3	92.5	87.5	77.1	98.5	96.1	94.6	90.7	93.1	97.4	92.0	79.6	87.5	68.4	1.5	64.0	49	
Caste/tribe																		
Scheduled caste	88.2	88.7	84.0	70.4	92.4	96.1	94.1	83.8	92.2	92.9	85.2	71.9	85.4	63.5	3.1	69.0	100	
Scheduled tribe	(70.6)	(93.6)	(91.9)	(75.6)	(97.8)	(100.0)	(100.0)	(89.0)	(90.3)	(100.0)	(97.2)	(82.2)	(93.0)	(78.0)	(0.0)	(65.4)	31	
Other backward class	89.0	95.0	86.4	68.1	98.7	94.1	92.6	87.7	96.2	95.8	91.4	73.7	91.9	65.9	0.6	65.8	285	
Other	93.1	96.3	91.3	78.8	99.0	98.6	96.5	93.6	97.4	98.3	95.1	83.6	93.3	78.9	1.0	79.0	75	
Total	88.2	93.8	87.1	70.6	97.4	95.6	94.0	87.9	95.3	95.9	91.1	75.3	90.6	68.1	1.1	68.7	495	

Note: Total includes children of birth order 4 or more, children whose mothers have less than 5 years of schooling, children belonging to Christian or "other" religions, and children whose caste/tribe is not known, who are not shown separately.

na = Not available

¹ Polio 0 is the polio vaccination given at birth and hepatitis B0 is the hepatitis vaccination given at birth

² Fully vaccinated with BCG, measles, and three doses each of DPT and polio vaccine (excluding polio vaccine given at birth)

³ Child has not received any vaccinations listed in the table

() Based on 25-49 unweighted cases

Table 53 Selected vaccinations by district

Percentage of children age 12-23 months who received specific vaccines at any time before the survey (according to a vaccination card or the mother's report) and percentage who received most vaccinations in a public health facility, by district, Telangana, 2015-16

District	3 doses of Hepatitis B	BCG	3 doses of DPT	3 doses of polio	Measles	All basic vaccinations ¹	No vaccinations ²	Number of children	Percentage who received most vaccinations in a public facility	Number of children who received any vaccinations
Adilabad	(71.4)	(88.6)	(81.5)	(80.4)	(84.3)	(70.0)	(4.3)	33	(90.8)	32
Hyderabad	77.6	100.0	89.9	78.9	91.0	71.3	0.0	78	54.3	78
Karimnagar	(83.7)	(100.0)	(92.0)	(89.4)	(94.9)	(84.2)	(0.0)	39	(82.2)	39
Khammam	(69.5)	(100.0)	(83.8)	(72.2)	(90.3)	(62.4)	(0.0)	34	(97.6)	34
Mahbubnagar	46.1	100.0	85.9	55.3	85.3	45.0	0.0	47	95.0	47
Medak	(79.5)	(96.8)	(89.7)	(89.5)	(93.7)	(81.4)	(0.0)	36	(100.0)	36
Nalgonda	(64.1)	(95.9)	(78.9)	(78.4)	(86.5)	(69.0)	(4.1)	41	(95.6)	40
Nizamabad	74.5	100.0	89.7	73.3	89.4	64.2	0.0	41	95.0	41
Rangareddy	69.2	94.9	93.0	74.0	91.3	68.1	1.5	105	76.7	104
Warangal	(71.4)	(97.8)	(84.0)	(66.9)	(97.8)	(66.9)	(2.2)	40	(92.3)	39
Telangana	70.6	97.4	87.9	75.3	90.6	68.1	1.1	495	83.7	489

¹ Fully vaccinated with BCG, measles, and three doses each of DPT and polio vaccine (excluding polio vaccine given at birth)

² Child has not received any vaccinations listed in the table

() Based on 25-49 unweighted cases

Table 54 Prevalence and treatment of symptoms of ARI and fever

Among children under age five, percentage who had symptoms of acute respiratory infection (ARI) and fever in the two weeks preceding the survey and percentage with symptoms of ARI and fever who received specific treatments, according to background characteristics, Telangana, 2015-16

Background characteristic	Children under age five			Children under age five with symptoms of ARI			Children under age five with fever	
	Percentage with symptoms of ARI ¹	Percentage with fever	Number of children	Percentage for whom treatment was sought from a health facility or provider ²	Percentage who received antibiotics	Number of children	Percentage for whom treatment was sought from a health facility or provider ²	Number of children
Age in months								
<6	2.7	10.2	188	*	*	5	*	19
6-11	1.5	19.9	232	*	*	3	(84.6)	46
12-23	2.7	21.7	495	*	*	14	73.7	108
24-35	1.9	18.0	448	*	*	8	77.6	81
36-47	2.0	13.8	510	*	*	10	71.3	71
48-59	1.5	14.3	474	*	*	7	73.8	68
Sex								
Male	1.8	16.3	1,252	*	*	22	74.6	204
Female	2.4	17.1	1,094	*	*	26	76.1	188
Residence								
Urban	2.2	14.6	1,106	*	*	24	80.9	161
Rural	1.9	18.6	1,241	(84.8)	(16.2)	24	71.4	231
Mother's schooling								
No schooling	0.9	15.0	450	*	*	4	78.2	68
<5 years complete	2.2	28.7	104	*	*	2	(70.2)	30
5-9 years complete	1.4	15.2	538	*	*	7	66.5	82
10-11 years complete	4.7	20.4	545	*	*	26	73.4	111
12 or more years complete	1.2	14.3	710	*	*	9	84.2	102
Religion								
Hindu	2.0	16.7	1,988	(89.8)	(38.1)	40	74.3	333
Muslim	1.6	15.5	288	*	*	5	(83.6)	45
Christian	5.0	18.6	67	*	*	3	*	13
Caste/tribe								
Scheduled caste	1.7	15.8	432	*	*	7	72.6	68
Scheduled tribe	2.6	22.2	191	*	*	5	59.9	42
Other backward class	2.3	16.6	1,344	(98.1)	(48.3)	32	77.8	223
Other	1.1	15.3	366	*	*	4	81.6	56
Total	2.0	16.7	2,347	(87.9)	(36.9)	48	75.3	392

Note: Total includes children belonging to "other" religions and children whose caste/tribe is not known, who are not shown separately.

¹ Symptoms of ARI (cough accompanied by short, rapid breathing or difficulty breathing which was chest related)

² Excludes pharmacy, shop, and traditional healer

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

Table 55 Prevalence and treatment of diarrhoea

Percentage of children under age five who had diarrhoea in the two weeks preceding the survey and among children under age five who had diarrhoea in the two weeks preceding the survey, percentage who received advice or treatment from a health facility or health provider, who received oral rehydration therapy (ORT), who were given other treatments and who were given no treatment, by background characteristics, Telangana, 2015-16

Background characteristic	Percentage of children with diarrhoea taken to a health facility or health provider ¹	Oral rehydration therapy (ORT)				Other treatments					Number of children with diarrhoea	Percent- age not receiving any treatment			
		Percent- age given fluid from ORS packets	Percent- age given gruel	Percent- age given either ORS or gruel	Percent- age given increased fluids	Any ORT	Percent- age given antibiotic drug	Percent- age given other drug ²	Percent- age given zinc supplements	Percent- age given intravenous solution			Percent- age given home remedy/ herbal/ other		
Age in months															
<6	10.8	*	*	*	*	*	*	*	*	*	*	*	*	20	*
6-11	17.8	(74.2)	(12.9)	(53.5)	(9.2)	(58.8)	(17.6)	(20.0)	(22.4)	(0.0)	(10.1)	(19.3)	41	(19.3)	
12-23	10.0	64.8	51.1	19.4	57.1	6.1	58.9	14.8	35.9	1.7	6.3	19.7	49	19.7	
24-35	7.0	448	(81.1)	(48.0)	(21.4)	(64.9)	(5.1)	(1.9)	(27.9)	(0.0)	(0.0)	(23.5)	32	(23.5)	
36-47	4.9	510	*	*	*	*	*	*	*	*	*	*	25	*	
48-59	5.9	474	(75.0)	(73.4)	(30.1)	(77.6)	(2.4)	(9.1)	(32.6)	(0.0)	(2.4)	(7.6)	28	(7.6)	
Sex															
Male	8.7	1,252	71.5	57.6	19.1	62.6	6.1	65.3	10.4	11.0	34.4	0.0	5.6	108	15.9
Female	8.0	1,094	77.2	55.7	16.1	63.4	3.2	64.8	8.9	13.2	28.0	1.0	4.2	88	16.6
Residence															
Urban	8.1	1,106	79.5	61.8	24.2	72.4	2.1	73.8	10.4	11.4	30.2	0.0	4.3	90	13.5
Rural	8.6	1,241	69.4	52.5	12.3	55.0	7.1	57.7	9.2	12.5	32.7	0.8	5.6	106	18.4
Mother's schooling															
No schooling	4.3	450	*	*	*	*	*	*	*	*	*	*	*	19	*
<5 years complete	9.6	104	*	*	*	*	*	*	*	*	*	*	*	10	*
5-9 years complete	10.9	538	72.9	57.9	18.5	60.1	3.1	61.6	9.1	16.4	33.9	0.0	2.6	59	14.3
10-11 years complete	11.0	545	75.0	60.2	20.0	70.0	9.6	75.4	14.0	9.3	26.2	0.0	6.0	60	13.7
12 or more years complete	6.8	710	(83.7)	(55.0)	(11.7)	(57.8)	(2.5)	(57.8)	(4.5)	(13.1)	(31.4)	(1.7)	(9.7)	48	(18.3)

Continued...

Table 55 Prevalence and treatment of diarrhoea—Continued

Percentage of children with diarrhoea under age five who had diarrhoea in the two weeks preceding the survey and among children under age five who had diarrhoea in the two weeks preceding the survey, percentage who received advice or treatment from a health facility or health provider, who received oral rehydration therapy (ORT), who were given other treatments and who were given no treatment, by background characteristics, Telangana, 2015-16

Background characteristic	Percentage of children with diarrhoea	Number of children with diarrhoea	Oral rehydration therapy (ORT)					Other treatments					Number of children with diarrhoea		
			Percentage of children with diarrhoea taken to a health facility or health provider ¹	Percentage of children with diarrhoea taken to a health facility or health provider ¹ who received ORT	Percentage of children with diarrhoea taken to a health facility or health provider ¹ who received ORT and/or increased fluids	Percentage of children with diarrhoea taken to a health facility or health provider ¹ who received ORT and/or increased fluids and/or gruel	Percentage of children with diarrhoea taken to a health facility or health provider ¹ who received ORT and/or increased fluids and/or gruel and/or increased fluids	Percentage of children with diarrhoea given zinc supplements	Percentage of children with diarrhoea given other drug ²	Percentage of children with diarrhoea given antibiotic drug	Percentage of children with diarrhoea given intravenous solution	Percentage of children with diarrhoea given home remedy/ herbal/ other		Percentage of children with diarrhoea not receiving any treatment	
Religion															
Hindu	8.5	1,988	72.9	55.0	16.2	60.5	4.5	63.0	9.4	10.6	31.3	0.5	5.4	17.5	169
Muslim	7.3	288	(85.3)	(65.0)	(31.5)	(78.6)	(5.7)	(78.6)	(5.6)	(17.3)	(36.3)	(0.0)	(3.2)	(10.1)	21
Christian	8.9	67	*	*	*	*	*	*	*	*	*	*	*	*	6
Caste/tribe															
Scheduled caste	8.2	432	(75.6)	(56.0)	(19.4)	(63.1)	(10.2)	(66.1)	(13.7)	(22.0)	(41.0)	(2.4)	(0.0)	(8.9)	36
Scheduled tribe	7.8	191	*	*	*	*	*	*	*	*	*	*	*	*	15
Other backward class	8.7	1,344	69.7	52.9	16.2	57.0	4.0	59.7	9.3	10.8	31.1	0.0	7.3	20.6	117
Other	7.5	366	*	*	*	*	*	*	*	*	*	*	*	*	27
Total	8.4	2,347	74.0	56.8	17.8	63.0	4.8	65.1	9.7	12.0	31.6	0.4	5.0	16.2	196

Note: ORT includes a solution prepared from an oral rehydration salt (ORS) packet and/or gruel and/or increased fluids. Total includes children belonging to "other" religions and children whose caste/tribe is not known, who are not shown separately.

¹ Excludes pharmacy, shop, and traditional healer

² Includes antimotility drugs, other drugs, and unknown drugs

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

Table 56. Feeding practices during diarrhoea

Percent distribution of children under age five who had diarrhoea in the two weeks preceding the survey by amount of liquids and food given compared with normal practice, according to background characteristics, Telangana, 2015-16

Background characteristic	Amount of liquids given				Amount of food given				Total	Percentage given increased fluids and continued feeding ¹	Percentage given ORT and continued feeding ¹	Number of children with diarrhoea				
	More usual	Same as usual	Some-what less	Much less	None	Total	More usual	Same as usual					Some-what less	Much less	None	Never gave food
Age in months																
6-11	(9.2)	(27.7)	(23.0)	(27.3)	(12.8)	100.0	(0.0)	(19.7)	(37.3)	(17.8)	(7.3)	(17.9)	100.0	(4.6)	(35.9)	41
12-23	6.1	40.4	25.1	19.0	9.3	100.0	4.3	26.4	30.9	19.0	1.8	17.5	100.0	2.4	35.6	49
24-35	(2.2)	(43.0)	(36.5)	(16.1)	(2.2)	100.0	(1.6)	(17.8)	(51.9)	(17.0)	(10.3)	(1.4)	100.0	(0.0)	(43.9)	32
48-59	(3.8)	(31.9)	(38.4)	(20.7)	(5.2)	100.0	(0.0)	(24.5)	(39.2)	(32.9)	(3.4)	(0.0)	100.0	(3.8)	(51.7)	28
Sex																
Male	6.1	28.3	37.8	21.3	6.6	100.0	0.6	21.8	42.5	16.6	4.2	14.3	100.0	2.9	46.2	108
Female	3.2	41.1	30.9	14.7	10.2	100.0	4.2	21.2	40.2	20.7	6.0	7.6	100.0	2.2	39.8	88
Residence																
Urban	2.1	31.6	39.1	20.7	6.5	100.0	0.7	14.9	53.9	21.3	0.8	8.4	100.0	2.1	53.0	90
Rural	7.1	36.0	30.9	16.3	9.6	100.0	3.5	27.2	31.0	16.0	8.6	13.7	100.0	3.0	35.2	106
Mother's schooling																
5-9 years complete	3.1	43.4	34.7	13.8	5.1	100.0	2.6	32.8	34.5	19.1	3.1	7.8	100.0	1.5	44.6	59
10-11 years complete	9.6	30.2	30.5	23.1	6.5	100.0	0.8	22.2	42.3	18.0	4.2	12.5	100.0	4.9	53.7	60
12 or more years complete	(2.5)	(25.8)	(41.9)	(19.9)	(9.8)	100.0	(4.7)	(9.3)	(55.3)	(18.9)	(4.7)	(7.2)	100.0	(2.5)	(41.8)	48
Religion																
Hindu	4.5	34.5	36.6	17.1	7.3	100.0	2.2	21.3	43.0	17.4	5.4	10.7	100.0	2.3	42.6	169
Muslim	(5.7)	(36.4)	(29.2)	(14.7)	(13.9)	100.0	(2.9)	(26.5)	(41.1)	(10.1)	(0.0)	(19.4)	100.0	(5.7)	(57.7)	21
Caste/tribe																
Scheduled caste	(10.2)	(21.9)	(27.4)	(25.6)	(14.8)	100.0	(4.3)	(21.0)	(32.3)	(27.7)	(4.5)	(10.2)	100.0	(5.5)	(40.0)	36
Other backward class	4.0	37.8	30.1	20.3	7.8	100.0	0.8	24.3	36.3	19.9	4.1	14.7	100.0	1.6	35.5	117
Total	4.8	34.0	34.7	18.3	8.2	100.0	2.2	21.6	41.5	18.4	5.0	11.3	100.0	2.6	43.3	196

Note: It is recommended that children should be given more liquids to drink during diarrhoea and food should not be reduced. Total includes children age <6 or 36-47 months, children whose mothers have no schooling or have less than 5 years of schooling, Christian children, children belonging to scheduled tribes or "other" castes/tribes, and children whose caste/tribe is not known, who are not shown separately.

ORT = Oral rehydration therapy, which includes a solution prepared from an oral rehydration salt (ORS) packet and/or gruel and/or increased fluids

¹ Continued feeding includes children who were given more, same as usual, or somewhat less food during the diarrhoea episode

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

Table 57 Knowledge of ORS packets

Percentage of all women and percentage of women who had a live birth in the five years preceding the survey who know about ORS packets for treatment of diarrhoea, by background characteristics, Telangana, 2015-16

Background characteristic	All women		Women who gave birth in the past five years	
	Percentage who know about ORS packets	Number of women	Percentage who know about ORS packets	Number of women
Age				
15-19	78.1	997	78.8	76
20-24	82.0	1,355	85.6	655
25-34	84.5	2,470	90.0	989
35-49	79.1	2,745	87.8	80
Residence				
Urban	82.8	3,619	88.6	842
Rural	79.9	3,948	87.1	958
Schooling				
No schooling	74.3	2,458	84.8	356
<5 years complete	81.2	337	83.4	82
5-9 years complete	82.3	1,493	85.8	391
10-11 years complete	86.5	1,435	87.6	419
12 or more years complete	85.8	1,844	92.0	552
Religion				
Hindu	80.8	6,529	86.9	1,554
Muslim	84.2	774	93.2	193
Christian	84.8	248	95.8	50
Caste/tribe				
Scheduled caste	82.1	1,395	90.0	335
Scheduled tribe	79.2	597	85.0	148
Other backward class	81.1	4,393	86.8	1,036
Other	82.8	1,137	91.0	271
Don't know	60.2	44	*	10
Total	81.3	7,567	87.8	1,800

Note: Total includes women belonging to "other" religions, who are not shown separately.

ORS = Oral rehydration salt

* Percentage not shown; based on fewer than 25 unweighted cases

Table 58 ICDS coverage and utilization of ICDS services

Percentage of children under age six years who received any service and received specific services from an anganwadi centre (AWC) in the 12 months preceding the survey, by background characteristics, Telangana, 2015-16

Background characteristic	Percentage of children age 0-71 months who received from an AWC			Children age 36-71 months		Children age 0-59 months		Children age 0-59 months who were weighed at an AWC			
	Any benefits ¹	Supplementary food ²	Any immunizations	Health check-ups	Number of children	Percentage who went for early childhood care/preschool to an AWC	Number of children	Percentage of children who were weighed at an AWC	Number of children	Percentage whose mothers received counselling from an AWC after child was weighed	Number of children weighed
Age in months											
<12	65.0	61.9	49.6	50.5	419	na	na	50.4	419	57.6	212
12-23	68.4	67.6	55.1	51.9	495	na	na	58.5	495	61.8	289
24-35	64.8	63.9	47.9	49.9	448	na	na	53.9	448	69.6	242
36-47	59.2	57.9	43.1	45.3	510	47.7	510	51.1	510	72.5	261
48-59	48.9	47.6	32.9	35.1	474	41.2	474	38.9	474	57.5	184
60-71	38.6	36.6	32.0	30.5	519	30.6	519	na	na	na	na
0-35	66.1	64.6	51.1	50.8	1,363	na	na	54.5	1,363	63.2	742
36-71	48.8	47.3	36.0	37.0	1,503	39.7	1,503	45.2	984	66.3	445
Sex											
Male	55.7	53.9	41.0	42.2	1,495	37.2	775	49.6	1,252	64.8	622
Female	58.5	57.4	45.6	45.1	1,371	42.5	728	51.7	1,094	63.9	566
Residence											
Urban	38.5	37.2	28.3	29.2	1,372	23.5	745	33.2	1,106	54.3	367
Rural	74.1	72.4	56.9	56.7	1,494	55.7	759	66.1	1,241	68.8	821
Mother's schooling											
No schooling	67.8	66.3	51.6	51.8	579	51.7	338	57.9	450	61.0	260
<5 years complete	67.9	66.2	44.9	52.6	132	55.3	84	60.0	104	62.3	62
5-9 years complete	62.4	60.5	48.2	45.8	669	43.4	377	55.9	538	68.0	300
10-11 years complete	57.6	55.9	45.9	45.4	655	40.1	331	51.9	545	70.0	283
12 or more years complete	43.2	42.1	30.9	33.2	831	21.3	373	39.6	710	58.3	282

Continued...

Table 58 ICDS coverage and utilization of ICDS services—Continued

Background characteristic	Percentage of children age 0-71 months who received from an AWC			Children age 36-71 months		Children age 0-59 months		Children age 0-59 months who were weighed at an AWC		
	Any benefits ¹	Supplementary food ²	Any immunizations	Health check-ups	Number of children	Percentage who went for early childhood care/preschool to an AWC	Number of children	Percentage of children who were weighed at an AWC	Number of children	Percentage whose mothers received counselling from an AWC after child was weighed
Religion										
Hindu	58.3	56.7	43.9	44.2	2,422	41.1	1,254	52.4	1,988	64.5
Muslim	46.8	45.5	35.0	37.0	355	33.7	200	36.5	288	64.0
Christian	66.2	65.4	56.3	53.7	84	(30.9)	47	57.2	67	(59.6)
Caste/tribe										
Scheduled caste	66.0	63.9	50.4	52.7	520	51.1	273	60.9	432	69.3
Scheduled tribe	68.8	66.6	50.0	52.8	228	49.6	115	56.6	191	61.7
Other backward class	57.7	56.5	43.6	43.6	1,679	40.1	888	51.7	1,344	63.8
Other	38.0	36.3	29.6	28.3	420	20.3	217	31.7	366	59.7
Don't know	(41.8)	(37.5)	(29.9)	(19.7)	19	*	11	*	13	*
Total	57.1	55.6	43.2	43.6	2,866	39.7	1,503	50.6	2,347	64.3

Note: Total includes children belonging to "other" religions, who are not shown separately.

na = Not applicable

ICDS = Integrated Child Development Services

¹ AWC benefits for children include distribution of supplementary food, growth monitoring, immunizations, health check-ups, and early childhood care/ preschool

² Supplementary food includes both food cooked and served at the AWC on a daily basis or given in the form of take home rations

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

Table 59 Utilization of ICDS services during pregnancy and while breastfeeding

Percentage of children under age six years whose mothers received specific benefits from an anganwadi centre (AWC) during pregnancy and while breastfeeding, according to background characteristics, Telangana, 2015-16

Background characteristic	Mother received from an AWC during pregnancy					Mother received from an AWC while breastfeeding ²				
	Any benefits	Supplementary food ¹	Health check-ups	Health and nutrition education	Number of children	Any benefits	Supplementary food ¹	Health check-ups	Health and nutrition education	Number of children breastfed
Residence										
Urban	43.3	43.1	31.3	34.9	1,372	39.9	39.6	27.7	32.0	1,368
Rural	78.8	78.5	62.1	70.8	1,494	75.5	74.7	58.5	66.2	1,494
Mother's schooling										
No schooling	70.6	70.1	55.5	63.5	579	65.8	65.3	49.6	56.7	579
<5 years complete	66.2	66.2	54.3	58.5	132	62.9	62.9	49.6	54.6	132
5-9 years complete	69.0	68.6	50.4	56.3	669	65.5	64.9	45.2	53.3	669
10-11 years complete	65.7	65.5	50.8	58.3	655	62.1	61.3	48.4	54.1	654
12 or more years complete	46.1	46.0	35.5	40.2	831	44.3	43.8	34.0	38.3	829
Religion										
Hindu	63.2	62.9	49.2	55.9	2,422	60.1	59.5	45.5	51.8	2,420
Muslim	51.7	51.2	34.1	37.3	355	47.0	46.3	31.0	35.8	354
Christian	64.8	64.8	48.6	56.6	84	62.1	62.1	47.5	52.9	84
Caste/tribe										
Scheduled caste	73.0	72.8	54.1	60.1	520	68.6	68.1	48.5	56.0	520
Scheduled tribe	66.3	65.9	51.5	58.9	228	64.4	63.4	51.0	55.7	228
Other backward class	62.9	62.6	49.8	56.1	1,679	59.6	59.2	45.7	52.3	1,677
Other	41.1	40.9	27.5	33.5	420	39.4	38.8	27.4	30.7	418
Don't know	(66.3)	(61.4)	(36.2)	(38.8)	19	(35.2)	(30.3)	(13.0)	(16.9)	19
Total	61.8	61.5	47.4	53.6	2,866	58.5	58.0	43.7	49.9	2,863

Note: Total includes children belonging to "other" religions, who are not shown separately.

ICDS = Integrated Child Development Services

¹ Supplementary food includes both food cooked and served at the AWC on a daily basis and food given in the form of take home rations

² Services are usually provided to breastfeeding mothers during the first six months of breastfeeding

() Based on 25-49 unweighted cases

Table 60 Nutritional status of children

Percentage of children under age five years classified as malnourished according to three anthropometric indices of nutritional status: height-for-age, weight-for-height, and weight-for-age, by background characteristics, Telangana, 2015-16

Background characteristic	Height-for-age ¹						Weight-for-height						Weight-for-age						
	Percentage below		Mean Z-score (SD)	Percentage below		Mean Z-score (SD)	Percentage below		Mean Z-score (SD)	Percentage below		Mean Z-score (SD)	Percentage above		Mean Z-score (SD)	Percentage above		Mean Z-score (SD)	
	-3 SD	-2 SD ²		-3 SD	-2 SD ²		-3 SD	-2 SD ²		-3 SD	-2 SD ²		+2 SD	+2 SD ²		+2 SD	+2 SD ²		
Age in months																			
<6	2.9	12.3	-0.1	17.4	39.8	1.5	-1.4	6.6	19.5	6.6	-1.4	6.6	19.5	2.2	-1.1	155			
6-8	1.7	12.5	-0.2	8.6	22.8	0.9	-1.1	5.8	16.0	5.8	-1.1	5.8	16.0	0.0	-1.0	105			
9-11	3.8	11.6	-0.1	11.3	29.5	0.0	-1.3	7.4	21.9	7.4	-1.3	7.4	21.9	0.0	-1.1	107			
12-17	4.7	18.7	-0.7	7.1	20.4	0.0	-1.1	6.7	26.3	6.7	-1.1	6.7	26.3	1.8	-1.2	192			
18-23	11.5	32.9	-1.4	6.2	25.3	0.5	-1.2	11.0	33.2	11.0	-1.2	11.0	33.2	0.0	-1.5	249			
24-35	12.3	34.8	-1.6	1.8	9.8	0.0	-0.8	7.2	27.4	7.2	-0.8	7.2	27.4	0.0	-1.4	397			
36-47	11.0	35.4	-1.6	0.7	10.8	0.2	-0.8	6.9	32.0	6.9	-0.8	6.9	32.0	0.7	-1.5	446			
48-59	10.6	29.3	-1.4	2.7	15.5	2.5	-1.0	9.7	32.2	9.7	-1.0	9.7	32.2	2.0	-1.5	409			
Sex																			
Male	9.0	27.9	-1.2	5.1	17.9	0.1	-1.1	7.6	29.2	7.6	-1.1	7.6	29.2	0.6	-1.4	1,107			
Female	9.4	28.3	-1.2	4.3	18.0	1.4	-1.0	8.3	27.6	8.3	-1.0	8.3	27.6	1.2	-1.3	954			
Birth interval in months³																			
First birth ⁴	6.6	22.5	-1.1	3.7	16.1	0.3	-0.9	6.5	26.0	6.5	-0.9	6.5	26.0	0.4	-1.3	874			
<24	13.2	37.2	-1.4	4.4	19.3	0.6	-1.0	10.8	31.4	10.8	-1.0	10.8	31.4	0.8	-1.5	369			
24-47	7.5	28.4	-1.1	5.3	18.1	0.2	-1.1	6.7	28.3	6.7	-1.1	6.7	28.3	0.1	-1.4	528			
48+	14.1	32.5	-1.2	6.6	22.1	3.6	-1.1	10.1	31.8	10.1	-1.1	10.1	31.8	5.0	-1.4	228			
Birth order³																			
1	6.7	22.6	-1.1	3.7	16.2	0.3	-0.9	6.5	26.0	6.5	-0.9	6.5	26.0	0.4	-1.3	867			
2-3	10.7	32.3	-1.2	5.3	19.0	1.1	-1.0	8.6	30.0	8.6	-1.0	8.6	30.0	1.2	-1.4	1,066			
4-5	10.8	28.9	-1.2	5.2	24.1	0.0	-1.1	10.9	29.5	10.9	-1.1	10.9	29.5	3.4	-1.5	64			
Size at birth³																			
Very small	(23.4)	(28.1)	(-1.7)	(2.5)	(21.1)	(0.0)	(-1.1)	(21.7)	(37.2)	(21.7)	(-1.1)	(21.7)	(37.2)	(0.0)	(-1.8)	35			
Small	17.7	42.3	-1.7	7.1	28.0	0.0	-1.4	19.7	48.2	19.7	-1.4	19.7	48.2	0.0	-1.9	105			
Average or larger	8.2	27.2	-1.1	4.4	17.3	0.8	-1.0	6.8	26.9	6.8	-1.0	6.8	26.9	1.0	-1.3	1,843			

Continued...

Table 60 Nutritional status of children—Continued

Percentage of children under age five years classified as malnourished according to three anthropometric indices of nutritional status: height-for-age, weight-for-age, and weight-for-age, by background characteristics, Telangana, 2015-16

Background characteristic	Height-for-age ¹					Weight-for-height					Weight-for-age					Number of children
	Percentage below -3 SD	Percentage below -2 SD ²	Mean Z-score (SD)	Percentage below -3 SD	Percentage below -2 SD ²	Percentage below -3 SD	Percentage below -2 SD ²	Mean Z-score (SD)	Percentage below -3 SD	Percentage below -2 SD ²	Percentage above +2 SD	Percentage above +2 SD	Mean Z-score (SD)	Percentage below -2 SD ²	Percentage above +2 SD	
Residence																
Urban	6.2	20.9	-0.9	3.7	14.6	1.2	14.6	-0.9	5.3	22.2	1.8	22.2	-1.1	870		
Rural	11.4	33.3	-1.4	5.6	20.4	0.4	20.4	-1.1	9.8	33.1	0.2	33.1	-1.5	1,191		
Mother's schooling⁵																
No schooling	15.7	40.2	-1.6	4.9	22.0	0.4	22.0	-1.2	13.5	41.1	0.0	41.1	-1.7	406		
<5 years complete	16.3	45.4	-1.6	5.2	26.7	0.7	26.7	-1.3	13.6	43.4	0.0	43.4	-1.8	98		
5-9 years complete	9.9	31.3	-1.4	4.3	19.8	0.7	19.8	-1.0	8.3	29.8	0.1	29.8	-1.5	473		
10-11 years complete	6.6	27.2	-1.1	5.8	17.4	0.3	17.4	-1.0	8.2	29.4	1.4	29.4	-1.4	473		
12 or more years complete	4.5	14.5	-0.6	4.3	13.0	1.4	13.0	-0.8	2.8	14.9	1.9	14.9	-0.9	575		
Religion																
Hindu	9.2	28.5	-1.2	4.7	18.0	0.9	18.0	-1.0	8.2	28.4	0.9	28.4	-1.4	1,731		
Muslim	9.1	23.7	-1.1	4.0	15.1	0.0	15.1	-1.0	5.0	24.2	0.8	24.2	-1.3	261		
Christian	8.6	34.2	-1.3	9.2	27.7	0.0	27.7	-1.3	10.9	48.4	0.0	48.4	-1.7	68		
Caste/tribe																
Scheduled caste	12.0	33.2	-1.4	6.2	20.8	0.3	20.8	-1.2	11.2	37.2	0.2	37.2	-1.6	387		
Scheduled tribe	10.2	34.7	-1.4	4.7	20.2	0.0	20.2	-1.1	10.6	35.4	0.0	35.4	-1.6	172		
Other backward class	9.1	27.7	-1.2	4.6	17.9	0.5	17.9	-1.0	7.2	27.1	0.6	27.1	-1.3	1,221		
Other	5.2	18.5	-0.8	3.8	12.6	3.1	12.6	-0.8	5.0	18.3	3.9	18.3	-1.0	268		
Mother's interview status																
Interviewed	8.9	27.9	-1.2	4.6	17.9	0.7	17.9	-1.0	7.8	28.3	0.9	28.3	-1.4	1,999		
Not interviewed but in the household	(19.4)	(32.7)	(-1.7)	(20.6)	(28.8)	(4.2)	(28.8)	(-1.3)	(26.3)	(40.2)	(0.0)	(40.2)	(-1.8)	27		
Not interviewed, and not in the household ⁶	(15.0)	(31.7)	(-1.2)	(1.9)	(11.9)	(0.0)	(11.9)	(-1.0)	(2.8)	(31.8)	(0.0)	(31.8)	(-1.4)	35		

Continued...

Table 60 Nutritional status of children—Continued

Percentage of children under age five years classified as malnourished according to three anthropometric indices of nutritional status: height-for-age, weight-for-height, and weight-for-age, by background characteristics, Telangana, 2015-16, and total for NFHS-3

Background characteristic	Height-for-age ¹			Weight-for-height			Weight-for-age			Number of children		
	Percentage below -3 SD	Percentage below -2 SD ²	Mean Z-score (SD)	Percentage below -3 SD	Percentage below -2 SD ²	Percentage above +2 SD	Mean Z-score (SD)	Percentage below -3 SD	Percentage below -2 SD ²		Percentage above +2 SD	
Mother's nutritional status⁷												
Underweight (BMI < 18.5)	12.4	35.4	-1.4	6.4	25.9	0.5	-1.2	12.5	38.7	0.1	-1.7	534
Normal (BMI 18.5-24.9)	8.5	27.8	-1.2	4.6	16.3	0.2	-1.0	7.6	27.5	0.1	-1.4	1,096
Overweight (BMI ≥ 25.0)	5.6	17.3	-0.7	3.1	11.9	2.5	-0.7	2.7	16.0	3.6	-0.9	370
Missing	(15.7)	(35.4)	(-1.1)	(5.9)	(17.8)	(4.5)	(-0.8)	(9.3)	(31.7)	(10.2)	(-1.1)	26
Child's living arrangements												
Living with both parents	9.2	28.4	-1.2	4.9	17.8	0.8	-1.0	8.0	28.3	1.0	-1.4	1,840
Living with one or neither parent	8.9	25.3	-1.1	3.6	19.5	0.4	-1.1	7.1	29.7	0.3	-1.4	221
Total	9.2	28.1	-1.2	4.8	18.0	0.7	-1.0	7.9	28.5	0.9	-1.4	2,061

Note: Table is based on children who stayed in the household the night before the interview. Each of the indices is expressed in standard deviation units (SD) from the median of the WHO Child Growth Standards adopted in 2006. The indices in this table are NOT comparable to those based on the previously used 1977 NCHS/CDC/WHO Reference. Table is based on children with valid dates of birth (month and year) and valid measurement of both height and weight. Total includes children of birth order 6 or more, children whose size at birth is not known, children belonging to "other" religions, and children whose caste/tribe is not known, who are not shown separately.

¹ Recumbent length is measured for children under age 2, or in the few cases when the age of the child is unknown and the child is less than 85 cm; standing height is measured for all other children

² Includes children who are below -3 standard deviations (SD) from the WHO Child Growth Standards population median

³ Excludes children whose mothers were not interviewed

⁴ First born twins (triplets, etc.) are counted as first births because they do not have a previous birth interval

⁵ For women who are not interviewed, information is taken from the Household Questionnaire. Excludes children whose mothers are not listed in the household schedule.

⁶ Includes children whose mothers are deceased

⁷ Excludes children whose mothers were not weighed and measured, children whose mothers were not interviewed, and children whose mothers are pregnant or gave birth within the preceding 2 months. Mother's nutritional status in terms of BMI (body mass index) is presented in Table 70.

() Based on 25-49 unweighted cases

Table 61 Initial breastfeeding

Among last-born children who were born in the two years preceding the survey, the percentage who were ever breastfed, and the percentage who started breastfeeding within one hour and one day of birth, and among last-born children born in the two years preceding the survey who were ever breastfed, the percentage who received a prelacteal feed, by background characteristics, Telangana, 2015-16

Background characteristic	Among last-born children born in the past two years:				Among last-born children born in the past two years who were ever breastfed:	
	Percentage ever breastfed	Percentage who started breastfeeding within one hour of birth ¹	Percentage who started breastfeeding within one day of birth ²	Number of last-born children	Percentage who received a prelacteal feed ³	Number of last-born ever breastfed children
Residence						
Urban	97.0	37.7	75.7	401	32.6	389
Rural	97.9	34.5	73.2	494	21.0	484
Sex						
Male	96.2	38.5	72.7	471	27.0	454
Female	98.9	33.0	76.1	424	25.2	419
Mother's schooling						
No schooling	97.3	28.4	72.6	150	17.0	146
<5 years complete	(97.5)	(41.4)	(74.3)	31	(8.1)	30
5-9 years complete	96.8	36.6	71.0	187	21.8	181
10-11 years complete	97.4	35.4	72.5	221	24.1	216
12 or more years complete	98.1	39.0	78.5	305	36.6	299
Religion						
Hindu	97.5	35.4	74.4	782	26.9	763
Muslim	97.4	34.4	75.4	85	23.1	82
Caste/tribe						
Scheduled caste	96.3	43.2	78.5	177	20.0	171
Scheduled tribe	96.9	25.6	69.3	75	27.1	73
Other backward class	97.6	35.9	72.5	509	23.3	497
Other	99.0	33.7	79.1	125	45.8	124
Assistance at delivery						
Health personnel ⁴	97.7	35.7	73.8	820	26.6	802
Dai (TBA)	(96.3)	(15.7)	(85.3)	25	(5.1)	24
Other/no one	94.6	49.3	76.8	49	29.9	47
Place of delivery						
Health facility	97.6	36.1	73.5	840	27.0	820
At home	96.5	30.8	86.6	53	13.6	51
Total	97.5	35.9	74.3	895	26.2	873

Note: Table is based on last-born children born in the past two years whether the children are living or dead at the time of interview. Total includes children belonging to Christian or "other" religions, children whose caste/tribe is not known, and children who were delivered in an "other" place of delivery, who are not shown separately.

TBA = Traditional birth attendant

¹ Includes children who started breastfeeding immediately after birth

² Includes children who started breastfeeding within one hour of birth

³ Children given something other than breastmilk during the first three days of life

⁴ Doctor, nurse, midwife, auxiliary nurse midwife, lady health visitor, or other health personnel

() Based on 25-49 unweighted cases

Table 62 Breastfeeding status by age

Percent distribution of youngest children under two years living with the mother by breastfeeding status, percentage currently breastfeeding, and percentage of all children under two years using a bottle with a nipple, according to age in months, Telangana, 2015-16

Age in months	Breastfeeding and consuming:						Total	Percentage currently breastfeeding	Number of youngest children under two years living with the mother	Percentage using a bottle with a nipple	Number of all children under two years
	Not breastfeeding	Exclusively breastfed	Plain water only	Non-milk liquids/juice	Other milk	Complementary foods					
<2	(2.8)	(86.7)	(4.4)	(2.0)	(2.8)	(1.4)	100.0	(97.2)	33	(2.6)	36
2-3	4.3	73.0	6.4	2.0	4.8	9.5	100.0	95.7	78	10.5	79
4-5	1.8	52.2	17.1	1.0	8.4	19.4	100.0	98.2	73	10.5	73
6-8	0.6	18.1	15.5	2.6	7.3	55.9	100.0	99.4	119	22.3	120
9-11	7.1	6.7	7.3	0.0	3.1	75.9	100.0	92.9	112	21.8	112
12-17	10.6	4.1	5.2	3.1	1.8	75.1	100.0	89.4	210	23.8	218
18-23	26.3	0.8	4.0	2.2	1.6	65.1	100.0	73.7	249	30.8	277
<4	3.9	77.1	5.8	2.0	4.2	7.0	100.0	96.1	111	8.1	115
<6	3.1	67.3	10.3	1.6	5.8	11.9	100.0	96.9	184	9.0	188
6-9	1.2	17.2	13.6	2.0	6.8	59.1	100.0	98.8	153	21.0	153
12-15	11.5	5.1	5.9	1.9	1.8	73.9	100.0	88.5	139	26.3	142
12-23	19.1	2.3	4.6	2.6	1.7	69.7	100.0	80.9	459	27.7	495
20-23	28.8	0.9	4.5	2.1	1.8	61.9	100.0	71.2	165	31.4	188

Note: Breastfeeding status refers to a "24-hour" period (yesterday and last night). Children who are classified as breastfeeding and consuming plain water only consumed no liquid or solid supplements. The categories of not breastfeeding, exclusively breastfed, breastfeeding and consuming plain water, non-milk liquids/juice, other milk, and complementary foods (solids and semi-solids) are hierarchical and mutually exclusive, and their percentages add to 100 percent. Thus children who receive breastmilk and non-milk liquids and who do not receive other milk and who do not receive complementary foods are classified in the non-milk liquid category even though they may also get plain water. Any children who get complementary food are classified in that category as long as they are breastfeeding as well.

() Based on 25-49 unweighted cases

Table 6.3 Median duration of breastfeeding and infant and young child feeding (IYCF) practices

Background characteristic	Median duration (months) of breastfeeding among last-born children born in the last three years ¹										Among breastfed children 6-23 months, percentage fed:					Among non-breastfed children 6-23 months, percentage fed:					Among all children 6-23 months, percentage fed:											
	Any breast-feeding		Exclu- sive breast-feeding		Predom- inant breast-feeding ²		Number of children		4+ food groups ³		Minimum meal fre- quency ⁴		Both 4+ food groups and minimum meal fre- quency ⁴		Milk or milk products ⁵		4+ food groups ³		Minimum meal fre- quency ⁴		Breast- milk, or milk products ⁷		4+ food groups ³		Minimum meal fre- quency ⁸		With 3 IYCF prac- tices ⁶		Number of children			
	na	na	na	na	na	na	na	na	9.7	33.0	5.0	119	*	*	*	*	*	*	99.7	9.5	33.6	4.9	120									
Age in months																																
6-8	na	na	na	na	na	na	na	9.7	33.0	5.0	119	*	*	*	*	*	*	99.7	9.5	33.6	4.9	120										
9-11	na	na	na	na	na	na	na	12.1	14.1	4.7	104	*	*	*	*	*	*	97.1	13.6	16.6	4.3	112										
12-17	na	na	na	na	na	na	na	30.7	21.4	5.4	188	*	*	*	*	*	*	94.7	29.9	23.9	5.4	210										
18-23	na	na	na	na	na	na	na	35.3	33.6	19.8	183	53.1	45.4	55.8	15.3	64	87.8	37.9	39.4	18.6	248											
Sex																																
Male	27.4	4.7	6.0	6.0	6.0	645	324	22.6	28.5	10.0	324	59.3	48.9	62.0	17.5	51	94.5	26.2	33.1	11.0	375											
Female	25.0	3.7	5.6	5.6	5.6	580	270	27.1	23.4	9.2	270	(46.2)	(26.6)	(42.8)	(4.4)	45	92.3	27.0	26.2	8.6	315											
Residence																																
Urban	24.6	4.3	5.9	5.9	5.9	560	269	32.3	28.8	14.8	269	(55.5)	(37.3)	(55.3)	(9.7)	56	92.4	33.1	33.4	13.9	324											
Rural	33.4	4.2	5.8	5.8	5.8	665	325	18.4	24.1	5.4	325	49.9	40.1	50.0	13.7	41	94.5	20.8	26.9	6.3	366											
Mother's schooling																																
No schooling	≥36.0	(3.2)	(5.1)	(5.1)	(5.1)	226	105	15.7	21.6	9.2	105	*	*	*	*	9	94.8	17.1	23.0	9.1	114											
<5 years complete	a	a	a	a	a	48	25	(21.9)	(15.6)	(3.8)	25	*	*	*	*	2	(95.1)	(22.5)	(17.3)	(3.5)	27											
5-9 years complete	(27.7)	(3.8)	5.7	5.7	5.7	257	126	24.1	28.6	9.4	126	*	*	*	*	22	93.2	25.0	30.8	8.9	148											
10-11 years complete	(24.1)	(4.4)	6.0	6.0	6.0	284	144	20.9	28.5	7.8	144	(43.5)	(33.3)	(50.8)	(4.7)	32	89.8	23.1	32.5	7.2	176											
12 or more years complete	23.2	5.0	6.1	6.1	6.1	411	193	33.1	26.8	12.2	193	(69.8)	(50.4)	(66.1)	(23.1)	32	95.7	35.6	32.4	13.7	225											
Religion																																
Hindu	25.7	4.4	5.9	5.9	5.9	1,062	519	25.0	26.7	9.2	519	51.4	42.0	52.4	12.1	85	93.2	27.4	30.3	9.6	603											
Muslim	*	*	(5.7)	(5.7)	(5.7)	127	56	15.4	18.3	7.3	56	*	*	*	*	7	96.3	16.0	22.5	7.5	64											

Continued...

Table 63 Median duration of breastfeeding and infant and young child feeding (IYCF) practices—Continued

Background characteristic	Median duration (months) of breastfeeding among last-born children born in the last three years ¹		Among breastfed children 6-23 months, percentage fed:				Among nonbreastfed children 6-23 months, percentage fed:				Among all children 6-23 months, percentage fed:						
	Any breast-feeding	Exclu- sive breast-feeding	Predom- inant breast-feeding ²	Number of children	4+ food groups ³	Minimum meal fre- quency ⁴	Both 4+ food groups and minimum meal fre- quency	Milk or milk products ⁵	4+ food groups ³	Minimum meal fre- quency ⁴	With 3 prac- tices ⁶	Number of children	Breast milk, or milk products ⁷	4+ food groups ³	Minimum meal fre- quency ⁸	With 3 prac- tices ⁶	Number of children
Caste/tribe																	
Scheduled caste	(29.7)	(5.0)	(5.9)	227	20.9	32.0	9.8	*	*	*	*	8	97.5	21.3	33.1	9.7	138
Scheduled tribe	a	a	a	107	24.3	25.0	13.8	*	*	*	*	3	97.4	24.4	27.4	13.2	55
Other backward class	26.6	4.4	6.3	708	23.8	24.9	8.1	32.5	47.7	46.0	8.9	67	91.0	26.0	28.6	8.2	392
Other	*	(3.1)	(5.1)	175	35.2	24.0	13.8	81	*	*	*	18	95.0	38.6	33.6	15.6	99
Total	25.7	4.2	5.8	1,225	24.7	26.2	9.6	594	53.2	53.0	11.4	96	93.5	26.6	30.0	9.9	690

Note: Total includes children belonging to Christian or "other" religions, and children whose caste/tribe is not known, who are not shown separately.
a = Omitted because the smoothed percentage is below 50% in the first age cell (<2 months) and therefore the median cannot be calculated
na = Not applicable
¹ Median and mean durations are based on the distributions at the time of the survey of the proportion of births by months since birth. Includes children living and deceased at the time of the survey. It is assumed that children not currently living with the mother are not currently breastfeeding.
² Either exclusively breastfed or received breastmilk and plain water and/or non-milk liquids only
³ Food groups are: a. infant formula, milk other than breastmilk, cheese or yogurt or other milk products; b. foods made from grains or roots, including porridge or gruel, fortified baby food; c. vitamin A-rich fruits and vegetables; d. other fruits and vegetables; e. eggs; f. meat, poultry, fish, shellfish, or organ meats; g. beans, peas, lentils, or nuts; h. foods made with oil, fat, ghee, or butter
⁴ Receiving solid or semi-solid food at least twice a day for breastfed infants 6-8 months and at least three times a day for breastfed children 9-23 months
⁵ Includes two or more feedings of commercial infant formula, fresh, tinned and powdered animal milk, and yogurt
⁶ Nonbreastfed children age 6-23 months are considered to be fed with a minimum standard of three Infant and Young Child Feeding Practices if they receive other milk or milk products at least twice a day, receive the minimum meal frequency, and receive solid or semi-solid foods from at least four food groups not including the milk or milk products food group
⁷ Breastfeeding, or not breastfeeding and receiving two or more feedings of commercial infant formula, fresh, tinned, and powdered animal milk, and yogurt
⁸ Children are fed the minimum recommended number of times per day according to their age and breastfeeding status as described in footnotes 4 and 6.
() Based on 25-49 unweighted cases
* Percentage not shown; based on fewer than 25 unweighted cases

Table 64 Child feeding practices and nutritional status of children by district

Among last-born children in the past 2 years, percentage breastfed within one hour of birth, percentage of youngest children under age 6 months living with the mother who are exclusively breastfed, and percentage of children under age five years classified as malnourished according to three anthropometric indices of nutritional status: height-for-age, weight-for-height, and weight-for-age, by district, Telangana, 2015-16

District	Among last born children in the past 2 years		Percentage of children under age 6 months exclusively breastfed ¹		Height-for-age ²	Weight-for-height	Weight-for-age	Number of children
	Percentage breastfed within one hour of birth	Number of children	Percentage	Number of children	Percentage below -2 SD ³	Percentage below -2 SD ³	Percentage below -2 SD ³	
Adilabad	23.8	66	(54.0)	20	38.2	22.1	35.8	147
Hyderabad	42.0	129	*	32	15.7	14.1	16.8	318
Karimnagar	37.1	79	*	17	24.3	19.3	25.4	202
Khammam	43.8	64	*	13	26.5	13.7	22.2	159
Mahbubnagar	33.7	102	(55.1)	22	37.0	18.6	34.5	239
Medak	51.2	66	*	9	33.4	20.7	37.0	147
Nalgonda	35.7	88	*	18	28.9	23.1	34.1	220
Nizamabad	29.2	70	*	12	36.6	22.0	36.3	141
Rangareddy	29.2	161	*	25	26.2	14.8	25.8	319
Warangal	38.9	70	*	15	26.6	16.6	29.1	170
Telangana	35.9	895	67.3	184	28.1	18.0	28.5	2,061

¹ Based on the youngest child living with the mother

² Recumbent length is measured for children under age 2 years, or in the few cases when the age of the child is unknown and the child is less than 85 cm; standing height is measured for all other children.

³ Based on the WHO Child Growth Standards population median

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

Table 65 Prevalence of anaemia in children

Percentage of children age 6-59 months classified as having anaemia, by background characteristics, Telangana, 2015-16

Background characteristic	Anaemia status by haemoglobin level			Any anaemia (<11.0 g/dl)	Number of children
	Mild (10.0-10.9 g/dl)	Moderate (7.0-9.9 g/dl)	Severe (<7.0 g/dl)		
Age in months					
6-11	25.1	42.3	3.9	71.3	183
12-23	24.4	47.9	3.4	75.7	410
24-35	28.8	37.5	2.4	68.6	387
36-47	22.3	25.3	2.4	50.0	423
48-59	23.7	19.2	1.2	44.0	392
Sex					
Male	25.2	35.1	1.9	62.2	985
Female	24.2	31.5	3.2	58.9	810
Birth order¹					
1	25.4	30.9	2.3	58.6	752
2-3	24.3	35.6	2.7	62.6	930
4-5	19.9	44.3	3.3	67.4	58
Residence					
Urban	24.6	25.7	1.3	51.6	765
Rural	24.8	39.2	3.4	67.5	1,030
Mother's schooling²					
No schooling	26.0	36.2	3.4	65.6	367
<5 years complete	23.3	45.2	1.7	70.2	90
5-9 years complete	25.2	36.2	4.0	65.4	425
10-11 years complete	23.6	33.0	1.9	58.5	421
12 or more years complete	24.9	27.9	1.2	54.1	460
Religion					
Hindu	24.2	34.2	2.9	61.3	1,510
Muslim	27.5	28.4	0.5	56.3	228
Christian	27.5	35.4	1.4	64.3	57
Caste/tribe					
Scheduled caste	30.4	39.2	2.6	72.3	332
Scheduled tribe	20.6	42.3	3.7	66.5	156
Other backward class	24.3	31.6	2.3	58.2	1,071
Other	21.2	28.2	2.5	51.8	228

Continued...

Table 65 Prevalence of anaemia in children—Continued

Percentage of children age 6-59 months classified as having anaemia, by background characteristics, Telangana, 2015-16

Background characteristic	Anaemia status by haemoglobin level			Any anaemia (<11.0 g/dl)	Number of children
	Mild (10.0-10.9 g/dl)	Moderate (7.0-9.9 g/dl)	Severe (<7.0 g/dl)		
Mother's interview status					
Interviewed	24.6	33.9	2.5	61.1	1,743
Not interviewed, and not in the household ³	(21.2)	(18.9)	(2.1)	(42.2)	32
Child's living arrangements					
Living with both parents	25.0	33.6	2.5	61.1	1,616
Living with one or neither parent	22.8	32.1	2.8	57.6	179
Mother's anaemia status⁴					
Not anaemic	24.7	26.7	2.7	54.1	776
Mildly anaemic	25.1	38.0	1.8	65.0	645
Moderately/severely anaemic	25.3	44.4	3.1	72.8	315
Total	24.7	33.5	2.5	60.7	1,795

Note: Table is based on children who stayed in the household the night before the interview. Prevalence of anaemia, based on haemoglobin levels, is adjusted for altitude using the CDC formulas (Centers for Disease Control (CDC). 1998. Recommendations to prevent and control iron deficiency in the United States. *Morbidity and Mortality Weekly Report* 47 (RR-3): 1-29). Haemoglobin levels shown in grams per decilitre (g/dl). Total includes children of birth order 6 or more, children belonging to "other" religions, children whose caste/tribe is not known, and children whose mothers were not interviewed but were in the household, who are not shown separately.

¹ Excludes children whose mothers were not interviewed

² For women who are not interviewed, information is taken from the Household Questionnaire. Excludes children whose mothers are not listed in the household schedule.

³ Includes children whose mothers are deceased

⁴ Mildly anaemic is classified as 10.0-11.9 g/dl for non-pregnant women and 10.0-10.9 g/dl for pregnant women. Moderately/severely anaemic is <10.0 g/dl. Adjusted for altitude and for smoking status. Excludes children whose mother's anaemia status is not known.

() Based on 25-49 unweighted cases

Table 66 Micronutrient intake among children

Percentage of youngest children age 9-23 months living with the mother who consumed vitamin A-rich and iron-rich foods in the day or night preceding the survey, percentage of children age 9-59 months who were given vitamin A supplements in the six months preceding the survey and percentage of children age 6-59 months who were given iron supplements in the past seven days, who were given deworming medication in the six months preceding the survey, and who live in households using iodized salt, by background characteristics, Telangana, 2015-16

Background characteristic	Youngest children age 9-23 months living with their mother			Children age 9-59 months			Children age 6-59 months in households with salt tested			
	Percentage who consumed foods rich in vitamin A in past 24 hours ¹	Percentage who consumed foods rich in iron in past 24 hours ²	Number of children	Percentage given vitamin A supplements in past 6 months	Number of children	Percentage given iron supplements in past 7 days	Percentage given deworming medication in past 6 months ³	Number of children	Percentage living in households using iodized salt ⁴	Number of children
Age in months										
6-8	na	na	na	na	na	37.3	25.9	120	96.5	128
9-11	29.6	16.7	112	81.6	112	45.3	30.3	112	93.9	122
12-17	48.7	27.5	210	89.3	218	35.7	25.3	218	97.7	231
18-23	56.1	39.7	249	85.0	277	39.7	28.4	277	94.7	295
24-35	na	na	na	76.1	448	38.8	20.8	448	95.4	481
36-47	na	na	na	73.7	510	35.6	26.0	510	94.9	548
48-59	na	na	na	67.0	474	36.4	27.1	474	96.0	508
Sex										
Male	49.5	30.5	310	75.7	1,100	37.0	25.5	1,164	95.8	1,243
Female	46.6	30.9	261	77.0	939	38.3	25.7	994	95.2	1,070
Birth order										
1	52.0	30.7	237	78.4	906	37.7	25.8	953	97.2	917
2-3	45.5	30.5	318	74.9	1,066	38.5	25.9	1,138	95.1	1,111
4-5	*	*	15	70.1	63	21.7	20.1	64	93.5	62
Breastfeeding status										
Breastfeeding	46.7	27.7	475	82.3	682	39.9	26.1	801	95.0	785
Not breastfeeding	55.6	45.6	96	73.3	1,357	36.2	25.3	1,358	96.4	1,310
Residence										
Urban	52.2	30.8	275	75.3	983	36.5	24.9	1,033	99.1	1,082
Rural	44.5	30.5	296	77.2	1,056	38.5	26.3	1,126	92.4	1,231

Continued...

Table 66 Micronutrient intake among children—Continued

Percentage of youngest children age 9-23 months living with the mother who consumed vitamin A-rich and iron-rich foods in the day or night preceding the survey, percentage of children age 9-59 months who were given vitamin A supplements in the six months preceding the survey and percentage of children age 6-59 months who were given iron supplements in the past seven days, who were given deworming medication in the six months preceding the survey, and who live in households using iodized salt, by background characteristics, Telangana, 2015-16

Background characteristic	Youngest children age 9-23 months living with their mother				Children age 6-59 months			Children age 6-59 months in households with salt tested	
	Percentage who consumed foods rich in vitamin A in past 24 hours ¹	Percentage who consumed foods rich in iron in past 24 hours ²	Number of children	Percentage given vitamin A supplements in past 6 months	Number of children	Percentage iron supplements given in past 7 days	Percentage given deworming medication in past 6 months ³	Number of children	Percentage living in households using iodized salt ⁴
Mother's schooling									
No schooling	35.4	25.2	97	69.0	405	34.4	24.5	421	89.0
<5 years complete	*	*	22	66.4	96	36.2	20.7	101	91.7
5-9 years complete	50.6	36.2	120	77.8	471	38.6	25.7	499	96.1
10-11 years complete	49.2	35.7	143	75.9	469	36.8	23.8	503	97.7
12 or more years complete	52.1	25.1	189	82.0	599	39.7	28.6	635	98.6
Religion									
Hindu	47.0	29.3	502	76.6	1,724	38.3	26.4	1,826	94.9
Muslim	52.0	41.9	49	75.4	252	30.3	19.7	267	99.0
Christian	*	*	19	70.9	60	44.9	25.8	63	96.6
Caste/tribe									
Scheduled caste	47.7	33.2	114	75.1	372	37.1	25.1	396	95.4
Scheduled tribe	53.5	21.6	45	75.9	165	35.0	21.3	175	90.7
Other backward class	46.7	34.3	325	77.9	1,168	37.0	25.7	1,235	95.4
Other	53.0	19.0	84	72.8	325	41.6	28.5	341	99.0
Total	48.2	30.7	571	76.3	2,039	37.6	25.6	2,159	95.5

Note: Information on iron supplements and deworming medication is based on the mother's recall. Information on vitamin A supplementation is based on the vaccination card (where available) and mother's recall. Total includes children of birth order 6 or more, children belonging to "other" religions, and children whose caste/tribe is not known, who are not shown separately.

na = Not applicable

¹ Includes meat and organ meats, fish, poultry, eggs, pumpkin, carrots, squash, and sweet potatoes that are yellow or orange inside, dark green leafy vegetables, ripe mango, cantaloupe, and jackfruit

² Includes meat and organ meats, fish, poultry, or eggs

³ Deworming for intestinal worms

⁴ Excludes children in households in which salt was not tested. Includes children whose mothers were not interviewed, except for birth order and breastfeeding status and mother's schooling. For mother's schooling, excludes children whose mothers are not listed in the household schedule.

* Percentage not shown; based on fewer than 25 unweighted cases

Table 67 Presence of iodized salt in household

Percentage of households with salt tested for iodine content that have iodized salt, according to background characteristics, Telangana, 2015-16

Background characteristic	Among households with tested salt:	
	Percentage with iodized salt	Number of households
Residence		
Urban	99.2	3,440
Rural	93.0	4,284
Religion of household head		
Hindu	95.5	6,749
Muslim	98.1	719
Christian	96.9	243
Caste/tribe of household head		
Scheduled caste	95.8	1,454
Scheduled tribe	90.9	597
Other backward class	95.7	4,375
Other	98.4	1,230
Don't know	96.3	67
Total	95.8	7,723

Note: Total includes households with household head belonging to "other" religions, which are not shown separately.

Table 68 Presence of iodized salt in household by district

Percentage of households with salt tested for iodine content that have iodized salt, by district, Telangana, 2015-16

District	Among households with tested salt:	
	Percentage with iodized salt	Number of households
Adilabad	97.1	573
Hyderabad	98.9	1,036
Karimnagar	98.8	869
Khammam	99.2	655
Mahbubnagar	80.4	701
Medak	98.2	568
Nalgonda	90.4	761
Nizamabad	95.4	497
Rangareddy	98.4	1,298
Warangal	97.4	767
Telangana	95.8	7,723

Table 69 Women's and men's food consumption

Percent distribution of women and men age 15-49 by frequency of consumption of specific foods, Telangana, 2015-16

Type of food	Frequency of consumption				Total	Number of respondents
	Daily	Weekly	Occasionally	Never		
WOMEN						
Milk or curd	55.6	18.8	17.4	8.3	100.0	7,567
Pulses or beans	27.7	62.2	9.4	0.7	100.0	7,567
Dark green leafy vegetables	13.2	64.8	20.7	1.3	100.0	7,567
Fruits	11.5	36.2	49.3	3.1	100.0	7,567
Eggs	6.2	54.3	31.3	8.2	100.0	7,567
Fish	0.6	22.9	53.8	22.7	100.0	7,567
Chicken or meat	1.5	51.3	37.2	10.0	100.0	7,567
Fish or chicken or meat	1.8	55.6	34.1	8.6	100.0	7,567
Fried foods	4.5	26.6	60.3	8.5	100.0	7,567
Aerated drinks	1.6	17.3	65.3	15.7	100.0	7,567
MEN						
Milk or curd	50.8	33.4	10.9	4.9	100.0	1,058
Pulses or beans	25.5	65.3	8.6	0.5	100.0	1,058
Dark green leafy vegetables	17.7	66.2	15.6	0.5	100.0	1,058
Fruits	8.5	51.7	38.4	1.4	100.0	1,058
Eggs	4.0	73.9	19.0	3.2	100.0	1,058
Fish	0.6	42.8	48.5	8.1	100.0	1,058
Chicken or meat	0.5	67.5	28.3	3.8	100.0	1,058
Fish or chicken or meat	0.8	72.8	23.6	2.8	100.0	1,058
Fried foods	2.7	30.7	52.1	14.5	100.0	1,058
Aerated drinks	5.7	37.0	49.9	7.4	100.0	1,058

Table 70 Nutritional status of adults

Percentage of women and men age 15-49 with specific body mass index (BMI) levels, by background characteristics, Telangana, 2015-16

Background characteristic	Body mass index: Women ¹					Body mass index: Men				
	<18.5 (total thin)	<17.0 (moderately/severely thin)	≥25.0 (over-weight or obese)	≥30.0 (obese)	Number of women	<18.5 (total thin)	<17.0 (moderately/severely thin)	≥25.0 (over-weight or obese)	≥30.0 (obese)	Number of men
Age										
15-19	48.5	25.5	7.3	1.1	883	54.9	35.9	4.7	0.5	161
20-29	28.0	12.6	17.0	4.4	2,363	21.1	6.3	22.2	2.1	310
30-39	15.1	6.7	37.8	11.6	1,883	12.5	3.7	31.9	7.8	267
40-49	11.3	5.0	44.5	14.5	1,598	8.7	2.2	31.7	7.8	227
Marital status										
Never married	44.7	22.9	7.8	1.4	1,248	38.9	21.7	12.7	0.7	347
Currently married	18.1	8.0	33.0	9.9	4,963	11.6	2.6	31.1	7.1	606
Widowed/divorced/separated/ deserted	19.0	9.0	30.5	10.9	516	*	*	*	*	13
Residence										
Urban	16.1	7.8	39.5	13.9	3,067	17.6	7.5	31.9	6.7	439
Rural	29.0	13.4	18.5	3.8	3,659	24.6	11.3	17.9	3.1	527
Schooling										
No schooling	21.4	10.1	27.0	7.5	2,303	19.1	4.8	16.4	2.0	172
<5 years complete	23.0	9.4	29.2	7.5	319	20.4	6.7	28.5	0.0	42
5-9 years complete	24.0	11.1	30.6	9.4	1,345	26.0	13.4	21.2	7.1	237
10-11 years complete	26.3	13.2	27.1	9.6	1,241	24.6	10.9	24.0	5.9	227
12 or more years complete	22.3	10.1	28.2	8.0	1,518	16.7	8.6	31.0	4.2	288
Religion										
Hindu	24.0	11.1	26.6	7.4	5,820	22.1	10.0	23.3	4.5	838
Muslim	15.8	8.4	39.9	15.4	671	15.4	6.5	29.5	3.5	102
Christian	21.0	10.6	31.2	11.5	223	(23.9)	(6.2)	(33.3)	(16.9)	27
Caste/tribe										
Scheduled caste	25.1	13.1	23.4	5.6	1,274	28.5	13.4	22.9	4.9	142
Scheduled tribe	29.2	12.2	18.2	5.3	546	23.7	6.6	12.5	3.0	86
Other backward class	23.5	10.9	27.4	7.7	3,902	21.2	10.2	23.4	4.0	612
Other	15.5	6.8	42.6	16.2	961	12.8	4.3	37.5	9.3	126
Don't know	18.1	9.7	31.3	14.7	44	nc	nc	nc	nc	0
Total age 15-49	23.1	10.8	28.1	8.4	6,726	21.4	9.6	24.2	4.7	966
Age 50-54	na	na	na	na	na	19.5	7.1	20.7	6.4	69
Total age 15-54	na	na	na	na	na	21.3	9.4	24.0	4.8	1,035

Note: The body mass index (BMI) is expressed as the ratio of weight in kilograms to the square of height in meters (kg/m²). Total includes women/men belonging to "other" religions, who are not shown separately.

nc = No cases

na = Not applicable

¹ Excludes pregnant women and women with a birth in the preceding 2 months

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

Table 71 Prevalence of anaemia in adults

Percentage of women and men age 15-49 with anaemia by background characteristics, Telangana, 2015-16

Background characteristic	Women					Men				
	Mild (10.0-11.9 g/dl) ¹	Moderate (7.0- 9.9 g/dl)	Severe (<7.0 g/dl)	Any anaemia (<12.0 g/dl) ²	Number of women	Mild (12.0- 12.9 g/dl)	Moderate (9.0- 11.9 g/dl)	Severe (<9.0 g/dl)	Any anaemia (<13.0 g/dl)	Number of men
Age										
15-19	38.5	17.7	3.6	59.8	903	7.0	11.3	1.0	19.3	160
20-29	36.6	16.9	1.8	55.3	2,513	7.2	3.3	1.9	12.5	304
30-39	37.3	18.2	3.5	58.9	1,866	7.1	5.9	0.8	13.7	269
40-49	35.3	17.4	2.0	54.6	1,584	9.8	7.7	1.0	18.6	226
Marital status										
Never married	39.1	17.9	4.0	61.0	1,214	6.3	6.7	1.6	14.6	341
Currently married	36.3	16.7	2.3	55.3	5,138	8.5	6.3	1.0	15.8	606
Widowed/divorced/ separated/ deserted	35.1	24.0	1.7	60.8	514	*	*	*	*	13
Maternity status										
Pregnant	21.4	26.7	1.6	49.8	228	na	na	na	na	na
Breastfeeding	38.1	12.6	0.8	51.5	852	na	na	na	na	na
Neither	37.1	17.8	2.8	57.8	5,785	na	na	na	na	na
Residence										
Urban	36.0	16.5	2.6	55.0	3,113	5.8	3.8	0.6	10.2	437
Rural	37.3	18.3	2.5	58.1	3,753	9.4	8.6	1.7	19.8	523
Schooling										
No schooling	36.8	17.6	2.8	57.2	2,311	9.9	11.3	1.7	22.9	170
<5 years complete	35.4	19.6	1.7	56.7	323	13.8	1.2	0.0	15.0	42
5-9 years complete	37.1	18.5	2.5	58.2	1,382	9.7	6.7	1.7	18.1	238
10-11 years complete	38.6	17.2	2.4	58.2	1,291	3.6	7.8	1.8	13.2	227
12 or more years complete	34.9	16.1	2.4	53.4	1,560	7.3	3.0	0.2	10.5	282
Religion										
Hindu	36.8	17.7	2.6	57.2	5,936	8.2	6.3	1.4	15.9	832
Muslim	33.3	14.5	1.8	49.6	686	4.7	4.6	0.0	9.3	101
Christian	41.3	20.5	2.6	64.5	231	(6.5)	(17.7)	(0.0)	(24.3)	27
Caste/tribe										
Scheduled caste	36.3	20.1	3.0	59.4	1,301	7.8	9.2	0.7	17.7	141
Scheduled tribe	42.5	21.0	2.7	66.2	557	9.9	13.5	0.0	23.4	84
Other backward class	37.0	17.2	2.6	56.8	3,999	7.0	4.9	1.5	13.4	611
Other	32.6	12.9	1.8	47.3	965	10.1	5.9	1.1	17.1	123
Don't know	39.4	14.2	1.1	54.7	44	nc	nc	nc	nc	0
Total age 15-49	36.7	17.5	2.5	56.7	6,866	7.8	6.4	1.2	15.4	959
Age 50-54	na	na	na	na	na	13.2	8.6	1.4	23.2	67
Total age 15-54	na	na	na	na	na	8.1	6.6	1.2	15.9	1,026

Note: Table is based on women and men who stayed in the household the night before the interview. Prevalence is adjusted for altitude and for smoking status, if known, using the CDC formulas (Centers for Disease Control (CDC). 1998. Recommendations to prevent and control iron deficiency in the United States. *Morbidity and Mortality Weekly Report* 47 (RR-3): 1-29). Total include women/men belonging to "other" religions, who are not shown separately.

Haemoglobin in g/dl = grams per decilitre.

nc = No cases

na = Not applicable

¹ For pregnant women the value is 10.0-10.9 g/dl

² For pregnant women the value is <11.0 g/dl

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

Table 72 Nutritional status and anaemia among children and women by district

Percentage of children age 6-59 months classified as having anaemia and percentage of women age 15-49 with anaemia and specific body mass index (BMI) levels by district, Telangana, 2015-16

District	Percentage of children having any anaemia (<11.0 g/dl)	Number of children	Percentage of women having any anaemia (<12.0 g/dl) ¹	Number of women	Women with BMI <18.5 (total thin)	Women with BMI ≥25.0 (overweight or obese)	Number of women ²
Adilabad	67.8	121	63.7	519	35.1	15.8	505
Hyderabad	54.9	267	54.0	976	12.9	47.9	960
Karimnagar	52.1	170	52.3	774	23.3	24.8	768
Khammam	72.2	143	71.1	589	20.2	27.3	567
Mahbubnagar	64.4	211	50.6	618	28.3	19.8	587
Medak	67.5	125	57.3	514	29.0	20.2	511
Nalgonda	69.1	198	56.2	630	24.9	25.0	613
Nizamabad	65.9	127	56.0	464	26.5	21.2	449
Rangareddy	48.2	294	54.8	1,124	19.3	33.5	1,119
Warangal	63.0	139	56.9	657	24.3	24.9	647
Telangana	60.7	1,795	56.7	6,866	23.1	28.1	6,726

Note: Table is based on children/women who stayed in the household the night before the interview. Prevalence of anaemia, based on haemoglobin levels, is adjusted for altitude (and smoking status for women) using the CDC formulas (Centers for Disease Control (CDC). 1998. Recommendations to prevent and control iron deficiency in the United States. *Morbidity and Mortality Weekly Report* 47 (RR-3): 1-29). Haemoglobin levels shown in grams per decilitre (g/dl). The body mass index (BMI) is expressed as the ratio of weight in kilograms to the square of height in meters (kg/m²).

¹ For pregnant women, the value is <11.0 g/dl

² Excludes pregnant women and women with a birth in the preceding 2 months

Table 73 Knowledge and prevention of HIV/AIDS

Percentage of women and men age 15-49 who have heard of HIV or AIDS and who, in response to prompted questions, say that people can reduce the risk of getting HIV/AIDS by using a condom every time they have sexual intercourse, who know that the risk of HIV/AIDS can be reduced by limiting sex to one uninfected partner, who have a comprehensive knowledge about HIV/AIDS, and who know that HIV/AIDS can be transmitted from a mother to a baby, by background characteristics, Telangana, 2015-16

Background characteristic	Percentage who have heard of HIV or AIDS		Percentage who know that people can reduce their chances of getting HIV/AIDS by using a condom every time they have sex				Percentage who know that the risk of HIV/AIDS can be reduced by limiting sex to one uninfected partner ¹				Percentage who have a comprehensive knowledge about HIV/AIDS ²				Percentage who know that HIV/AIDS can be transmitted from a mother to her baby		Number	
	Women	Men	Women	Men	Women	Men	Women	Men	Women	Men	Women	Men	Women	Men	Women	Men	Women	Men
Age																		
15-24	94.8	97.6	65.7	83.7	77.5	83.6	34.9	53.8	82.8	90.6	416	329						
15-19	94.2	97.2	62.3	82.1	75.6	81.9	27.8	52.4	79.3	90.3	183	174						
20-24	95.3	98.1	68.4	85.6	78.9	85.5	40.5	55.3	85.5	91.0	233	155						
25-29	90.6	97.7	64.5	86.1	78.1	83.4	32.4	56.3	82.7	85.2	231	184						
30-39	90.0	96.4	56.6	84.7	68.3	83.3	28.4	51.7	78.7	87.1	332	296						
40-49	83.4	91.7	48.1	70.3	66.0	67.5	21.3	37.5	73.5	79.3	306	248						
Residence																		
Urban	94.8	96.1	68.6	85.4	80.8	83.7	39.1	55.1	86.2	90.9	614	506						
Rural	85.7	95.8	50.1	77.5	64.8	76.0	20.8	45.0	73.3	81.6	671	552						
Schooling																		
No schooling	77.7	86.1	37.5	58.3	51.0	61.0	14.7	30.8	64.9	65.1	440	179						
<5 years complete	88.9	92.8	56.4	67.5	72.1	70.4	19.4	33.9	75.9	71.3	58	44						
5-9 years complete	92.1	97.3	56.9	78.7	76.0	77.7	24.3	45.5	80.8	86.8	255	257						
10-11 years complete	98.8	99.4	73.8	89.4	87.3	84.8	36.3	51.8	87.6	91.8	233	245						
12 or more years complete	100.0	98.0	81.2	91.4	89.5	88.7	52.4	64.1	94.2	94.4	299	333						
Regular media exposure³																		
Yes	91.3	96.4	61.4	82.4	75.3	80.7	31.1	50.5	81.3	86.7	1,179	1,014						
No	76.5	85.8	32.5	54.1	41.8	56.8	12.5	34.2	59.9	70.4	107	44						
Marital status																		
Never married	94.0	97.1	64.6	85.1	78.5	85.1	33.6	57.3	82.4	91.9	255	380						
Currently married	89.6	95.3	58.7	79.2	71.5	76.4	29.4	45.8	79.6	82.7	938	665						
Widowed/divorced/separated/deserted	84.5	*	46.6	*	66.0	*	19.7	*	70.7	*	92	13						

Continued...

Table 73 Knowledge and prevention of HIV/AIDS—Continued

Percentage of women and men age 15-49 who have heard of HIV or AIDS and who, in response to prompted questions, say that people can reduce the risk of getting HIV/AIDS by using a condom every time they have sexual intercourse, who know that the risk of HIV/AIDS can be reduced by limiting sex to one uninfected partner, who have a comprehensive knowledge about HIV/AIDS, and who know that HIV/AIDS can be transmitted from a mother to a baby, by background characteristics, Telangana, 2015-16

Background characteristic	Percentage who have heard of HIV or AIDS		Percentage who know that people can reduce their chances of getting HIV/AIDS by using a condom every time they have sex		Percentage who know that the risk of HIV/AIDS can be reduced by limiting sex to one uninfected partner ¹		Percentage who have a comprehensive knowledge about HIV/AIDS ²		Percentage who know that HIV/AIDS can be transmitted from a mother to her baby		Number	
	Women	Men	Women	Men	Women	Men	Women	Men	Women	Men	Women	Men
Whether been away from home for one month or more at a time in the past 12 months⁴												
Been away	90.6	98.8	66.3	85.8	73.0	84.4	23.9	50.4	83.1	93.4	124	134
Not been away	90.0	95.5	58.2	80.6	72.4	79.0	30.1	49.8	79.1	85.0	1,161	924
Whether been away from home for six months or more at a time in the past 12 months⁴												
Been away	93.8	98.0	68.7	84.8	78.8	79.7	28.4	50.9	89.7	94.7	66	80
Not been away	89.9	95.7	58.4	81.0	72.1	79.7	29.6	49.7	78.9	85.3	1,219	977
Religion												
Hindu	89.7	96.2	58.0	80.9	72.2	80.7	28.7	49.4	79.5	85.5	1,133	910
Muslim	93.0	93.1	67.9	82.9	73.3	72.3	39.9	49.7	78.2	90.6	121	120
Christian	(96.7)	(100.0)	(64.5)	(84.4)	(84.6)	(77.9)	(24.0)	(65.6)	(90.6)	(84.9)	27	27
Caste/tribe												
Scheduled caste	88.1	93.5	51.0	76.1	67.5	76.4	23.9	50.2	78.9	78.7	198	160
Scheduled tribe	82.2	95.3	39.5	72.8	53.6	66.2	20.2	42.5	69.1	87.3	108	87
Other backward class	90.5	96.3	60.2	81.9	73.2	80.5	29.5	48.6	80.0	86.7	796	666
Other	97.6	97.3	75.5	89.0	88.0	87.6	42.1	59.7	86.5	90.5	175	145
Total	90.1	95.9	59.0	81.3	72.5	79.7	29.5	49.8	79.5	86.0	1,285	1,058

Note: Total includes women/men belonging to "other" religions and women/men who don't know their caste/tribe, who are not shown separately.

¹ Partner who has no other sex partners

² Comprehensive knowledge means knowing that consistent use of condoms every time they have sex and having just one uninfected faithful sex partner can reduce the chance of getting HIV/AIDS, knowing that a healthy-looking person can have HIV/AIDS, and rejecting two common misconceptions about transmission or prevention of HIV/AIDS

³ Exposure to radio, television, or newspapers/magazines at least once a week or cinema at least once a month

⁴ For women, visits to parental/in-laws' home excluded

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

Table 7.4. Accepting attitudes toward those living with HIV/AIDS

Among women and men age 15-49 who have heard of HIV/AIDS, percentage expressing specific accepting attitudes toward people with HIV/AIDS, by background characteristics, Telangana, 2015-16

Background characteristic	Percentage of women who:				Percentage of men who:				Number of men who have heard of HIV/AIDS			
	Are willing to care for a relative with HIV/AIDS in own home	Would buy fresh vegetables from a shopkeeper or vendor who has HIV/AIDS	Say that a female teacher who has HIV/AIDS but is not sick should be allowed to continue teaching	Would not want to keep secret that a family member got infected with HIV/AIDS	Express accepting attitudes on all four indicators	Number of women who have heard of HIV/AIDS	Are willing to care for a relative with HIV/AIDS in own home	Would buy fresh vegetables from a shopkeeper or vendor who has HIV/AIDS		Say that a female teacher who has HIV/AIDS but is not sick should be allowed to continue teaching	Would not want to keep secret that a family member got infected with HIV/AIDS	Express accepting attitudes on all four indicators
Age												
15-24	70.6	67.7	76.8	34.7	17.5	395	67.3	66.9	72.9	24.2	9.7	321
15-19	67.2	60.3	72.4	33.6	16.2	172	71.6	66.0	73.5	25.2	12.0	169
20-24	73.2	73.5	80.1	35.5	18.5	222	62.5	68.0	72.2	23.1	7.1	152
25-29	61.4	66.2	69.7	38.6	16.7	209	58.8	65.7	73.6	29.8	11.0	180
30-39	62.4	58.6	70.8	38.4	12.1	299	70.0	65.5	69.4	31.4	12.7	285
40-49	62.8	52.8	63.3	37.8	14.5	255	67.5	59.3	67.7	30.1	10.0	228
Residence												
Urban	70.3	71.7	77.5	37.1	19.7	583	70.7	71.8	77.4	26.2	10.8	486
Rural	59.8	51.8	64.3	37.0	10.8	575	62.8	57.9	64.8	30.8	10.8	529
Schooling												
No schooling	52.5	43.1	55.2	38.7	7.2	341	57.6	45.8	54.7	36.3	6.8	154
<5 years complete	62.9	61.0	61.0	48.9	15.8	52	(64.3)	(41.5)	(51.1)	(28.3)	(1.5)	41
5-9 years complete	61.7	59.3	70.5	37.5	15.4	235	59.6	56.0	64.4	29.8	8.9	250
10-11 years complete	71.2	66.6	76.4	34.6	19.1	230	68.8	65.8	72.3	25.4	10.4	244
12 or more years complete	77.8	81.5	86.8	34.6	21.4	299	74.8	82.1	84.9	26.3	15.7	326
Regular media exposure¹												
Yes	66.4	63.1	72.5	37.1	16.0	1,076	67.0	65.3	71.4	28.4	11.0	977
No	48.3	44.7	50.8	36.6	6.5	82	(55.5)	(46.4)	(56.7)	(31.8)	(6.1)	37
Marital status												
Never married	76.2	70.2	80.7	33.0	19.7	239	70.3	69.9	75.3	24.0	10.2	369
Currently married	62.3	60.6	69.3	37.5	14.7	840	64.6	61.3	68.4	31.3	11.2	634
Widowed/divorced/separated/deserted	61.2	49.3	59.0	44.7	8.0	78	*	*	*	*	*	12

Continued...

Table 74 Accepting attitudes toward those living with HIV/AIDS—Continued

Among women and men age 15-49 who have heard of HIV/AIDS, percentage expressing specific accepting attitudes toward people with HIV/AIDS, by background characteristics, Telangana, 2015-16

Background characteristic	Percentage of women who:				Percentage of men who:					
	Are willing to care for a relative with HIV/AIDS in own home	Would buy fresh vegetables from a shopkeeper or vendor who has HIV/AIDS	Say that a female teacher who has HIV/AIDS but is not sick should be allowed to continue teaching	Would not want to keep secret that a family member got infected with HIV/AIDS	Are willing to care for a relative with HIV/AIDS in own home	Would buy fresh vegetables from a shopkeeper or vendor who has HIV/AIDS	Say that a female teacher who has HIV/AIDS but is not sick should be allowed to continue teaching	Would not want to keep secret that a family member got infected with HIV/AIDS	Express accepting attitudes on all four indicators	Number of men who have heard of HIV/AIDS
Religion										
Hindu	66.8	60.3	71.0	37.2	65.0	63.5	70.0	29.8	10.6	875
Muslim	53.7	74.8	69.4	36.0	75.9	74.6	80.3	20.0	12.1	112
Christian	(51.7)	(63.5)	(76.2)	(35.4)	(77.7)	(59.3)	(58.3)	(23.1)	(14.0)	27
Caste/tribe										
Scheduled caste	65.4	52.9	67.6	33.4	67.9	62.7	69.4	23.3	9.7	150
Scheduled tribe	51.3	47.9	63.5	49.6	53.8	46.8	53.8	35.3	12.4	83
Other backward class	65.4	63.0	71.6	36.8	65.7	64.7	71.6	30.3	10.5	641
Other	70.8	74.0	75.8	35.2	76.6	76.7	79.0	22.3	12.4	141
Total	65.1	61.8	71.0	37.0	66.6	64.6	70.9	28.6	10.8	1,014

Note: Total includes women/men belonging to "other" religions and women/men who don't know their caste/tribe, who are not shown separately.

¹ Exposure to radio, television, or newspapers/magazines at least once a week or cinema at least once a month

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

Table 75 Sexual behaviour, HIV testing, blood transfusion, and injections

Indicators of higher risk sexual behaviour, use of blood transfusion, prior HIV testing, and any injections for women and men age 15-49 by residence, Telangana, 2015-16

Behaviour	Urban			Rural			Total		
	Male	Female	Total	Male	Female	Total	Male	Female	Total
Among those who had sexual intercourse in the past 12 months:									
Percentage who had two or more partners in the past 12 months	3.2	1.4	2.1	4.8	1.1	2.6	4.1	1.2	2.4
Percentage who had higher-risk intercourse in the past 12 months ¹	4.5	0.8	2.4	6.3	0.4	2.8	5.5	0.6	2.6
Percentage who had two or more partners and higher-risk intercourse in the past 12 months ¹	1.7	0.8	1.2	3.5	0.4	1.6	2.7	0.6	1.4
Number who had sexual intercourse in the past 12 months	232	323	556	286	419	705	519	742	1,261
Among those who had higher-risk sexual intercourse in the past 12 months:									
Percentage who reported using a condom at last higher-risk intercourse ¹	*	*	*	*	*	*	(23.9)	*	(28.8)
Number who had higher-risk sexual intercourse in the past 12 months	11	3	13	18	2	20	29	4	33
Among those who have ever had sexual intercourse:									
Mean number of sexual partners in lifetime	1.7	2.0	1.9	1.9	2.3	2.1	1.8	2.2	2.0
Number who have ever had sexual intercourse	240	396	636	332	524	856	572	919	1,491
Among all men									
Percentage who paid for sexual intercourse in the past 12 months	0.6	na	na	0.7	na	na	0.6	na	na
Number of men	506	na	na	552	na	na	1,058	na	na
Among women with a birth in the past five years who received ANC during pregnancy, percentage who were tested for HIV during ANC									
	na	47.4	na	na	33.7	na	na	39.8	na
Number of women with a birth in the past five years who received ANC during pregnancy									
	na	129	na	na	161	na	na	290	na
Percentage ever tested for HIV prior to NFHS-4	11.9	36.0	25.1	13.7	28.0	21.5	12.8	31.8	23.3
Percentage who have ever had a blood transfusion	12.7	8.3	10.3	11.4	12.0	11.7	12.0	10.2	11.0
Percentage who received any injection in the past 12 months	45.4	53.8	50.0	62.3	67.2	65.0	54.2	60.8	57.8
Mean number of injections in the past 12 months	4.1	5.3	4.8	5.1	6.1	5.7	4.7	5.8	5.3
Number of respondents	506	614	1,120	552	671	1,223	1,058	1,285	2,343
Among those who received an injection in the past 12 months, percentage for whom for the last injection, a disposable syringe was used									
	53.7	72.2	64.7	56.2	78.9	69.1	55.2	76.1	67.2
Number who received an injection in the past 12 months	229	331	560	344	451	794	573	781	1,354

na = Not applicable

ANC = Antenatal care

¹ Sexual intercourse with a partner who was neither a spouse nor who lived with the respondent

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

Table 76 Knowledge of HIV/AIDS and sexual behaviour among youth

Indicators of HIV/AIDS knowledge and sexual behaviour for women and men age 15-24 by residence, Telangana, 2015-16

Knowledge and behaviour	Urban			Rural			Total		
	Male	Female	Total	Male	Female	Total	Male	Female	Total
Knowledge									
Percentage with comprehensive knowledge about HIV/AIDS ¹	57.0	42.5	48.8	51.0	27.8	38.2	53.8	34.9	43.3
Percentage who know a condom source	81.1	33.9	54.3	80.1	21.9	48.0	80.5	27.7	51.0
Sexual behaviour									
Percentage who have ever had sexual intercourse	12.5	30.4	22.6	18.2	50.6	36.0	15.6	40.9	29.7
Percentage who had sexual intercourse before age 15	0.7	0.6	0.7	1.5	3.2	2.4	1.1	1.9	1.6
HIV testing, injections and blood transfusion									
Percentage ever tested for HIV prior to NFHS-4	5.7	25.2	16.7	8.5	35.8	23.5	7.2	30.7	20.3
Percentage who have ever had a blood transfusion	10.6	4.7	7.2	8.2	9.8	9.1	9.3	7.3	8.2
Percentage who received any injection in the past 12 months	43.9	51.0	48.0	68.0	62.1	64.8	56.8	56.7	56.8
Mean number of injections in the past 12 months	3.7	6.5	5.4	4.8	4.2	4.5	4.4	5.2	4.9
Number age 15-24	153	200	354	176	216	392	329	416	745
Among those who received an injection in the past 12 months, percentage for whom a disposable syringe was used	(55.7)	66.7	62.4	59.1	81.4	70.9	57.9	75.1	67.5
Number who received an injection in the past 12 months	67	102	170	120	134	254	187	236	423
Among those who have ever had sexual intercourse, percentage who used a condom at first sexual intercourse	*	7.5	7.4	(22.4)	7.0	10.2	16.2	7.2	9.1
Number who have ever had sexual intercourse	22	76	98	33	123	156	55	199	254
Among those who had sexual intercourse in the past 12 months, percentage who had higher-risk sexual intercourse in the past 12 months ²	*	0.0	4.4	(30.8)	0.0	6.5	(27.2)	0.0	5.7
Number who had sexual intercourse in the past 12 months	15	56	71	24	91	116	39	147	186
Among the never married:									
Percentage who have never had sexual intercourse	95.1	92.5	93.8	90.1	94.0	91.6	92.4	93.1	92.7
Percentage who had sexual intercourse in the past 12 months	1.9	0.0	1.0	4.9	0.0	3.1	3.5	0.0	2.0
Number never married	138	134	272	159	97	256	297	231	528

¹ Comprehensive knowledge means knowing that consistent use of condoms every time they have sex and having just one uninfected faithful sex partner can reduce the chance of getting HIV/AIDS, knowing that a healthy-looking person can have HIV/AIDS, and rejecting two common misconceptions about transmission or prevention of HIV/AIDS

² Sexual intercourse with a partner who was neither a spouse nor lived with the respondent

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

Table 77 Prevalence of tuberculosis

Number of persons per 100,000 usual household residents suffering from any tuberculosis and medically treated tuberculosis by age, sex, and main type of cooking fuel, according to residence, Telangana, 2015-16

Age and sex	Number of persons per 100,000 suffering from:		
	Tuberculosis ¹	Medically treated tuberculosis ²	Number of usual residents
URBAN			
Sex			
Female	109	94	6,890
Male	207	207	7,012
Age			
<15	80	80	3,481
15-59	176	165	9,272
60 +	257	257	1,149
Cooking fuel			
Solid fuel ³	591	591	962
Other fuel	127	118	12,940
Total	159	151	13,901
RURAL			
Sex			
Female	258	238	8,256
Male	600	588	8,183
Age			
<15	49	49	4,048
15-59	338	317	10,087
60 +	1,493	1,469	2,303
Cooking fuel			
Solid fuel ³	548	517	8,280
Other fuel	307	307	8,159
Total	428	413	16,439
TOTAL			
Sex			
Female	190	173	15,145
Male	419	412	15,195
Age			
<15	63	63	7,529
15-59	260	244	19,359
60 +	1,082	1,065	3,452
Cooking fuel			
Solid fuel ³	553	524	9,242
Other fuel	196	191	21,098
Total	305	293	30,340

¹ Includes medically treated tuberculosis

² Suffering from tuberculosis and received medical treatment

³ Includes coal, lignite, charcoal, wood, straw/shrubs/grass, agricultural crop waste, and dung cakes

Table 78 Knowledge and attitudes toward tuberculosis

Background characteristic	Among women who have heard of TB, percentage who:				Among men who have heard of TB, percentage who:								
	Percentage of women who have heard of TB	Number of women	Report that TB is spread through the air by coughing or sneezing	Have misconceptions about transmission of TB ¹	Believe that TB can be cured	Would want a family member's TB kept secret	Number of women who have heard of TB	Percentage of men who have heard of TB	Report that TB is spread through the air by coughing or sneezing	Have misconceptions about transmission of TB	Believe that TB can be cured	Would want a family member's TB kept secret	Number of men who have heard of TB
Age													
15-19	75.9	997	77.2	45.2	82.3	43.9	756	68.0	71.1	47.6	88.9	67.2	119
20-34	73.9	3,825	73.2	41.7	82.1	41.3	2,827	75.0	66.5	47.1	82.9	58.5	371
35-49	67.9	2,745	72.4	42.6	82.7	41.4	1,863	71.3	58.6	50.9	84.3	56.0	277
Residence													
Urban	79.7	3,619	77.8	41.9	87.5	37.2	2,886	74.4	70.5	54.8	87.4	61.9	376
Rural	64.9	3,948	68.6	43.3	76.5	46.8	2,560	70.7	58.4	42.6	81.4	56.0	390
Schooling													
No schooling	56.7	2,458	61.4	37.5	73.8	44.0	1,393	59.0	53.8	42.4	72.9	50.6	106
<5 years complete	66.9	337	64.8	36.1	71.7	43.8	226	59.1	(44.6)	(48.2)	(80.9)	(50.7)	26
5-9 years complete	74.6	1,493	72.9	40.1	81.6	44.0	1,114	66.8	51.4	52.0	77.9	53.9	171
10-11 years complete	78.7	1,435	78.5	44.4	84.3	41.2	1,129	79.2	67.2	43.4	87.9	60.7	194
12 or more years complete	85.9	1,844	82.2	48.3	90.5	38.0	1,584	80.9	76.6	52.6	90.8	64.9	269
Religion													
Hindu	71.4	6,529	73.1	42.8	81.8	42.3	4,662	72.9	63.9	45.8	84.3	56.1	664
Muslim	74.8	774	76.2	38.6	86.0	38.7	579	70.1	69.7	66.5	84.2	76.7	85
Christian	78.5	248	72.8	47.7	83.2	36.7	195	(66.8)	*	*	*	*	18

Continued...

Table 78 Knowledge and attitudes toward tuberculosis – Continued

Background characteristic	Among women who have heard of TB, percentage who:				Among men who have heard of TB, percentage who:							
	Percentage of women who have heard of TB	Report that TB is spread through the air by coughing or sneezing	Have misconceptions about transmission of TB ¹	Believe that TB can be cured	Would want a family member's TB kept secret	Number of women who have heard of TB	Percentage of men who have heard of TB	Report that TB is spread through the air by coughing or sneezing	Have misconceptions about transmission of TB ¹	Believe that TB can be cured	Would want a family member's TB kept secret	Number of men who have heard of TB
Caste/tribe												
Scheduled caste	69.4	71.8	43.5	80.1	42.9	968	68.1	59.2	49.5	86.2	50.0	109
Scheduled tribe	65.6	75.5	36.9	76.2	45.3	392	83.2	62.4	33.3	75.6	64.5	72
Other backward class	72.3	73.4	43.0	83.2	43.0	3,177	70.8	61.7	51.3	83.8	58.9	472
Other	78.0	75.6	41.8	84.5	34.1	888	78.4	81.6	46.1	90.3	64.0	113
Don't know	47.5	(41.3)	(65.3)	(70.4)	(27.9)	21	nc	nc	nc	nc	nc	0
Total	72.0	73.5	42.5	82.3	41.7	5,446	72.5	64.3	48.6	84.3	58.9	766

Note: Total includes women/men belonging to "other" religions, who are not shown separately.

nc = No cases

¹ Misconceptions about the transmission of TB include that it can be spread by sharing utensils, by touching a person with TB, through food, through sexual contact, through mosquito bites, or through any other means except through the air when coughing or sneezing

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

Table 79 Health insurance coverage among women and men

Percentage of women and men age 15-49 who are covered by any health scheme or health insurance by background characteristics, Telangana, 2015-16

Background characteristic	Percentage of women covered by any health scheme or health insurance	Number of women	Percentage of men covered by any health scheme or health insurance	Number of men
Age				
15-19	66.9	997	74.6	174
20-24	47.6	1,355	70.7	155
25-34	56.3	2,470	64.3	339
35-49	68.3	2,745	74.3	389
Residence				
Urban	49.0	3,619	62.1	506
Rural	71.0	3,948	78.4	552
Schooling				
No schooling	73.3	2,458	79.2	179
<5 years complete	71.7	337	78.2	44
5-9 years complete	58.4	1,493	70.1	257
10-11 years complete	56.8	1,435	70.5	245
12 or more years complete	45.9	1,844	65.4	333
Religion				
Hindu	61.9	6,529	71.3	910
Muslim	48.0	774	71.1	120
Christian	62.6	248	(44.5)	27
Caste/tribe				
Scheduled caste	66.9	1,395	71.0	160
Scheduled tribe	70.3	597	87.0	87
Other backward class	60.3	4,393	71.5	666
Other	48.0	1,137	56.0	145
Don't know	62.0	44	nc	0
Total age 15-49	60.5	7,567	70.6	1,058
Age 50-54	na	na	72.7	75
Total age 15-54	na	na	70.7	1,133

Note: Total includes women/men belonging to "other" religions, who are not shown separately.

na = Not applicable

nc = No cases

() Based on 25-49 unweighted cases

Table 80 Source of health care and health insurance coverage among households

Percent distribution of households by the source of health care that household members generally use when they get sick, percentage of households with at least one usual member covered by health insurance or a health scheme, and among households with at least one usual member covered by health insurance or a health scheme, type of health insurance/scheme coverage, according to residence, Telangana, 2015-16

Source	Residence		Total
	Urban	Rural	
Public health sector	26.6	31.3	29.2
Government/municipal hospital	15.4	10.9	12.9
Government dispensary	2.0	1.5	1.7
UHC/UHP/UFWC	2.9	1.0	1.9
CHC/rural hospital/Block PHC	1.9	4.3	3.2
PHC/additional PHC	2.8	11.4	7.5
Sub-centre	0.1	1.1	0.7
Vaidya/hakim/homeopath (AYUSH)	0.0	0.1	0.0
Government mobile clinic	0.0	0.1	0.0
Other public health sector	1.4	1.0	1.2
NGO or trust hospital/clinic	0.4	0.2	0.3
Private health sector	71.7	63.9	67.4
Private hospital	47.9	36.3	41.5
Private doctor/clinic	18.1	17.7	17.9
Private paramedic	1.0	2.2	1.7
Vaidya/hakim/homeopath (AYUSH)	0.3	0.1	0.2
Traditional healer	0.1	0.0	0.0
Pharmacy/drugstore	0.1	0.3	0.2
Other private health sector	4.3	7.3	5.9
Other source	1.2	4.5	3.1
Shop	0.1	0.0	0.0
Home treatment	0.1	0.1	0.1
Other	1.0	4.5	2.9
Total	100.0	100.0	100.0
Health insurance			
Percentage of households in which at least one usual member is covered by a health scheme or health insurance	53.5	76.7	66.4
Number of households	3,465	4,321	7,786
Type of coverage among households in which at least one usual member is covered by a health scheme/health insurance			
Employees' State Insurance Scheme (ESIS)	10.7	0.9	4.4
Central Government Health Scheme (CGHS)	3.2	0.6	1.5
State health insurance scheme	83.6	96.8	92.1
Rashtriya Swasthya Bima Yojana (RSBY)	0.4	1.9	1.3
Community health insurance programme	0.1	0.4	0.3
Other health insurance through employer	1.5	0.1	0.6
Medical reimbursement from employer	1.2	0.2	0.6
Other privately purchased commercial health insurance	2.8	0.5	1.3
Other	0.7	0.1	0.3
Number of households	1,855	3,315	5,170

UHC = Urban health centre; UHP = Urban health post; UFWC = Urban family welfare centre; CHC = Community health centre; PHC = Primary health centre; AYUSH = Ayurveda, yoga and naturopathy, *unani*, *siddha* and homeopathy; ICDS = Integrated Child Development Services; NGO = Nongovernmental organization; TBA = Traditional birth attendant

Table 81 Health problems

Number of women and men age 15-49 per 100,000 who reported that they have diabetes, asthma, goitre or any other thyroid disorder, any heart disease, or cancer, by background characteristics, Telangana, 2015-16

Background characteristic	Number of women per 100,000						Number of men per 100,000					
	Diabetes	Asthma	Goitre or any other thyroid disorder	Any heart disease	Cancer	Number of women	Diabetes	Asthma	Goitre or any other thyroid disorder	Any heart disease	Cancer	Number of men
Age												
15-19	365	1,598	1,596	569	0	997	0	3,087	2,007	348	0	174
20-34	889	2,700	4,040	1,443	35	3,825	881	882	123	668	0	494
35-49	4,617	5,183	6,052	2,326	167	2,745	4,691	1,626	725	3,000	204	389
Residence												
Urban	3,083	4,023	6,127	1,423	59	3,619	2,431	692	918	1,226	0	506
Rural	1,337	2,935	2,909	1,855	95	3,948	1,867	2,277	413	1,698	144	552
Schooling												
No schooling	2,752	4,061	3,981	2,073	89	2,458	3,969	2,135	755	1,764	0	179
<5 years complete	4,691	4,071	6,017	3,729	0	337	2,255	1,416	1,131	2,629	0	44
5-9 years complete	1,742	3,538	4,975	1,505	59	1,493	1,451	2,689	0	1,308	0	257
10-11 years complete	1,742	2,654	3,317	1,942	110	1,435	2,229	1,519	1,049	1,571	325	245
12 or more years complete	1,623	3,092	5,237	589	68	1,844	1,595	298	752	1,217	0	333
Marital status												
Never married	575	1,859	2,273	946	0	1,392	0	1,575	921	393	0	380
Currently married	2,470	3,719	5,034	1,745	88	5,619	3,397	1,383	515	2,117	120	665
Widowed/divorced/separated/deserted	3,163	4,789	3,974	2,425	175	556	*	*	*	*	*	13
Religion												
Hindu	2,080	3,743	4,440	1,687	90	6,529	1,701	1,700	761	1,561	87	910
Muslim	2,962	1,994	3,995	1,161	0	774	4,510	494	0	0	0	120
Christian	2,275	659	5,958	2,255	0	248	(6,193)	(0)	(0)	(5,017)	(0)	27

Continued...

Table 81 Health problems—Continued

Number of women and men age 15-49 per 100,000 who reported that they have diabetes, asthma, goitre or any other thyroid disorder, heart disease, or cancer, by background characteristics, Telangana, 2015-16

Background characteristic	Number of women per 100,000					Number of men per 100,000						
	Diabetes	Asthma	Goitre or any other thyroid disorder	Any heart disease	Cancer	Number of women	Diabetes	Asthma	Goitre or any other thyroid disorder	Any heart disease	Cancer	Number of men
Caste/tribe												
Scheduled caste	1,453	3,003	3,817	1,764	90	1,395	1,701	1,139	414	1,269	0	160
Scheduled tribe	3,316	3,106	4,205	2,322	163	597	798	1,656	1,870	1,795	0	87
Other backward class	2,028	3,642	4,620	1,628	64	4,393	2,151	1,822	311	1,516	119	666
Other	3,013	3,466	4,612	1,234	77	1,137	3,357	460	1,776	1,303	0	145
Don't know	2,139	3,600	6,331	1,570	0	44	nc	nc	nc	nc	nc	0
Total age 15-49	2,172	3,455	4,448	1,648	78	7,567	2,137	1,519	655	1,473	75	1,058
Age 50-54	na	na	na	na	na	na	13,550	6,594	0	2,666	645	75
Total age 15-54	na	na	na	na	na	na	2,897	1,857	611	1,552	113	1,133

Note: Total includes women/men belonging to "other" religions, who are not shown separately.

na = Not applicable

nc = No cases

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

Table 82 Health examinations

Percentage of women age 15-49 who have ever undergone specific health examinations by background characteristics, Telangana, 2015-16

Background characteristic	Type of examination			Number of women
	Cervix	Breast	Oral cavity	
Age				
15-19	5.1	1.7	5.6	997
20-24	25.1	7.8	6.5	1,355
25-34	36.1	10.3	11.2	2,470
35-49	42.3	12.5	13.5	2,745
Residence				
Urban	30.7	11.1	12.5	3,619
Rural	33.7	8.1	8.6	3,948
Schooling				
No schooling	38.7	9.9	10.1	2,458
<5 years complete	44.2	9.5	10.3	337
5-9 years complete	31.6	9.8	10.6	1,493
10-11 years complete	28.7	9.2	9.4	1,435
12 or more years complete	24.8	9.0	11.7	1,844
Religion				
Hindu	33.6	10.1	10.7	6,529
Muslim	20.4	5.3	8.1	774
Christian	33.5	9.1	11.7	248
Caste/tribe				
Scheduled caste	27.5	7.2	8.4	1,395
Scheduled tribe	36.4	8.0	10.7	597
Other backward class	32.8	10.1	10.3	4,393
Other	34.5	11.0	13.8	1,137
Don't know	20.4	7.3	4.8	44
Total	32.3	9.5	10.5	7,567

Note: Total includes women belonging to "other" religions, who are not shown separately.

Table 83.1 Blood pressure status: Women

Among women age 15-49, prevalence of hypertension, percent distribution of blood pressure values, and percentages having normal blood pressure and taking medication to lower blood pressure, by background characteristics, Telangana, 2015-16

Background characteristic	Prevalence of hypertension ¹	Pre-hypertensive			Hypertensive			Total	Percentage with normal blood pressure and taking medicine to lower blood pressure	Number of women
		Normal SBP <120 mmHg and DBP <80 mmHg	SBP 120-139 mmHg or DBP 80-89 mmHg	Stage 1: SBP 140-159 mmHg or DBP 90-99 mmHg	Stage 2: SBP 160-179 mmHg or DBP 100-109 mmHg	Stage 3: SBP ≥180 mmHg or DBP ≥110 mmHg				
Age										
15-19	2.1	93.1	5.8	0.6	0.1	0.4	100.0	0.9	919	
20-24	3.6	83.0	14.1	2.5	0.3	0.3	100.0	0.5	1,239	
25-29	8.6	75.0	18.6	4.6	1.0	0.8	100.0	1.8	1,307	
30-34	11.2	65.3	26.1	6.8	1.5	0.3	100.0	1.0	956	
35-39	16.8	53.7	32.9	10.4	1.9	1.1	100.0	1.3	942	
40-44	27.4	43.5	34.4	16.2	3.7	2.2	100.0	2.5	783	
45-49	32.0	38.8	37.9	15.4	5.1	2.7	100.0	3.6	813	
Residence										
Urban	14.7	64.9	24.3	8.1	1.8	1.0	100.0	1.8	3,170	
Rural	11.9	68.4	22.1	6.7	1.6	1.0	100.0	1.4	3,790	
Schooling										
No schooling	18.9	56.7	28.7	10.3	2.6	1.7	100.0	2.4	2,340	
<5 years complete	14.8	60.8	27.2	6.8	4.4	0.7	100.0	1.3	325	
5-9 years complete	13.0	65.7	23.9	8.2	1.5	0.7	100.0	0.9	1,398	
10-11 years complete	8.7	76.1	17.6	5.1	0.4	0.7	100.0	1.4	1,305	
12 or more years complete	8.3	76.2	17.9	4.2	1.2	0.5	100.0	1.0	1,591	
Religion										
Hindu	12.7	67.5	22.8	7.1	1.6	1.0	100.0	1.6	6,012	
Muslim	15.9	62.5	24.9	9.2	1.9	1.4	100.0	1.1	701	
Christian	17.7	62.9	23.7	8.9	4.2	0.4	100.0	1.3	234	
Caste/tribe										
Scheduled caste	14.0	65.0	24.2	7.9	2.0	0.9	100.0	1.9	1,320	
Scheduled tribe	13.0	69.7	19.9	7.7	2.0	0.7	100.0	1.7	566	
Other backward class	12.7	67.7	22.3	7.1	1.7	1.1	100.0	1.4	4,041	
Other	14.3	64.0	26.4	7.6	1.0	1.0	100.0	1.6	988	
Don't know	9.1	63.5	29.5	3.2	3.9	0.0	100.0	0.0	44	
Total	13.2	66.8	23.1	7.4	1.7	1.0	100.0	1.6	6,959	

Note: Total includes women belonging to "other" religions, who are not shown separately.

SBP = Systolic blood pressure; DBP = Diastolic blood pressure

¹ A woman is classified as having hypertension if she has SBP ≥140 mmHg or DBP ≥90 mmHg at the time of the survey, or she is currently taking antihypertension medication to control her blood pressure. The term hypertension as used in this table is not meant to be a clinical diagnosis of the disease, but rather to provide an indication of the disease burden in the population at the time of the survey.

Table 83.2 Blood pressure status: Men

Among men age 15-49, prevalence of hypertension, percent distribution of blood pressure values, and percentage having normal blood pressure and taking medication to lower blood pressure, by background characteristics, Telangana, 2015-16

Background characteristic	Prevalence of hypertension ¹	Pre-hypertensive		Hypertensive			Total	Percentage with normal blood pressure and taking medicine to lower blood pressure	Number of men
		Normal SBP <120 mmHg and DBP <80 mmHg	SBP 120-139 mmHg or DBP 80-89 mmHg	Stage 1: SBP 140-159 mmHg or DBP 90-99 mmHg	Stage 2: SBP 160-179 mmHg or DBP 100-109 mmHg	Stage 3: SBP ≥180 mmHg or DBP ≥110 mmHg			
Age									
15-19	2.1	75.8	23.1	1.0	0.0	0.0	100.0	1.1	161
20-24	4.6	56.9	38.9	3.5	0.0	0.0	100.0	0.0	144
25-29	8.4	57.4	34.2	5.2	1.4	1.8	100.0	0.0	166
30-34	25.7	37.8	38.5	17.6	3.2	2.8	100.0	1.5	142
35-39	36.6	28.6	35.3	22.9	7.7	5.5	100.0	0.0	127
40-44	37.7	32.7	32.0	20.4	11.8	3.1	100.0	1.5	107
45-49	35.7	29.6	39.2	20.7	3.8	6.7	100.0	0.8	121
Residence									
Urban	22.3	45.7	33.4	13.5	4.1	3.4	100.0	0.6	441
Rural	17.5	49.0	35.0	10.8	3.0	2.2	100.0	0.7	528
Schooling									
No schooling	22.1	43.1	35.5	14.2	3.3	3.8	100.0	0.4	172
<5 years complete	34.2	40.1	25.7	24.5	5.1	4.6	100.0	0.0	42
5-9 years complete	23.4	49.6	30.1	14.3	3.2	2.9	100.0	2.1	238
10-11 years complete	15.1	47.9	38.2	7.5	3.8	2.7	100.0	0.0	230
12 or more years complete	16.7	49.3	35.0	10.6	3.4	1.7	100.0	0.3	287
Religion									
Hindu	19.6	47.7	34.2	12.1	3.6	2.4	100.0	0.7	839
Muslim	21.0	43.2	36.4	11.6	3.9	4.8	100.0	0.6	102
Christian	(15.4)	(57.8)	(26.8)	(10.3)	(0.0)	(5.1)	100.0	(0.0)	27
Caste/tribe									
Scheduled caste	17.9	50.8	33.0	9.6	3.2	3.4	100.0	1.2	143
Scheduled tribe	10.8	53.7	35.5	9.9	0.8	0.0	100.0	0.0	86
Other backward class	20.0	47.7	33.4	12.4	4.0	2.6	100.0	0.5	614
Other	26.0	38.9	39.0	14.3	3.4	4.5	100.0	1.3	126
Total age 15-49	19.7	47.5	34.2	12.0	3.5	2.7	100.0	0.7	968
Age 50-54	38.7	36.9	32.3	24.8	4.4	1.6	100.0	0.0	68
Total age 15-54	20.9	46.8	34.1	12.9	3.6	2.6	100.0	0.6	1,036

SBP = Systolic blood pressure; DBP = Diastolic blood pressure

¹ A man is classified as having hypertension if he has SBP ≥140 mmHg or DBP ≥90 mmHg at time of survey, or he is currently taking antihypertension medication to control his blood pressure. The term hypertension as used in this table is not meant to be a clinical diagnosis of the disease, but rather to provide an indication of the disease burden in the population at the time of the survey.

() Based on 25-49 unweighted cases

Table 84.1 Random blood glucose levels: Women

Among women age 15-49, percent distribution of random blood glucose values, by background characteristics, Telangana, 2015-16

Background characteristic	Random blood glucose values			Total	Number of women
	≤140 mg/dl (normal)	141-160 mg/dl (high)	>160 mg/dl (very high)		
Age					
15-19	97.6	1.7	0.7	100.0	902
20-24	96.7	2.6	0.7	100.0	1,222
25-29	96.6	1.4	2.0	100.0	1,292
30-34	94.0	2.3	3.6	100.0	948
35-39	90.7	4.1	5.2	100.0	917
40-44	87.0	5.9	7.1	100.0	777
45-49	85.3	3.9	10.8	100.0	807
Residence					
Urban	91.8	3.1	5.1	100.0	3,113
Rural	94.4	2.8	2.8	100.0	3,752
Schooling					
No schooling	90.9	4.0	5.1	100.0	2,310
<5 years complete	92.7	2.4	4.8	100.0	323
5-9 years complete	92.8	2.8	4.3	100.0	1,382
10-11 years complete	95.4	2.2	2.3	100.0	1,290
12 or more years complete	95.2	2.2	2.6	100.0	1,560
Religion					
Hindu	93.6	2.8	3.6	100.0	5,935
Muslim	90.4	4.2	5.5	100.0	686
Christian	91.5	3.0	5.6	100.0	231
Caste/tribe					
Scheduled caste	93.6	2.5	3.9	100.0	1,302
Scheduled tribe	92.2	3.0	4.8	100.0	557
Other backward class	93.2	3.1	3.7	100.0	3,998
Other	93.3	2.8	3.9	100.0	965
Don't know	96.5	3.5	0.0	100.0	44
Total	93.2	3.0	3.9	100.0	6,865

Note: Total includes women belonging to "other" religions, who are not shown separately.

Table 84.2 Random blood glucose levels: Men

Among men age 15-49, percent distribution of random blood glucose values, by background characteristics, Telangana, 2015-16

Background characteristic	Random blood glucose values			Total	Number of men
	≤140 mg/dl (normal)	141-160 mg/dl (high)	>160 mg/dl (very high)		
Age					
15-19	98.9	0.4	0.8	100.0	160
20-24	98.1	1.4	0.5	100.0	140
25-29	98.6	1.2	0.2	100.0	164
30-34	93.3	2.2	4.5	100.0	142
35-39	88.1	2.8	9.2	100.0	127
40-44	90.7	2.1	7.2	100.0	105
45-49	86.4	3.7	9.8	100.0	121
Residence					
Urban	93.2	1.7	5.1	100.0	437
Rural	94.6	2.0	3.3	100.0	523
Schooling					
No schooling	93.6	2.6	3.8	100.0	170
<5 years complete	88.2	7.9	3.9	100.0	42
5-9 years complete	90.1	2.5	7.4	100.0	238
10-11 years complete	95.9	0.7	3.4	100.0	227
12 or more years complete	96.8	1.0	2.2	100.0	282
Religion					
Hindu	94.5	1.7	3.8	100.0	832
Muslim	88.7	3.6	7.7	100.0	101
Christian	(97.4)	(0.0)	(2.6)	100.0	27
Caste/tribe					
Scheduled caste	94.4	1.9	3.7	100.0	141
Scheduled tribe	98.0	1.6	0.4	100.0	84
Other backward class	94.2	2.0	3.8	100.0	611
Other	89.9	1.4	8.7	100.0	123
Total age 15-49	94.0	1.9	4.1	100.0	959
Age 50-54	69.4	9.7	20.9	100.0	67
Total age 15-54	92.4	2.4	5.2	100.0	1,026

() Based on 25-49 unweighted cases

Table 85 Tobacco and alcohol use by women and men

Percentage of women and men age 15-49 by their use of tobacco and alcohol, percent distribution of those who smoke cigarettes or *bidis* by number of cigarettes/*bidis* smoked in the 24 hours preceding the survey, and among those who drink alcohol, the percent distribution of the frequency of alcohol consumption, by residence, Telangana, 2015-16

Tobacco use	Women			Men		
	Urban	Rural	Total	Urban	Rural	Total
Use of tobacco/alcohol						
Smokes cigarettes	0.1	0.1	0.1	16.6	11.4	13.9
Smokes <i>bidis</i>	0.0	0.3	0.1	3.3	10.2	6.9
Smokes cigars or pipe	0.1	0.4	0.2	0.6	1.3	1.0
Smokes <i>hookah</i>	0.0	0.2	0.1	0.0	0.0	0.0
Chews <i>paan</i> with tobacco	0.3	1.9	1.1	1.3	1.4	1.3
Uses other chewing tobacco	0.2	0.8	0.5	0.4	0.6	0.5
Uses <i>gutkha</i> or <i>paan masala</i> with tobacco	0.4	0.8	0.6	5.4	5.4	5.4
Uses <i>khaini</i>	0.1	0.3	0.2	2.9	7.5	5.3
Uses snuff	0.0	0.0	0.0	0.1	0.0	0.0
Uses other tobacco product	0.0	0.1	0.1	0.0	0.1	0.1
Uses any type of tobacco	1.1	4.4	2.8	25.0	31.4	28.3
Drinks alcohol	2.7	14.3	8.8	46.0	61.2	53.9
Number of respondents	3,619	3,948	7,567	506	552	1,058
Among those who smoke cigarettes/<i>bidis</i>, number of cigarettes/<i>bidis</i> smoked in the past 24 hours						
0	*	*	*	0.0	6.7	3.6
1-2	*	*	*	25.9	22.3	24.0
3-5	*	*	*	40.3	15.6	27.0
6-9	*	*	*	10.1	6.7	8.3
10 or more	*	*	*	23.8	47.8	36.7
Don't know	*	*	*	0.0	0.9	0.5
Total	100.0	100.0	100.0	100.0	100.0	100.0
Number of cigarette/ <i>bidi</i> smokers	2	12	15	90	104	194
Among those who drink alcohol, frequency of drinking						
Almost every day	25.4	32.5	31.5	8.0	17.6	13.7
About once a week	28.6	28.5	28.5	43.3	47.0	45.5
Less than once a week	46.0	38.9	40.0	48.7	35.4	40.8
Total	100.0	100.0	100.0	100.0	100.0	100.0
Number who drink alcohol	97	566	663	233	338	571

* Percentage not shown; based on fewer than 25 unweighted cases

Table 86 Methods of menstrual protection

Percentage of women age 15-24 who use various methods of menstrual protection and percentage using a hygienic method, according to background characteristics, Telangana, 2015-16

Background characteristic	Type of menstrual protection ¹						Percentage using a hygienic method ²	Number of women
	Cloth	Locally prepared napkins	Sanitary napkins	Tampons	Other	Nothing		
Age								
15-19	27.8	20.9	63.0	2.9	0.1	0.3	81.2	997
20-24	37.2	15.6	58.0	2.9	0.0	0.2	72.6	1,355
Residence								
Urban	22.9	17.7	71.5	2.1	0.0	0.1	86.7	1,096
Rural	42.2	17.9	50.3	3.5	0.1	0.4	67.2	1,256
Schooling								
No schooling	74.1	2.7	29.0	0.0	0.0	1.2	31.4	176
<5 years complete	77.3	7.8	28.7	0.0	0.0	0.0	35.0	51
5-9 years complete	38.2	16.7	56.4	2.9	0.1	0.2	72.3	504
10-11 years complete	31.1	17.2	60.4	2.5	0.0	0.4	77.1	675
12 or more years complete	22.0	22.3	69.5	3.8	0.0	0.1	88.3	946
Religion								
Hindu	33.5	17.5	59.1	3.0	0.0	0.3	75.5	2,002
Muslim	32.5	20.7	62.6	1.5	0.0	0.0	79.1	278
Christian	24.7	12.9	83.1	2.6	0.0	1.4	86.4	63
Caste/tribe								
Scheduled caste	32.9	18.6	58.4	4.3	0.1	0.5	76.0	450
Scheduled tribe	47.1	12.2	55.0	0.7	0.0	0.4	65.1	205
Other backward class	33.3	18.7	58.7	2.8	0.0	0.2	75.7	1,384
Other	24.1	16.1	73.0	2.5	0.0	0.0	86.8	307
Total	33.2	17.8	60.1	2.9	0.0	0.3	76.2	2,352

Note: Table is based on women age 15-24 who have ever menstruated. Total includes women belonging to "other" religions and women who don't know their caste/tribe, who are not shown separately.

¹ Respondents may report multiple methods so the sum may exceed 100 percent

² Locally prepared napkins, sanitary napkins, and tampons are considered to be hygienic methods of protection

Table 87 Employment and cash earnings of women and men

Percentage of women and men age 15-49 who were employed at any time in the 12 months preceding the survey and percent distribution of women and men employed in the 12 months preceding the survey by type of earnings and sector of employment, according to age, Telangana, 2015-16

Age	Percentage employed	Number of respondents	Percent distribution of employed respondents by type of earnings				Percent distribution of employed respondents by sector				Number of employed respondents
			Cash and in-kind		In-kind only		Total	Agriculture	Non-agriculture	Total	
			Cash only	and in-kind	only	Not paid					
WOMEN											
15-19	28.1	183	84.5	1.3	4.4	9.8	100.0	56.6	43.4	100.0	51
20-24	30.8	233	95.6	3.1	1.4	0.0	100.0	51.9	48.1	100.0	72
25-29	49.1	231	97.3	0.0	0.5	2.2	100.0	54.5	45.5	100.0	113
30-34	52.8	162	93.5	2.8	0.8	2.9	100.0	59.0	41.0	100.0	86
35-39	62.9	170	91.7	2.0	2.0	4.3	100.0	55.2	44.8	100.0	107
40-44	62.4	151	92.9	1.6	1.1	4.4	100.0	42.6	57.4	100.0	94
45-49	55.8	155	94.9	2.5	0.0	2.6	100.0	62.7	37.3	100.0	87
Total	47.4	1,285	93.5	1.8	1.3	3.4	100.0	54.4	45.6	100.0	610
MEN											
15-19	30.1	174	85.9	3.0	0.0	11.2	100.0	35.2	64.8	100.0	52
20-24	70.9	155	94.7	2.4	0.0	2.9	100.0	18.4	81.6	100.0	110
25-29	91.8	184	93.4	3.7	0.8	2.2	100.0	27.2	72.8	100.0	169
30-34	94.9	155	96.5	3.5	0.0	0.0	100.0	31.1	68.9	100.0	147
35-39	98.7	140	91.4	4.8	0.8	3.0	100.0	31.1	68.9	100.0	139
40-44	97.9	116	92.4	4.3	0.0	3.3	100.0	41.0	59.0	100.0	114
45-49	98.0	132	86.6	7.0	0.6	5.9	100.0	42.1	57.9	100.0	130
Total	81.4	1,058	92.2	4.2	0.4	3.3	100.0	31.9	68.1	100.0	861

Table 88 Control over and magnitude of women's cash earnings

Percentage of currently married women and men age 15-49 by who decide how women's and men's cash earnings are used and by the magnitude of women's cash earnings compared with their husbands' cash earnings, according to background characteristics, Telangana, 2015-16

Background characteristic	Percentage of currently married women who report that they:					Percentage of currently married men who report that:						
	Alone or jointly with their husband decide how their own earnings are used	Number employed for cash	Alone or jointly with their husband decide how their husbands' earnings are used	Number of women whose husbands are employed for cash	Earn more or about the same as their husband	Number employed for cash whose husbands are employed for cash	They alone or jointly with their wife decide how their wife's earnings are used	Number of men with wives who are employed for cash	Their wife alone or jointly with them decides how their own earnings are used	Number employed for cash	Their wife earns more or about the same as them	Number employed for cash and whose wives are employed for cash
Age												
15-19	*	9	(34.7)	23	*	8	nc	0	nc	0	nc	0
20-29	68.9	140	66.8	261	43.2	108	(93.2)	20	63.4	146	*	18
30-39	76.6	155	71.8	201	43.4	115	86.2	71	69.9	259	54.8	66
40-49	79.6	141	79.7	147	34.8	85	82.7	73	70.6	225	53.1	65
Residence												
Urban	81.3	123	70.8	275	33.2	83	(85.3)	54	73.8	291	(47.4)	47
Rural	71.5	322	69.8	357	43.4	234	85.7	110	64.2	338	53.5	101
Schooling												
No schooling	74.2	239	72.3	256	44.3	174	89.1	60	70.5	148	65.5	56
<5 years complete	(76.1)	31	(67.8)	28	*	19	*	8	(63.9)	37	*	7
5-9 years complete	70.8	76	70.5	129	26.6	59	(83.5)	41	74.9	151	(36.2)	40
10-11 years complete	73.9	54	70.2	116	(48.3)	38	(82.9)	29	67.1	134	*	23
12 or more years complete	(79.2)	45	65.4	104	(36.9)	27	(91.3)	25	63.4	160	*	22
Religion												
Hindu	73.4	411	70.2	571	41.8	291	87.4	143	67.9	553	50.1	132
Muslim	(80.7)	22	71.8	46	*	16	*	10	70.7	58	*	10
Caste/tribe												
Scheduled caste	75.2	80	65.2	93	31.4	54	(78.0)	39	70.8	93	(69.1)	34
Scheduled tribe	65.7	58	72.4	61	40.2	48	*	17	63.5	59	*	17
Other backward class	74.5	259	70.0	396	44.1	181	83.8	86	68.4	393	37.9	77
Other	(79.2)	42	74.8	79	(41.1)	30	*	21	70.9	85	*	20
Total	74.2	445	70.2	632	40.7	316	85.5	163	68.6	629	51.5	148

Note: Total includes Christian women/men and women/men who don't know their caste/tribe, who are not shown separately.

nc = No cases

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

Table 89 Decision making

Percent distribution of currently married women and men age 15-49 by who usually make decisions about specific issues, by residence, Telangana, 2015-16

Decision	According to women, person who usually makes the decision					According to men, person who usually makes the decision						
	Mainly respondent	Respondent and husband jointly	Mainly husband	Someone else	Other	Total	Mainly respondent	Respondent and wife jointly	Mainly wife	Someone else	Other	Total
	URBAN											
Own health care	12.3	57.5	29.6	0.4	0.2	100.0	29.0	62.0	9.1	0.0	0.0	100.0
Major household purchases	15.3	59.2	24.8	0.5	0.2	100.0	27.7	67.6	4.1	0.6	0.0	100.0
Visits to her family or relatives	5.6	64.6	29.0	0.6	0.2	100.0	na	na	na	na	na	na
	RURAL											
Own health care	10.0	57.0	31.7	0.8	0.5	100.0	35.4	50.5	12.4	1.7	0.0	100.0
Major household purchases	15.0	55.6	26.7	2.2	0.4	100.0	30.0	60.1	9.1	0.5	0.3	100.0
Visits to her family or relatives	8.2	62.7	27.6	1.2	0.3	100.0	na	na	na	na	na	na
	TOTAL											
Own health care	11.0	57.2	30.7	0.6	0.4	100.0	32.5	55.7	10.9	0.9	0.0	100.0
Major household purchases	15.1	57.2	25.8	1.5	0.3	100.0	29.0	63.5	6.8	0.5	0.1	100.0
Visits to her family or relatives	7.0	63.5	28.3	0.9	0.2	100.0	na	na	na	na	na	na

na = Not applicable

Table 90 Decision making by background characteristics

Percentage of currently married women and men age 15-49 who usually make specific decisions either by themselves or jointly with their spouse, by background characteristics, Telangana, 2015-16

Background characteristic	Percentage of women who usually make specific decisions alone or jointly with their husband			Percentage of men who usually make specific decisions alone or jointly with their wife			Number of women	Percentage who participate in three decisions	Percentage who participate in none of the three decisions	Percentage who participate in both decisions	Percentage who participate in neither decision	Number of men
	Own health care	Making major household purchases	Visits to her family or relatives	Own health care	Making major household purchases	Percentage who participate in three decisions						
Age												
15-19	(50.8)	(46.3)	(48.7)	(25.8)	(34.8)	33	nc	nc	nc	nc	0	
20-24	66.2	64.1	68.7	55.5	23.7	148	(90.5)	(97.8)	(88.3)	(0.0)	32	
25-29	59.0	72.1	70.0	51.4	21.3	206	87.6	92.3	83.7	3.8	123	
30-39	71.4	74.0	72.0	56.6	14.8	293	88.8	92.9	84.5	2.8	268	
40-49	75.5	78.8	73.3	67.2	16.8	259	87.6	91.3	83.3	4.3	242	
Residence												
Urban	69.8	74.5	70.2	59.3	17.9	427	90.9	95.3	88.5	2.3	302	
Rural	67.0	70.6	70.9	55.4	19.7	511	85.9	90.1	80.4	4.3	363	
Schooling												
No schooling	70.8	73.1	71.9	59.6	17.2	372	86.1	90.3	81.6	5.2	158	
<5 years complete	59.6	76.3	62.2	52.4	21.0	46	(88.1)	(84.9)	(78.7)	(5.8)	40	
5-9 years complete	66.4	70.1	67.7	56.5	23.6	185	83.7	93.8	79.5	1.9	155	
10-11 years complete	68.1	71.4	70.4	52.9	19.2	162	90.3	93.0	86.6	3.2	145	
12 or more years complete	67.3	73.2	73.1	57.9	16.7	173	92.6	94.6	89.8	2.7	167	
Employment (past 12 months)												
Employed	71.1	75.9	74.4	60.4	15.8	467	88.3	92.5	84.1	3.3	653	
Employed, for cash	71.7	76.7	74.9	61.2	15.2	445	88.8	92.8	84.5	3.0	629	
Employed, not for cash	(57.4)	(58.0)	(64.0)	(44.1)	(28.8)	21	(77.3)	(85.6)	(74.2)	(11.4)	23	
Not employed	65.5	68.9	66.7	53.9	21.9	471	*	*	*	*	13	

Continued...

Table 90 Decision making by background characteristics – Continued

Percentage of currently married women and men age 15-49 who usually make specific decisions either by themselves or jointly with their spouse, by background characteristics, Telangana, 2015-16

Background characteristic	Percentage of women who usually make specific decisions alone or jointly with their husband			Percentage of men who usually make specific decisions alone or jointly with their wife			Number of women	Percentage who participate in three decisions	Percentage who participate in none of the three decisions	Percentage who participate in both decisions	Percentage who participate in neither decision	Number of men
	Own health care	Making major household purchases	Visits to her family or relatives	Own health care	Making major household purchases	Percentage who participate in three decisions						
Number of living children												
0	57.9	61.4	61.4	45.4	27.4	79	93.6	94.5	90.3	2.2	93	
1-2	67.7	72.7	71.7	57.5	19.0	598	87.1	92.8	83.0	3.1	409	
3-4	72.8	75.1	70.5	58.9	15.4	243	89.2	89.9	84.2	5.0	153	
Household structure¹												
Nuclear	71.6	77.2	74.5	61.8	15.5	587	88.6	93.5	85.0	2.9	413	
Non-nuclear	62.6	64.4	64.0	49.4	24.5	351	87.5	90.8	82.6	4.2	252	
Religion												
Hindu	68.6	72.0	70.9	57.3	18.8	831	87.5	92.4	83.6	3.7	585	
Muslim	64.9	74.7	66.5	54.0	18.9	84	96.3	96.3	92.6	0.0	58	
Caste/tribe												
Scheduled caste	65.0	73.6	72.7	55.2	19.4	134	86.2	91.8	82.0	3.9	100	
Scheduled tribe	65.4	67.9	65.3	51.0	21.6	76	83.6	93.8	80.1	2.8	60	
Other backward class	68.2	71.4	70.3	57.5	19.5	591	88.5	92.6	84.5	3.4	419	
Other	72.5	77.3	73.5	61.2	14.8	130	92.0	91.9	87.0	3.1	87	
Total	68.3	72.4	70.6	57.1	18.9	938	88.2	92.5	84.1	3.4	665	

Note: Total includes women/men who have 5 or more living children, women/men belonging to Christian or "other" religions, and women/men who don't know their caste/tribe, who are not shown separately. nc = No cases

¹ Nuclear households are households comprised of a married couple or a man or a woman living alone or with unmarried children (biological, adopted, or fostered) with or without unrelated individuals. () Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

Table 91 Women's access to money and credit

Percentage of women age 15-49 who have access to money, who know of a microcredit programme, who have taken a loan from a microcredit programme, and who are allowed to go to three specified places alone by background characteristics, Telangana, 2015-16

Background characteristic	Women's access to money		Women's knowledge and use of microcredit programmes			Number of women
	Percentage who have money that they can decide how to use	Percentage who have a bank or savings account that they themselves use	Percentage who know of a microcredit programme	Percentage who have taken a loan from a microcredit programme	Percentage of women allowed to go to three specified places alone ¹	
Age						
15-19	7.6	41.4	37.8	0.8	19.4	183
20-24	17.5	56.2	58.5	9.4	30.2	233
25-29	25.3	58.5	60.6	24.0	45.4	231
30-39	29.4	67.4	69.2	33.4	55.0	332
40-49	31.5	65.7	65.7	34.4	58.9	306
Residence						
Urban	29.7	60.7	57.0	13.5	42.0	614
Rural	18.6	58.7	63.6	31.6	47.0	671
Schooling						
No schooling	24.3	59.5	61.5	34.4	56.0	440
<5 years complete	22.3	63.1	56.3	30.5	48.5	58
5-9 years complete	18.8	59.2	59.8	25.5	42.5	255
10-11 years complete	24.8	53.5	58.7	17.0	35.2	233
12 or more years complete	27.3	64.5	61.6	7.2	36.2	299
Employment (past 12 months)						
Employed	29.8	67.6	67.5	34.8	55.2	610
Employed, for cash	30.9	68.1	68.0	35.4	56.5	581
Employed, not for cash	(7.1)	(56.7)	(58.5)	(23.8)	(29.9)	29
Not employed	18.5	52.5	54.0	12.3	35.0	675
Number of living children						
0	15.2	50.9	49.3	4.4	25.2	342
1-2	27.6	62.7	66.4	28.1	49.5	658
3-4	24.5	62.8	61.0	34.9	56.7	264

Continued...

Table 91 Women's access to money and credit—Continued

Percentage of women age 15-49 who have access to money, who know of a microcredit programme, who have taken a loan from a microcredit programme, and who are allowed to go to three specified places alone by background characteristics, Telangana, 2015-16

Background characteristic	Women's access to money		Women's knowledge and use of microcredit programmes			Number of women
	Percentage who have money that they can decide how to use	Percentage who have a bank or savings account that they themselves use	Percentage who know of a microcredit programme	Percentage who have taken a loan from a microcredit programme	Percentage of women allowed to go to three specified places alone ¹	
Household structure²						
Nuclear	23.7	61.6	62.9	24.6	45.5	806
Non-nuclear	24.2	56.4	56.4	20.3	43.1	479
Religion						
Hindu	23.8	59.3	61.3	24.5	45.6	1,133
Muslim	24.9	64.6	52.0	10.3	36.4	121
Christian	(23.0)	(60.2)	(72.8)	(19.6)	(40.5)	27
Caste/tribe						
Scheduled caste	24.6	60.3	63.1	31.0	44.5	198
Scheduled tribe	21.4	56.5	60.6	22.9	52.8	108
Other backward class	23.2	59.7	59.5	22.3	44.1	796
Other	28.9	60.6	61.4	16.4	41.8	175
Total	23.9	59.7	60.4	23.0	44.6	1,285

Note: Total includes women who have 5 or more living children, women belonging to "other" religions, and women who don't know their caste/tribe, who are not shown separately.

¹ To the market, to the health facility, and to places outside the village/community

² Nuclear households are households comprised of a married couple or a man or a woman living alone or with unmarried children (biological, adopted, or fostered) with or without unrelated individuals.

() Based on 25-49 unweighted cases

Table 92 Ownership of assets

Percentage of women and men age 15-49 who own a house or land either alone or jointly, percentage of women who own a mobile phone that they themselves use, and among women who own a mobile phone, percentage who can read SMS messages, according to background characteristics, Telangana, 2015-16

Background characteristic	Women				Among women who have a mobile phone, percentage who can read SMS messages	Men			
	Own a house alone or jointly	Own land alone or jointly	Have a mobile phone that they themselves use	Number of women		Number of women	Own a house alone or jointly	Own land alone or jointly	Number of men
Age									
15-19	33.4	20.8	26.1	183	(91.4)	48	60.9	40.0	174
20-24	45.2	31.1	51.7	233	87.6	121	59.3	39.7	155
25-29	47.4	30.5	60.4	231	71.5	139	69.9	47.9	184
30-34	41.8	29.0	57.8	162	60.7	94	77.4	54.1	155
35-39	51.1	35.0	54.7	170	46.1	93	76.6	51.3	140
40-44	49.1	31.8	46.3	151	51.5	70	78.2	57.7	116
45-49	62.3	37.8	32.2	155	(36.0)	50	79.4	53.1	132
Residence									
Urban	38.9	20.3	63.1	614	74.3	388	60.1	32.7	506
Rural	54.0	40.2	33.8	671	50.7	227	80.9	62.9	552
Religion									
Hindu	47.1	32.1	47.2	1,133	65.3	534	72.7	51.1	910
Muslim	45.0	16.8	57.0	121	68.2	69	61.0	31.2	120
Christian	(43.8)	(34.6)	(40.0)	27	*	11	(57.2)	(37.8)	27
Caste/tribe									
Scheduled caste	49.3	28.5	38.3	198	47.0	76	78.3	51.2	160
Scheduled tribe	38.4	33.5	25.6	108	(55.7)	28	80.9	67.3	87
Other backward class	49.3	31.9	47.8	796	65.2	381	70.6	45.6	666
Other	36.8	26.8	73.0	175	81.1	128	58.6	47.3	145
Schooling									
No schooling	50.9	35.7	28.3	440	1.6	124	84.9	58.5	179
<5 years complete	46.7	40.2	46.1	58	(12.1)	27	81.9	59.4	44
5-9 years complete	51.6	27.9	48.6	255	56.9	124	66.5	45.3	257
10-11 years complete	42.6	28.2	56.4	233	91.4	131	71.7	50.6	245
12 or more years complete	39.9	25.8	69.4	299	99.6	208	64.9	42.5	333
Total age 15-49	46.8	30.7	47.8	1,285	65.6	614	70.9	48.5	1,058
Age 50-54	na	na	na	na	na	na	85.3	53.4	75
Total age 15-54	na	na	na	na	na	na	71.9	48.8	1,133

Note: Total includes women/men belonging to "other" religions and women/men who don't know their caste/tribe, who are not shown separately.

na = Not applicable

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

Table 93 Gender role attitudes

Percentage of women and men age 15-49 with specific attitudes toward wife beating and refusal by a wife to have sex with her husband by reason, and percentage of men age 15-49 who agree that a man can behave in specific ways if his wife refuses to have sex with him, according to marital status, Telangana, 2015-16

Reason/behaviour	Ever-married		Never married		Total	
	Women	Men	Women	Men	Women	Men
Percentage who agree that a husband is justified in hitting or beating his wife if:						
She goes out without telling him	43.9	30.9	39.9	37.4	43.1	33.2
She neglects the house or children	71.2	55.1	64.6	59.8	69.9	56.8
She argues with him	47.9	39.8	38.5	42.3	46.0	40.7
She refuses to have sexual intercourse with him	20.3	19.1	16.4	18.7	19.5	19.0
She doesn't cook properly	25.2	17.8	22.6	16.9	24.7	17.5
He suspects her of being unfaithful	34.0	47.4	27.0	51.0	32.6	48.7
She shows disrespect for in-laws	67.3	49.6	59.0	54.5	65.7	51.4
Percentage who agree with at least one specified reason	85.5	73.7	75.8	77.8	83.6	75.2
Percentage who agree that a wife is justified in refusing to have sex with her husband when she:						
Knows her husband has a sexually transmitted disease	78.5	84.0	60.6	79.7	75.0	82.4
Knows her husband has sex with other women	77.8	78.0	60.5	75.5	74.4	77.1
Is tired or not in the mood	74.9	67.9	53.2	61.0	70.6	65.5
Percentage who agree with all three reasons	65.8	59.8	46.5	54.4	62.0	57.9
Percentage who agree with none of the three reasons	13.3	10.2	32.6	13.7	17.1	11.5
Percentage who agree that when a wife refuses to have sex with her husband, he has the right to:						
Get angry and reprimand her	na	43.9	na	40.2	na	42.6
Refuse to give her financial support	na	31.3	na	29.6	na	30.7
Use force to have sex even if she doesn't want to	na	26.1	na	24.6	na	25.6
Have sex with another woman	na	26.3	na	24.7	na	25.8
Percentage who agree with all four behaviours	na	13.8	na	13.2	na	13.6
Percentage who agree with none of the four behaviours	na	41.7	na	46.5	na	43.4
Number of respondents	1,031	678	255	380	1,285	1,058
na = Not applicable						

Table 94 Gender role attitudes by background characteristics

Percentage of women and men age 15-49 who agree that a husband is justified in hitting or beating his wife for at least one specified reason, who agree that a wife is justified in refusing to have sex with her husband for all specified reasons, and percentage of men who agree that when a wife refuses to have sex with her husband, the husband does not have the right to any of the four specified behaviours, by background characteristics, Telangana, 2015-16

Background characteristic	Percentage who agree that a husband is justified in hitting or beating his wife for at least one specified reason ¹		Percentage who agree that a wife is justified in refusing to have sex with her husband for all specified reasons ²		Percentage who agree that when a wife refuses to have sex with her husband, he does not have the right to any of the four specified behaviours ³	Number	
	Women	Men	Women	Men	Men	Women	Men
Age							
15-19	78.9	77.0	41.0	55.2	32.6	183	174
20-24	82.7	79.4	65.2	48.3	42.4	233	155
25-29	83.4	77.0	72.5	58.3	35.3	231	184
30-39	84.4	76.1	64.5	63.2	39.9	332	296
40-49	86.3	68.8	61.3	58.9	43.9	306	248
Residence							
Urban	82.1	73.1	63.1	58.2	43.0	614	506
Rural	84.9	77.1	60.9	57.5	35.7	671	552
Schooling							
No schooling	83.7	71.8	59.2	58.8	37.9	440	179
<5 years complete	94.8	66.5	65.7	68.0	41.2	58	44
5-9 years complete	86.9	80.1	59.5	51.7	42.8	255	257
10-11 years complete	83.7	75.0	63.0	57.4	37.6	233	245
12 or more years complete	78.3	74.5	66.7	61.1	38.0	299	333
Employment (past 12 months)							
Employed	84.1	76.6	60.7	60.0	39.8	610	861
Employed, for cash	83.6	76.9	60.7	60.6	40.6	581	829
Employed, not for cash	(93.8)	(68.3)	(60.5)	(42.6)	(19.5)	29	31
Not employed	83.1	68.9	63.1	48.5	36.5	675	197
Marital status							
Never married	75.8	77.8	46.5	54.4	38.4	255	380
Currently married	85.2	73.4	65.6	59.6	40.1	938	665
Widowed/divorced/ separated/deserted	88.8	*	67.9	*	*	92	13

Continued...

Table 94 Gender role attitudes by background characteristics—Continued

Percentage of women and men age 15-49 who agree that a husband is justified in hitting or beating his wife for at least one specified reason, who agree that a wife is justified in refusing to have sex with her husband for all specified reasons, and percentage of men who agree that when a wife refuses to have sex with her husband, the husband does not have the right to any of the four specified behaviours, by background characteristics, Telangana, 2015-16

Background characteristic	Percentage who agree that a husband is justified in hitting or beating his wife for at least one specified reason ¹		Percentage who agree that a wife is justified in refusing to have sex with her husband for all specified reasons ²		Percentage who agree that when a wife refuses to have sex with her husband, he does not have the right to any of the four specified behaviours ³	Number	
	Women	Men	Women	Men	Men	Women	Men
Number of living children							
0	78.9	77.0	50.2	56.9	39.4	342	473
1-2	85.7	73.0	67.9	61.3	37.3	658	419
3-4	86.0	75.2	61.6	52.6	43.8	264	155
Household structure⁴							
Nuclear	81.7	73.2	62.5	57.0	39.0	806	671
Non-nuclear	86.7	78.6	61.1	59.3	39.6	479	386
Religion							
Hindu	83.3	74.9	61.9	58.7	36.9	1,133	910
Muslim	84.6	78.4	61.1	54.5	53.9	121	120
Christian	(95.2)	(69.5)	(66.6)	(43.9)	(49.2)	27	27
Caste/tribe							
Scheduled caste	87.0	70.8	67.0	59.4	41.3	198	160
Scheduled tribe	86.2	74.9	53.4	38.9	36.9	108	87
Other backward class	81.7	75.3	61.4	58.5	38.1	796	666
Other	86.1	79.8	65.4	64.7	43.1	175	145
Total	83.6	75.2	62.0	57.9	39.2	1,285	1,058

Note: Total includes women/men who have 5 or more living children, women/men belonging to "other" religions, and women/men who don't know their caste/tribe, who are not shown separately.

¹ Specified reasons are: she goes out without telling him, she neglects the house or children, she argues with him, she refuses to have sexual intercourse with him, she doesn't cook properly, he suspects she is unfaithful, and she shows disrespect for in-laws.

² Specified reasons are: knows husband has a sexually transmitted disease, knows husband has sex with other women, and is tired or not in the mood.

³ Specified behaviours are: gets angry and reprimands her, refuses to give her financial support, uses force to have sex, and has sex with another woman.

⁴ Nuclear households are households comprised of a married couple or a man or a woman living alone or with unmarried children (biological, adopted, or fostered) with or without unrelated individuals.

() Based on 25-49 unweighted cases

* Percentage not shown; based on fewer than 25 unweighted cases

Table 95 Experience of physical and sexual violence

Percentage of women age 15-49 who have ever experienced physical or sexual violence; among those who have experienced physical violence, the person committing the violence; and among those who have experienced sexual violence, the person committing the violence the first time, by marital status, Telangana, 2015-16

Type of violence/perpetrator	Ever-married	Never married	Total
Type of violence experienced			
Physical violence only	42.7	22.9	38.9
Sexual violence only	1.3	0.0	1.1
Physical and sexual violence	6.1	0.0	4.9
Physical or sexual violence	50.1	22.9	44.9
Number of women	726	169	895
Person committing physical violence¹			
Current husband	85.6	*	77.1
Former husband	0.2	*	0.2
Father/step-father	11.7	*	14.5
Mother/step-mother	24.7	*	29.4
Sister/brother	7.0	*	8.6
Other relative	0.6	*	0.5
Former boyfriend	0.2	*	0.2
Mother in-law	1.0	*	0.9
Father in-law	0.2	*	0.2
Other in-law	0.7	*	0.7
Teacher	4.0	*	5.6
Other	0.3	*	0.2
Number who experienced physical violence since age 15	354	39	393
Person committing sexual violence the first time			
Current husband	88.2	nc	88.2
Former husband	1.1	nc	1.1
Father/step-father	4.4	nc	4.4
Brother/step-brother	1.8	nc	1.8
Teacher	2.4	nc	2.4
Stranger	1.2	nc	1.2
Other	0.9	nc	0.9
Number who experienced sexual violence	54	0	54

Note: All women were asked about their experience of physical violence since age 15. Ever-married women were also asked about their experience of spousal physical violence at any age.

nc = No cases

¹ Women can report more than one person who committed the violence

* Percentage not shown; based on fewer than 25 unweighted cases

Table 96 Experience of violence during pregnancy

Among women age 15-49 who have ever been pregnant, percentage who have ever experienced physical violence during pregnancy, by background characteristics, Telangana, 2015-16

Background characteristic	Percentage who experienced violence during pregnancy	Number of women who have ever been pregnant
Age		
20-24	2.2	93
25-29	6.5	145
30-39	4.5	223
40-49	7.3	203
Residence		
Urban	4.4	291
Rural	7.0	389
Marital status		
Currently married	4.3	621
Widowed/divorced/separated/deserted	22.5	60
Number of living children		
1-2	5.3	476
3-4	7.3	169
Schooling		
No schooling	8.6	293
<5 years complete	(8.6)	34
5-9 years complete	8.2	120
10-11 years complete	1.3	113
12 or more years complete	0.6	120
Religion		
Hindu	6.2	609
Muslim	3.5	52
Caste/tribe		
Scheduled caste	14.2	112
Scheduled tribe	7.1	55
Other backward class	4.4	416
Other	0.9	92
Total	5.9	681

Note: Total includes women age 15-19, women with no living children, women with 5 or more living children, women belonging to Christian or "other" religions, and women who don't know their caste/tribe, who are not shown separately.

() Based on 25-49 unweighted cases

Table 97 Forms of spousal violence

Percentage of ever-married women age 15-49 who have experienced various forms of violence committed by their husband ever and in the 12 months preceding the survey, Telangana, 2015-16

Type of violence	Ever ¹	In the past 12 months		
		Often	Sometimes	Often or sometimes
Physical violence				
Any form of physical violence	41.8	5.6	28.6	34.3
Pushed her, shook her, or threw something at her	21.4	2.4	14.5	16.9
Twisted her arm or pulled her hair	13.8	1.3	9.0	10.3
Slapped her	35.7	2.5	25.8	28.3
Punched her with his fist or with something that could hurt her	11.3	1.9	7.0	8.9
Kicked her, dragged her, or beat her up	16.2	2.0	10.1	12.1
Tried to choke her or burn her on purpose	1.4	0.5	0.7	1.2
Threatened her or attacked her with a knife, gun, or any other weapon	2.1	0.4	0.9	1.3
Sexual violence				
Any form of sexual violence	6.5	1.0	4.1	5.1
Physically forced her to have sexual intercourse with him even when she did not want to	5.2	0.5	3.3	3.8
Physically forced her to perform any sexual acts she did not want to	2.0	0.4	1.2	1.5
Forced her with threats or in any other way to perform sexual acts she did not want to	2.7	0.2	1.7	1.9
Emotional violence				
Any form of emotional violence	20.0	2.8	13.5	16.3
Said or did something to humiliate her in front of others	10.6	1.0	7.2	8.2
Threatened to hurt or harm her or someone close to her	10.8	1.8	6.7	8.4
Insulted her or made her feel bad about herself	12.1	1.6	8.2	9.8
Any form of physical and/or sexual violence	43.0	6.0	29.2	35.2
Any form of physical and sexual violence	5.3	0.6	3.5	4.1
Any form of physical and/or sexual and/or emotional violence	46.0	6.8	31.7	38.5
Any form of physical and sexual and emotional violence	3.8	0.5	2.7	3.2
Any violence by women against their husband ²	4.7	0.9	3.1	4.0
Number of ever-married women	726	726	726	726

Note: Husband refers to the current husband for currently married women and the most recent husband for widowed, divorced, separated, or deserted women.

¹ Includes in the past 12 months

² Any violence by women against their husband when he was not already beating or physically hurting her

Table 98 Spousal violence by background characteristics

Percentage of ever-married women age 15-49 who have ever experienced emotional, physical, or sexual violence committed by their husband, according to background characteristics, Telangana, 2015-16

Background characteristic	Emotional violence	Physical violence	Sexual violence	Physical or sexual violence	Emotional, physical, or sexual violence	Number of women
Age						
15-19	(28.6)	(40.7)	(2.9)	(42.2)	(42.2)	29
20-24	14.9	26.3	4.8	28.2	29.9	103
25-29	19.3	45.0	7.4	45.8	48.2	154
30-39	21.9	42.2	6.4	43.6	46.8	231
40-49	19.9	47.0	7.1	47.6	52.0	208
Residence						
Urban	10.9	36.9	3.1	36.9	39.8	313
Rural	27.0	45.6	9.0	47.6	50.7	413
Schooling						
No schooling	24.6	51.4	9.8	52.2	54.2	301
<5 years complete	(38.8)	(58.8)	(15.2)	(66.0)	(66.0)	36
5-9 years complete	21.2	40.4	6.4	42.2	45.1	135
10-11 years complete	15.1	30.7	1.9	31.3	40.0	122
12 or more years complete	7.9	27.3	0.9	27.3	28.4	132
Employment (past 12 months)						
Not employed	9.8	32.1	3.4	32.7	34.5	345
Employed for cash	29.7	51.0	8.9	52.7	56.7	368
Marital status						
Currently married	19.0	40.8	5.7	42.0	45.3	665
Widowed	20.7	45.7	10.9	47.1	47.1	47
Marital duration¹						
Married only once	18.9	40.6	5.6	41.7	45.0	660
0-4 years	12.5	24.6	2.3	25.3	27.1	122
5-9 years	18.0	41.2	6.1	43.5	45.5	119
10+ years	21.0	45.0	6.4	45.9	50.1	420
Number of living children						
0	15.1	28.3	4.0	30.0	30.8	68
1-2	19.8	42.4	6.1	43.6	46.7	476
3-4	23.2	46.6	8.4	47.5	51.5	169
Household structure²						
Nuclear	19.5	43.5	6.6	44.7	48.5	442
Non-nuclear	20.9	39.3	6.2	40.2	42.1	283
Religion						
Hindu	20.7	42.7	6.5	44.0	47.3	652
Muslim	6.7	25.1	6.9	25.1	25.1	53

Continued...

Table 98 Spousal violence by background characteristics—Continued

Percentage of ever-married women age 15-49 who have ever experienced emotional, physical, or sexual violence committed by their husband, according to background characteristics, Telangana, 2015-16

Background characteristic	Emotional violence	Physical violence	Sexual violence	Physical or sexual violence	Emotional, physical, or sexual violence	Number of women
Caste/tribe						
Scheduled caste	27.8	56.7	7.3	58.8	60.8	125
Scheduled tribe	37.4	56.3	13.4	58.0	61.6	58
Other backward class	16.6	38.0	5.5	39.0	42.0	442
Other	15.0	32.2	5.5	32.2	36.7	96
Respondent's father beat her mother						
Yes	23.5	55.2	5.8	56.2	58.9	294
No	18.0	31.9	7.2	33.2	36.3	416
Total	20.0	41.8	6.5	43.0	46.0	726

Note: Husband refers to the current husband for currently married women and the most recent husband for widowed, divorced, separated, or deserted women. Total includes women employed but not for cash, currently married women who have been married more than once, women with 5 or more living children, Christian women, women who don't know their caste/tribe, and women who don't know whether their father beat their mother, who are not shown separately.

¹ Currently married women only

² Nuclear households are households comprised of a married couple or a man or a woman living alone or with unmarried children (biological, adopted, or fostered) with or without unrelated individuals.

() Based on 25-49 unweighted cases

Table 99 Spousal violence by husband's characteristics and empowerment indicators

Percentage of ever-married women age 15-49 who have ever suffered emotional, physical, or sexual violence committed by their husband, according to his characteristics, marital characteristics, and selected empowerment indicators, Telangana, 2015-16

Background characteristic	Emotional violence	Physical violence	Sexual violence	Physical or sexual violence	Emotional, physical, or sexual violence	Number of women
Husband's schooling						
No schooling	26.7	51.9	9.6	53.0	55.1	215
<5 years complete	8.7	46.2	7.5	48.1	48.1	47
5-9 years complete	22.6	46.0	6.9	47.5	50.4	141
10-11 years complete	21.4	29.6	6.1	31.0	37.8	124
12 or more years complete	12.3	34.4	2.5	35.0	37.5	197
Husband's alcohol consumption						
Does not drink	7.9	28.2	3.7	29.1	31.3	310
Gets drunk sometimes	24.2	48.6	5.8	49.7	52.1	314
Gets drunk often	48.2	67.1	18.2	69.5	77.2	93
Spousal age difference¹						
Wife 1-4 years younger	19.0	34.9	5.0	35.4	39.5	230
Wife 5-9 years younger	16.9	42.0	5.6	43.3	46.3	322
Wife 10+ years younger	28.1	48.6	7.7	50.3	53.9	85
Spousal schooling difference						
Husband has more schooling	21.5	45.0	7.3	46.3	49.1	312
Wife has more schooling	17.4	37.2	3.3	38.9	39.8	145
Both have equal schooling	11.2	23.8	2.2	25.2	33.3	91
Neither attended school	23.6	49.1	9.7	49.5	52.1	176
Number of marital control behaviours displayed by husband²						
0	8.9	28.2	2.7	28.2	30.6	268
1-2	20.6	44.5	5.6	45.7	49.2	332
3-4	39.3	60.8	15.5	64.3	68.4	106
5-6	(58.7)	(80.9)	(23.7)	(83.0)	(83.0)	19
Number of decisions in which women participate³						
0	17.9	36.5	7.6	39.2	42.4	118
1-2	26.4	53.7	8.2	54.5	58.1	156
3	16.3	37.0	4.1	37.8	41.0	391
Number of reasons for which wife beating is justified⁴						
0	11.6	30.6	2.4	30.6	32.3	112
1-2	14.8	35.1	3.5	36.3	38.0	196
3-4	25.0	47.9	8.6	49.0	53.7	233
5-6	28.4	51.6	8.9	53.1	57.0	122
7	16.7	41.4	10.2	44.1	45.5	63

Continued...

Table 99 Spousal violence by husband's characteristics and empowerment indicators—Continued

Percentage of ever-married women age 15-49 who have ever suffered emotional, physical, or sexual violence committed by their husband, according to his characteristics, marital characteristics, and selected empowerment indicators, Telangana, 2015-16

Background characteristic	Emotional violence	Physical violence	Sexual violence	Physical or sexual violence	Emotional, physical, or sexual violence	Number of women
Number of reasons given for refusing to have sexual intercourse with husband⁵						
0	25.8	40.4	10.0	42.9	44.5	94
1-2	23.2	50.4	5.9	51.1	53.7	165
3	17.8	39.1	5.9	40.1	43.6	467
Afraid of husband						
Most of the time	41.1	54.0	13.0	57.7	60.8	136
Sometimes	17.8	45.2	5.9	45.7	48.2	412
Never	9.0	24.6	2.7	25.3	29.5	177
Total	20.0	41.8	6.5	43.0	46.0	726

Note: Husband refers to the current husband for currently married women and the most recent husband for widowed, divorced, separated, or deserted women. Total includes women whose husbands drink alcohol but never get drunk and currently married women who are older or the same age as their husband, who are not shown separately.

¹ Currently married women only

² Behaviours include: Is jealous or angry if she talks to other men, frequently accuses her of being unfaithful, does not permit her to meet her female friends, tries to limit her contact with her family, insists on knowing where she is at all times, and does not trust her with any money

³ Currently married women only. Decisions included are decisions about own health care, major household purchases, and visits to her family or relatives.

⁴ Reasons given for which wife beating is justified include: she goes out without telling him, she neglects the house or children, she argues with him, she refuses to have sexual intercourse with him, she doesn't cook properly, he suspects she is unfaithful, and she shows disrespect for in-laws

⁵ Reasons given for refusing to have sexual intercourse with husband include: she knows her husband has a sexually transmitted disease, she knows her husband has sex with other women, and she is tired or not in the mood

() Based on 25-49 unweighted cases

Table 100 Injuries to women due to spousal violence

Percentage of ever-married women age 15-49 who have experienced spousal violence by types of injuries resulting from what their husband did to them, the type of violence, and whether they have experienced the violence ever and in the 12 months preceding the survey, Telangana, 2015-16

Type of spousal violence experienced	Percentage of women who have had:					Number of ever-married women
	Cuts, bruises, or aches	Severe burns	Eye injuries, sprains, dislocations, or minor burns	Deep wounds, broken bones, broken teeth, or any other serious injury	Any of these injuries	
Experienced physical violence¹						
Ever ²	21.2	5.5	7.6	3.5	23.4	304
In the past 12 months	21.2	5.6	8.5	3.4	23.9	249
Experienced sexual violence						
Ever ²	31.7	9.9	12.5	8.6	37.1	47
In the past 12 months	(29.8)	(11.1)	(12.6)	(5.5)	(34.5)	37
Experienced physical or sexual violence						
Ever ²	20.9	5.4	7.4	3.4	23.0	312
In the past 12 months	20.7	5.4	8.3	3.3	23.3	255
Experienced physical and sexual violence						
Ever ²	(36.7)	(12.0)	(15.2)	(10.4)	(43.2)	39
In the past 12 months	(36.3)	(13.6)	(15.4)	(6.7)	(42.0)	30

Note: Husband refers to the current husband for currently married women and the most recent husband for widowed, divorced, separated, or deserted women.

¹ Excludes women who reported violence only in response to a direct question on violence during pregnancy

² Includes violence in the past 12 months

() Based on 25-49 unweighted cases

Table 101 Help seeking

Percentage of women age 15-49 who have ever experienced physical or sexual violence by whether they have ever sought help, and among those who have sought help from any source, the source from which help was sought, according to the type of violence experienced and marital status, Telangana, 2015-16

Source	Type of violence experienced			Marital status		Total
	Physical only	Sexual only	Both physical and sexual	Ever-married	Never married	
Help seeking						
Never sought help and never told anyone	77.0	*	53.2	73.9	*	74.4
Never sought help but told someone	9.8	*	8.6	8.9	*	9.4
Sought help	13.3	*	38.2	17.3	*	16.2
Number of women who experienced violence	348	10	44	363	39	402
Sources of help among those who sought any help¹						
Own family	60.7	*	*	68.9	*	68.1
Husband's family	20.9	*	*	18.4	*	17.7
Friend	4.5	*	*	4.0	*	3.9
Neighbour	12.7	*	*	10.2	*	9.9
Religious leader	7.1	*	*	7.1	*	6.8
Police	3.0	*	*	2.2	*	2.2
Social service organization	4.3	*	*	3.2	*	3.1
Other	4.6	*	*	1.3	*	3.2
Number of women who experienced violence and sought help	46	2	17	63	2	65

¹ Women can report more than one source from which they sought help.

* Percentage not shown; based on fewer than 25 unweighted cases

APPENDIX

ESTIMATES OF SAMPLING ERRORS

The estimates from a sample survey are affected by two types of errors: (1) non-sampling errors, and (2) sampling errors. Non-sampling errors are the results of mistakes made in implementing data collection and data processing, such as failure to locate and interview the correct household, misunderstanding of the questions on the part of either the interviewer or the respondent, and data entry errors. Although numerous efforts were made during the implementation of the fourth National Family Health Survey (NFHS-4) to minimize this type of error, non-sampling errors are impossible to avoid and difficult to evaluate statistically. Sampling errors, on the other hand, can be evaluated statistically. The sample of respondents selected in NFHS-4 is only one of many samples that could have been selected from the same population, using the same design and expected sample size. Each of these samples would yield results that differ somewhat from the results of the actual sample selected. Sampling errors are a measure of the variability among all possible samples. Although the degree of variability is not known exactly, it can be estimated from the survey results.

A sampling error is usually measured in terms of the standard error for a particular statistic (mean, percentage, etc.), which is the square root of the variance. The standard error can be used to calculate confidence intervals within which the true value for the population can reasonably be assumed to fall. For example, for any given statistic calculated from a sample survey, the value of that statistic will fall within a range of plus or minus two times the standard error of that statistic in 95 percent of all possible samples of identical size and design. If the sample of respondents had been selected as a simple random sample, it would have been possible to use straightforward formulas for calculating sampling errors. However, the NFHS-4 sample is the result of a multi-stage stratified design, and, consequently, it was necessary to use more complex formulae. The computer software used to calculate sampling errors for NFHS-4 is programmed in SAS. This procedure uses the Taylor linearization method for variance estimation for survey estimates that are means or proportions. The Jackknife repeated replication method is used for variance estimation of more complex statistics such as total fertility rates and child mortality rates.

The Taylor linearization method treats any proportion or mean as a ratio estimate, $r = y/x$, where y represents the total sample value for variable y , and x represents the total number of cases in the group or subgroup under consideration. The variance of r is computed using the formula given below, with the standard error being the square root of the variance:

$$SE^2(r) = var(r) = \frac{1-f}{x^2} \sum_{h=1}^H \left[\frac{m_h}{m_h - 1} \left(\sum_{i=1}^{m_h} z_{hi}^2 - \frac{z_h^2}{m_h} \right) \right]$$

in which

$$z_{hi} = y_{hi} - rx_{hi}, \text{ and } z_h = y_h - rx_h$$

where

- h represents the stratum, which varies from 1 to H ,
- m_h is the total number of clusters selected in the h^{th} stratum,
- y_{hi} is the sum of the weighted values of variable y in the i^{th} cluster in the h^{th} stratum,
- x_{hi} is the sum of the weighted number of cases in the i^{th} cluster in the h^{th} stratum, and
- f is the overall sampling fraction, which is so small that it is ignored.

The Jackknife repeated replication method derives estimates of complex rates from each of several replications of the parent sample, and calculates standard errors for these estimates using simple formulae. Each replication considers all but one cluster in the calculation of the estimates. Pseudo-independent replications are thus created. In the NFHS-4 sample for Telangana, there were 380 clusters. Hence, 380 replications were created. The variance of a rate r is calculated as follows:

$$SE^2(r) = var(r) = \frac{1}{k(k-1)} \sum_{i=1}^k (r_i - r)^2$$

in which

$$r_i = kr - (k-1)r_{(i)}$$

where

- r is the estimate computed from the full sample of 380 clusters,
- $r_{(i)}$ is the estimate computed from the reduced sample of 379 clusters (i^{th} cluster excluded), and
- k is the total number of clusters.

In addition to the standard error, the design effect (DEFT) for each estimate is also computed, which is defined as the ratio between the standard error using the given sample design and the standard error that would result if a simple random sample had been used. A DEFT value of 1.0 indicates that the sample design is as efficient as a simple random sample, while a value greater than 1.0 indicates the increase in the sampling error due to the use of a more complex and less statistically efficient design. The relative standard error (SE/R) and confidence limits ($R \pm 2SE$) for each estimate are also computed.

Sampling errors for NFHS-4 are calculated for selected variables considered to be of primary interest. The results are presented in this appendix for Telangana as a whole and for the urban and rural areas of the state. For each variable, the type of statistic (mean, proportion, rate, or ratio) and the base population are given in Table A.1. Table A.2 presents the value of the statistic (R), its standard error (SE), the number of unweighted (N) and weighted (WN) cases, the design effect (DEFT), the relative standard error (SE/R), and the 95 percent confidence limits ($R \pm 2SE$); for each variable. The DEFT is considered undefined when the standard error for a simple random sample is zero (when the estimate is close to 0 or 1).

Table A.1 List of variables for sampling errors, Telangana, 2015-16

Variable	Estimate	Base population
HOUSEHOLDS		
Using an improved source of drinking water	Proportion	Households
Using an improved sanitation facility	Proportion	Households
Using iodized salt	Proportion	Households
Sex ratio (females per 1,000 males)	Ratio	<i>De facto</i> household population, all ages
WOMEN		
No schooling (Females age 6 years and above)	Proportion	<i>De facto</i> household population of females age 6 and above
Urban residence	Proportion	Women age 15-49
No schooling (Women age 15-49)	Proportion	Women age 15-49
Completed 10 or more years of schooling	Proportion	Women age 15-49
Never married, including married <i>gauna</i> not performed	Proportion	Women age 15-49
Currently married	Proportion	Women age 15-49
Married before age 18	Proportion	Women age 20-49
Currently using any contraceptive method	Proportion	Currently married women age 15-49
Currently using a modern contraceptive method	Proportion	Currently married women age 15-49
Currently using a traditional contraceptive method	Proportion	Currently married women age 15-49
Currently using pill	Proportion	Currently married women age 15-49
Currently using IUD/PPIUD	Proportion	Currently married women age 15-49
Currently using condom/ <i>Nirodha</i>	Proportion	Currently married women age 15-49
Currently using female sterilization	Proportion	Currently married women age 15-49
Using public health sector source of contraception	Proportion	Women age 15-49 currently using modern methods of contraception
Unmet need for family planning	Proportion	Currently married women age 15-49
Want no more children	Proportion	Currently married women age 15-49
Want to delay next birth at least 2 years	Proportion	Currently married women age 15-49
Mother received four or more antenatal care (ANC) visits	Proportion	Women with at least one birth in last five years (last birth)
Took iron and folic acid (IFA) for 100 days or more	Proportion	Women with at least one birth in last five years (last birth)
Birth registration	Proportion	<i>De jure</i> children under age 5 years
Births delivered by a health personnel	Proportion	Births in last 5 years
Institutional delivery	Proportion	Births in last 5 years
Postnatal check for mother within 2 days of birth	Proportion	Women with at least one birth in last five years (last birth)
Postnatal check for newborn within 2 days of birth	Proportion	Women with at least one birth in last five years (last birth)
Exclusive breastfeeding	Proportion	Children under age 6 months
Children with diarrhoea	Proportion	Children under age 5 years
Treated with oral rehydration salt (ORS) packets	Proportion	Children under age 5 years with diarrhoea in last 2 weeks
Children with diarrhoea taken to a health provider	Proportion	Children under age 5 years with diarrhoea in last 2 weeks
Child received BCG vaccination	Proportion	Children age 12-23 months
Child received DPT vaccination (3 doses)	Proportion	Children age 12-23 months
Child received polio vaccination (3 doses)	Proportion	Children age 12-23 months
Child received measles vaccination	Proportion	Children age 12-23 months
Child received hepatitis B vaccination (3 doses)	Proportion	Children age 12-23 months
Child with all basic vaccinations	Proportion	Children age 12-23 months
Children given vitamin A supplement in past 6 months	Proportion	Children age 9-59 months
Height-for-age, stunting (below -2SD)	Proportion	Children under age 5 years who were measured
Weight-for-height, wasting (below -2SD)	Proportion	Children under age 5 years who were measured
Weight-for-age, underweight (below -2SD)	Proportion	Children under age 5 years who were measured
Children with any anaemia	Proportion	Children age 6-59 months with an anaemia test
Women with any anaemia	Proportion	Women age 15-49 with an anaemia test
Body mass index (BMI) <18.5 kg/m ²	Proportion	Women age 15-49 who were measured
Body mass index (BMI) ≥25.0 kg/m ²	Proportion	Women age 15-49 who were measured
Have heard of HIV/AIDS	Proportion	Women age 15-49
Have comprehensive knowledge about HIV/AIDS	Proportion	Women age 15-49
Ever experienced physical or sexual violence	Proportion	Women age 15-49
Total fertility rate (last 3 years)	Rate	Women
Neonatal mortality	Rate	Births in last 5 years
Postneonatal mortality	Rate	Births in last 5 years
Infant mortality	Rate	Births in last 5 years
Child mortality	Rate	Births in last 5 years
Under-five mortality	Rate	Births in last 5 years
MEN		
No schooling (Males age 6 years and above)	Proportion	<i>De facto</i> household population of males age 6 and above
Urban residence	Proportion	Men age 15-49
No schooling (Men age 15-49)	Proportion	Men age 15-49
Completed 10 or more years of schooling	Proportion	Men age 15-49
Never married, including married <i>gauna</i> not performed	Proportion	Men age 15-49
Currently married	Proportion	Men age 15-49
Married before age 21	Proportion	Men age 25-49
Want no more children	Proportion	Currently married men age 15-49
Want to delay next birth at least 2 years	Proportion	Currently married men age 15-49
Men with any anaemia	Proportion	Men age 15-49 with an anaemia test
Body mass index (BMI) <18.5 kg/m ²	Proportion	Men age 15-49 who were measured
Body mass index (BMI) ≥25.0 kg/m ²	Proportion	Men age 15-49 who were measured
Have heard of HIV/AIDS	Proportion	Men age 15-49
Have comprehensive knowledge about HIV/AIDS	Proportion	Men age 15-49

Table A.2 Sampling errors: Total sample, Telangana, 2015-16

Variable	Value (R)	Standard error (SE)	Number of cases		Design effect (DEFT)	Relative standard error (SE/R)	Confidence limits	
			Unweighted (N)	Weighted (WN)			R-2SE	R+2SE
HOUSEHOLDS								
Using an improved source of drinking water	0.776	0.012	7,786	7,786	2.553	0.016	0.752	0.800
Using an improved sanitation facility	0.502	0.016	7,786	7,786	2.784	0.031	0.471	0.534
Using iodized salt	0.958	0.004	7,722	7,723	1.960	0.005	0.949	0.967
Sex ratio (females per 1,000 males)	1007.265	10.379	14,709	14,837	1.257	0.010	986.507	1028.024
WOMEN								
No schooling (Females age 6 years and above)	0.377	0.007	13,528	13,481	1.675	0.018	0.363	0.391
Urban residence	0.478	0.011	7,567	7,567	1.973	0.024	0.456	0.501
No schooling (Women age 15-49)	0.325	0.009	7,567	7,567	1.622	0.027	0.307	0.342
Completed 10 or more years of schooling	0.433	0.011	7,567	7,567	1.967	0.026	0.411	0.456
Never married, including married <i>gauna</i> not performed	0.184	0.006	7,567	7,567	1.426	0.035	0.171	0.197
Currently married	0.743	0.007	7,567	7,567	1.424	0.010	0.728	0.757
Married before age 18	0.433	0.009	6,514	6,570	1.492	0.021	0.415	0.452
Currently using any contraceptive method	0.572	0.010	5,603	5,619	1.451	0.017	0.552	0.591
Currently using a modern contraceptive method	0.569	0.010	5,603	5,619	1.438	0.017	0.550	0.588
Currently using a traditional contraceptive method	0.002	0.001	5,603	5,619	1.378	0.389	0.000	0.004
Currently using pill	0.003	0.001	5,603	5,619	1.087	0.273	0.001	0.004
Currently using IUD/PPIUD	0.003	0.001	5,603	5,619	1.372	0.316	0.001	0.005
Currently using condom/ <i>Nirodh</i>	0.005	0.001	5,603	5,619	1.140	0.223	0.003	0.007
Currently using female sterilization	0.542	0.010	5,603	5,619	1.444	0.018	0.522	0.561
Using public health sector source of contraception	0.667	0.015	3,437	3,523	1.823	0.022	0.638	0.696
Unmet need for family planning	0.073	0.004	5,603	5,619	1.162	0.055	0.065	0.081
Want no more children	0.678	0.008	5,603	5,619	1.342	0.012	0.662	0.695
Want to delay next birth at least 2 years	0.054	0.004	5,603	5,619	1.246	0.070	0.047	0.062
Mother received four or more antenatal care (ANC) visits	0.750	0.015	1,778	1,793	1.495	0.020	0.719	0.780
Took iron and folic acid (IFA) for 100 days or more	0.528	0.022	1,778	1,793	1.850	0.041	0.485	0.572
Birth registration	0.829	0.012	2,564	2,577	1.361	0.015	0.804	0.853
Births delivered by a health personnel	0.914	0.008	2,419	2,411	1.182	0.009	0.898	0.930
Institutional delivery	0.915	0.008	2,419	2,411	1.160	0.008	0.900	0.931
Postnatal check for mother within 2 days of birth	0.827	0.014	1,778	1,793	1.619	0.017	0.798	0.856
Postnatal check for newborn within 2 days of birth	0.265	0.015	1,778	1,793	1.452	0.057	0.235	0.295
Exclusive breastfeeding	0.673	0.037	193	184	1.057	0.054	0.600	0.746
Children with diarrhoea	0.084	0.007	2,349	2,347	1.246	0.088	0.069	0.098
Treated with oral rehydration salt (ORS) packets	0.568	0.040	202	196	1.088	0.070	0.488	0.647
Children with diarrhoea taken to a health provider	0.740	0.036	202	196	1.149	0.049	0.668	0.813
Child received BCG vaccination	0.974	0.009	484	495	1.322	0.010	0.955	0.993
Child received DPT vaccination (3 doses)	0.879	0.017	484	495	1.180	0.020	0.845	0.914
Child received polio vaccination (3 doses)	0.753	0.024	484	495	1.211	0.032	0.706	0.801
Child received measles vaccination	0.906	0.015	484	495	1.172	0.017	0.875	0.937
Child received hepatitis B vaccination (3 doses)	0.706	0.026	484	495	1.270	0.037	0.653	0.758
Child with all basic vaccinations	0.681	0.026	484	495	1.213	0.038	0.629	0.732
Children given vitamin A supplement in past 6 months	0.763	0.017	2,022	2,039	1.673	0.023	0.728	0.798
Height-for-age, stunting (below -2SD)	0.281	0.014	2,087	2,061	1.283	0.049	0.253	0.308
Weight-for-height, wasting (below -2SD)	0.180	0.010	2,087	2,061	1.157	0.056	0.160	0.200
Weight-for-age, underweight (below -2SD)	0.285	0.014	2,087	2,061	1.288	0.049	0.257	0.313
Children with any anaemia	0.607	0.015	1,809	1,795	1.298	0.025	0.577	0.638
Women with any anaemia	0.567	0.008	6,990	6,866	1.294	0.014	0.552	0.583
Body mass index (BMI) <18.5 kg/m ²	0.231	0.006	6,822	6,726	1.235	0.027	0.218	0.244
Body mass index (BMI) ≥25.0 kg/m ²	0.282	0.008	6,822	6,726	1.507	0.029	0.266	0.299
Have heard of HIV/AIDS	0.901	0.011	1,302	1,285	1.385	0.013	0.878	0.924
Have comprehensive knowledge about HIV/AIDS	0.295	0.023	1,302	1,285	1.826	0.078	0.249	0.342
Ever experienced physical or sexual violence	0.449	0.030	895	895	1.829	0.068	0.388	0.510
Total fertility rate (last 3 years)	1.787	0.053	21,912	21,956	1.161	0.030	1.681	1.893
Neonatal mortality	21.945	3.124	2,480	2,477	0.963	0.142	15.698	28.192
Postneonatal mortality	7.844	1.949	2,482	2,484	1.028	0.248	3.946	11.743
Infant mortality	29.789	3.816	2,481	2,478	1.010	0.128	22.157	37.422
Child mortality	3.923	1.387	2,492	2,524	1.153	0.354	1.148	6.698
Under-five mortality	33.596	4.194	2,486	2,482	1.048	0.125	25.207	41.984
MEN								
No schooling (Males age 6 years and above)	0.205	0.007	13,091	13,207	1.843	0.032	0.191	0.218
Urban residence	0.478	0.019	1,054	1,058	1.254	0.040	0.439	0.517
No schooling (Men age 15-49)	0.170	0.013	1,054	1,058	1.136	0.077	0.143	0.196
Completed 10 or more years of schooling	0.546	0.022	1,054	1,058	1.435	0.040	0.502	0.590
Never married, including married <i>gauna</i> not performed	0.359	0.016	1,054	1,058	1.102	0.045	0.327	0.392
Currently married	0.629	0.017	1,054	1,058	1.127	0.027	0.595	0.662
Married before age 21	0.248	0.016	719	728	0.977	0.063	0.217	0.280
Want no more children	0.722	0.020	670	665	1.174	0.028	0.681	0.762
Want to delay next birth at least 2 years	0.045	0.010	670	665	1.190	0.213	0.026	0.064
Men with any anaemia	0.154	0.013	964	959	1.011	0.082	0.129	0.179
Body mass index (BMI) <18.5 kg/m ²	0.214	0.013	972	966	0.980	0.061	0.188	0.241
Body mass index (BMI) ≥25.0 kg/m ²	0.242	0.020	972	966	1.371	0.082	0.202	0.282
Have heard of HIV/AIDS	0.959	0.009	1,054	1,058	1.419	0.009	0.942	0.976
Have comprehensive knowledge about HIV/AIDS	0.498	0.028	1,054	1,058	1.802	0.056	0.443	0.554

Table A.3 Sampling errors: Urban sample, Telangana, 2015-16

Variable	Value (R)	Standard error (SE)	Number of cases		Design effect (DEFT)	Relative standard error (SE/R)	Confidence limits	
			Unweighted (N)	Weighted (WN)			R-2SE	R+2SE
HOUSEHOLDS								
Using an improved source of drinking water	0.801	0.019	2,739	3,465	2.490	0.024	0.763	0.839
Using an improved sanitation facility	0.644	0.029	2,739	3,465	3.144	0.045	0.587	0.702
Using iodized salt	0.992	0.002	2,718	3,440	1.341	0.002	0.988	0.997
Sex ratio (females per 1,000 males)	976.073	17.031	5,508	6,942	1.293	0.017	942.010	1010.135
WOMEN								
No schooling (Females age 6 years and above)	0.233	0.011	4,866	6,083	1.948	0.049	0.210	0.256
Currently using any contraceptive method	0.588	0.015	1,981	2,637	1.345	0.025	0.558	0.617
Currently using a modern contraceptive method	0.583	0.015	1,981	2,637	1.324	0.025	0.554	0.612
Currently using a traditional contraceptive method	0.005	0.002	1,981	2,637	1.226	0.409	0.001	0.008
Currently using pill	0.005	0.002	1,981	2,637	0.996	0.315	0.002	0.008
Currently using IUD/PPIUD	0.006	0.002	1,981	2,637	1.226	0.348	0.002	0.011
Currently using condom/Nirodh	0.008	0.002	1,981	2,637	1.049	0.265	0.004	0.012
Currently using female sterilization	0.549	0.015	1,981	2,637	1.322	0.027	0.519	0.578
Using public health sector source of contraception	0.577	0.026	1,224	1,675	1.861	0.046	0.524	0.630
Unmet need for family planning	0.088	0.007	1,981	2,637	1.121	0.081	0.074	0.102
Want no more children	0.693	0.014	1,981	2,637	1.307	0.020	0.666	0.720
Mother received four or more antenatal care (ANC) visits	0.776	0.027	619	841	1.613	0.034	0.722	0.829
Took iron and folic acid (IFA) for 100 days or more	0.588	0.039	619	841	1.978	0.066	0.511	0.666
Birth registration	0.899	0.016	960	1,221	1.346	0.018	0.868	0.931
Births delivered by a health personnel	0.947	0.010	851	1,129	1.092	0.011	0.928	0.967
Institutional delivery	0.963	0.009	851	1,129	1.236	0.009	0.945	0.981
Postnatal check for mother within 2 days of birth	0.852	0.027	619	841	1.916	0.032	0.798	0.906
Postnatal check for newborn within 2 days of birth	0.217	0.024	619	841	1.479	0.112	0.168	0.265
Children with diarrhoea	0.081	0.014	831	1,106	1.401	0.167	0.054	0.108
Treated with oral rehydration salt (ORS) packets	0.618	0.061	71	90	1.025	0.099	0.495	0.741
Children with diarrhoea taken to a health provider	0.795	0.058	71	90	1.143	0.073	0.679	0.911
Child received BCG vaccination	0.976	0.015	179	246	1.381	0.016	0.945	1.007
Child received DPT vaccination (3 doses)	0.895	0.027	179	246	1.194	0.030	0.842	0.949
Child received polio vaccination (3 doses)	0.744	0.040	179	246	1.224	0.054	0.663	0.824
Child received measles vaccination	0.918	0.025	179	246	1.227	0.027	0.868	0.967
Child received hepatitis B vaccination (3 doses)	0.697	0.045	179	246	1.310	0.065	0.606	0.787
Child with all basic vaccinations	0.678	0.044	179	246	1.247	0.065	0.590	0.765
Children given vitamin A supplement in past 6 months	0.753	0.031	733	983	1.721	0.041	0.691	0.814
Height-for-age, stunting (below -2SD)	0.209	0.023	688	870	1.318	0.109	0.163	0.254
Weight-for-height, wasting (below -2SD)	0.146	0.017	688	870	1.235	0.115	0.113	0.180
Weight-for-age, underweight (below -2SD)	0.222	0.025	688	870	1.441	0.115	0.171	0.272
Children with any anaemia	0.516	0.027	600	765	1.310	0.052	0.462	0.570
Women with any anaemia	0.550	0.013	2,441	3,113	1.253	0.023	0.525	0.576
Body mass index (BMI) <18.5 kg/m ²	0.161	0.010	2,384	3,067	1.327	0.063	0.141	0.181
Body mass index (BMI) ≥25.0 kg/m ²	0.397	0.015	2,384	3,067	1.440	0.037	0.367	0.426
Have heard of HIV/AIDS	0.948	0.012	499	614	1.170	0.012	0.925	0.972
Have comprehensive knowledge about HIV/AIDS	0.391	0.041	499	614	1.869	0.105	0.309	0.473
Total fertility rate (last 3 years)	1.671	0.078	8,060	10,544	1.150	0.046	1.516	1.826
Neonatal mortality	12.223	3.535	876	1,167	0.912	0.289	5.153	19.294
Postneonatal mortality	8.123	3.102	879	1,174	1.026	0.382	1.920	14.327
Infant mortality	20.347	4.895	876	1,167	0.991	0.241	10.556	30.137
Child mortality	4.625	2.498	902	1,222	1.183	0.540	0.000	9.621
Under-five mortality	24.877	5.731	877	1,168	1.063	0.230	13.415	36.340
MEN								
No schooling (Males age 6 years and above)	0.115	0.011	4,924	6,186	2.171	0.091	0.094	0.137
Want no more children	0.698	0.037	226	302	1.201	0.053	0.625	0.772
Men with any anaemia	0.102	0.022	326	437	1.192	0.211	0.059	0.145
Body mass index (BMI) <18.5 kg/m ²	0.176	0.021	329	439	1.003	0.119	0.134	0.218
Body mass index (BMI) ≥25.0 kg/m ²	0.319	0.037	329	439	1.330	0.116	0.244	0.393
Have heard of HIV/AIDS	0.961	0.015	380	506	1.541	0.016	0.930	0.992
Have comprehensive knowledge about HIV/AIDS	0.551	0.047	380	506	1.825	0.085	0.457	0.645

Table A.4 Sampling errors: Rural sample, Telangana, 2015-16

Variable	Value (R)	Standard error (SE)	Number of cases		Design effect (DEFT)	Relative standard error (SE/R)	Confidence limits	
			Unweighted (N)	Weighted (WN)			R-2SE	R+2SE
HOUSEHOLDS								
Using an improved source of drinking water	0.756	0.015	5,047	4,321	2.557	0.020	0.725	0.787
Using an improved sanitation facility	0.389	0.013	5,047	4,321	1.900	0.034	0.363	0.415
Using iodized salt	0.930	0.008	5,004	4,284	2.149	0.008	0.914	0.945
Sex ratio (females per 1,000 males)	1034.695	11.957	9,201	7,895	1.112	0.012	1010.782	1058.609
WOMEN								
No schooling (Females age 6 years and above)	0.495	0.007	8,662	7,398	1.352	0.014	0.481	0.509
Currently using any contraceptive method	0.558	0.012	3,622	2,983	1.492	0.022	0.533	0.582
Currently using a modern contraceptive method	0.557	0.012	3,622	2,983	1.491	0.022	0.533	0.582
Currently using a traditional contraceptive method	0.000	0.000	3,622	2,983	0.897	1.001	0.000	0.001
Currently using pill	0.001	0.001	3,622	2,983	1.022	0.576	0.000	0.002
Currently using IUD/PPIUD	0.001	0.000	3,622	2,983	0.990	0.583	0.000	0.002
Currently using condom/Nirodh	0.002	0.001	3,622	2,983	1.019	0.399	0.000	0.003
Currently using female sterilization	0.536	0.012	3,622	2,983	1.504	0.023	0.511	0.560
Using public health sector source of contraception	0.749	0.013	2,213	1,848	1.377	0.017	0.723	0.774
Unmet need for family planning	0.060	0.004	3,622	2,983	1.070	0.070	0.051	0.068
Want no more children	0.666	0.010	3,622	2,983	1.308	0.015	0.645	0.686
Mother received four or more antenatal care (ANC) visits	0.727	0.016	1,159	952	1.214	0.022	0.695	0.758
Took iron and folic acid (IFA) for 100 days or more	0.475	0.019	1,159	952	1.279	0.040	0.438	0.513
Birth registration	0.765	0.017	1,604	1,356	1.338	0.022	0.731	0.799
Births delivered by a health personnel	0.884	0.012	1,568	1,282	1.218	0.013	0.861	0.907
Institutional delivery	0.873	0.012	1,568	1,282	1.175	0.013	0.850	0.897
Postnatal check for mother within 2 days of birth	0.805	0.015	1,159	952	1.257	0.018	0.776	0.834
Postnatal check for newborn within 2 days of birth	0.307	0.017	1,159	952	1.234	0.054	0.274	0.341
Children with diarrhoea	0.086	0.007	1,518	1,241	0.910	0.080	0.072	0.099
Treated with oral rehydration salt (ORS) packets	0.525	0.051	131	106	1.106	0.097	0.423	0.627
Children with diarrhoea taken to a health provider	0.694	0.042	131	106	1.034	0.060	0.611	0.777
Child received BCG vaccination	0.972	0.011	305	249	1.161	0.011	0.950	0.994
Child received DPT vaccination (3 doses)	0.863	0.021	305	249	1.079	0.025	0.821	0.906
Child received polio vaccination (3 doses)	0.763	0.026	305	249	1.051	0.034	0.712	0.815
Child received measles vaccination	0.894	0.018	305	249	1.032	0.020	0.857	0.930
Child received hepatitis B vaccination (3 doses)	0.714	0.027	305	249	1.056	0.038	0.659	0.769
Child with all basic vaccinations	0.683	0.027	305	249	1.023	0.040	0.628	0.738
Children given vitamin A supplement in past 6 months	0.772	0.018	1,289	1,056	1.395	0.023	0.737	0.808
Height-for-age, stunting (below -2SD)	0.333	0.015	1,399	1,191	1.111	0.045	0.303	0.363
Weight-for-height, wasting (below -2SD)	0.204	0.012	1,399	1,191	1.028	0.057	0.181	0.227
Weight-for-age, underweight (below -2SD)	0.331	0.014	1,399	1,191	1.049	0.044	0.302	0.360
Children with any anaemia	0.675	0.015	1,209	1,030	1.096	0.022	0.645	0.706
Women with any anaemia	0.581	0.009	4,549	3,753	1.288	0.016	0.562	0.600
Body mass index (BMI) <18.5 kg/m ²	0.290	0.008	4,438	3,659	1.141	0.027	0.274	0.305
Body mass index (BMI) ≥25.0 kg/m ²	0.186	0.008	4,438	3,659	1.286	0.040	0.171	0.201
Have heard of HIV/AIDS	0.857	0.019	803	671	1.513	0.022	0.820	0.895
Have comprehensive knowledge about HIV/AIDS	0.208	0.020	803	671	1.391	0.096	0.168	0.248
Total fertility rate (last 3 years)	1.883	0.072	13,852	11,412	1.130	0.038	1.739	2.027
Neonatal mortality	30.630	4.700	1,604	1,310	1.009	0.153	21.229	40.031
Postneonatal mortality	7.582	2.444	1,603	1,310	0.967	0.322	2.694	12.471
Infant mortality	38.212	5.420	1,605	1,311	1.026	0.142	27.373	49.052
Child mortality	3.221	1.335	1,590	1,301	0.943	0.414	0.551	5.890
Under-five mortality	41.310	5.766	1,609	1,314	1.031	0.140	29.778	52.842
MEN								
No schooling (Males age 6 years and above)	0.283	0.006	8,167	7,022	1.230	0.021	0.271	0.295
Want no more children	0.741	0.021	444	363	1.030	0.029	0.698	0.784
Men with any anaemia	0.198	0.014	638	523	0.836	0.071	0.170	0.225
Body mass index (BMI) <18.5 kg/m ²	0.246	0.017	643	527	0.966	0.068	0.213	0.280
Body mass index (BMI) ≥25.0 kg/m ²	0.179	0.019	643	527	1.243	0.106	0.141	0.217
Have heard of HIV/AIDS	0.958	0.009	674	552	1.128	0.009	0.940	0.975
Have comprehensive knowledge about HIV/AIDS	0.450	0.031	674	552	1.604	0.068	0.389	0.512



Technical assistance for NFHS-4 was provided by the USAID-supported DHS Program at ICF, and assistance for the HIV components was provided by NACO and NARI. Funding assistance was provided by Ministry of Health and Family Welfare, Government of India and:



The opinions in this publication do not necessarily reflect the views of the funding agencies.
For additional information on NFHS-4, visit <http://www.rchiips.org/nfhs>
For related information, visit <http://www.iipsindia.org> or <http://www.mohfw.nic.in>