

Key Indicators for Madhya Pradesh from NFHS-3	NFHS-3 (2005-06)	Residence		Education				NFHS-2 (1998-99)	NFHS-1 (1992-93)
		Urban	Rural	No education <sup>4</sup>	<5 years complete <sup>4</sup>	5-9 years complete <sup>4</sup>	10 years complete and above <sup>4</sup>		
<b>Marriage and Fertility</b>									
1. Women age 20-24 married by age 18 (%)	57.3	34.9	66.5	79.2	74.5	53.8	11.3	64.7	na
2. Men age 25-29 married by age 21 (%)	54.0	27.4	64.1	74.9	(67.7)	49.7	34.6	na	na
3. Total fertility rate (children per woman)	3.1	2.6	3.3	3.8	3.1	2.8	1.9	3.4	na
4. Women age 15-19 who were already mothers or pregnant at the time of the survey (%)	13.6	7.2	16.2	30.1	9.7	10.7	3.8	na	na
5. Median age at first birth for women age 25-49	19.4	20.3	19.1	18.8	18.9	19.6	22.8	18.7	na
6. Married women with 2 living children wanting no more children <sup>1</sup> (%)	84.3	88.8	82.0	78.1	82.2	84.5	97.3	60.1	na
6a. Two sons	92.1	91.2	92.5	89.2	*	93.0	97.3	79.5	na
6b. One son, one daughter	89.7	94.3	87.1	84.4	(95.9)	89.7	97.3	63.0	na
6c. Two daughters	37.6	57.0	27.8	(27.6)	*	(30.1)	(97.9)	19.5	na
<b>Family Planning (currently married women, age 15-49)</b>									
<b>Current use</b>									
7. Any method (%)	55.9	61.1	54.1	55.9	55.6	51.5	66.5	44.1	na
8. Any modern method (%)	52.8	56.3	51.5	53.8	52.8	48.4	56.8	42.8	na
8a. Female sterilization (%)	44.3	36.8	46.9	49.8	44.9	38.5	27.1	36.0	na
8b. Male sterilization (%)	1.3	1.6	1.2	1.3	0.8	1.3	1.3	1.9	na
8c. IUD (%)	0.7	1.6	0.3	0.3	0.0	0.7	3.3	0.7	na
8d. Pill (%)	1.7	3.6	1.0	0.8	2.5	2.9	3.4	1.0	na
8e. Condom (%)	4.8	12.6	2.0	1.7	4.5	5.0	21.8	3.1	na
<b>Unmet need for family planning</b>									
9. Total unmet need (%)	11.3	10.8	11.5	10.3	12.9	13.2	11.2	17.1	na
9a. For spacing (%)	5.4	4.8	5.6	4.3	5.6	8.0	5.4	9.2	na
9b. For limiting (%)	5.9	6.1	5.8	6.0	7.3	5.2	5.8	7.9	na
<b>Maternal and Child Health</b>									
<b>Maternity care (for births in the last 3 years)</b>									
10. Mothers who had at least 3 antenatal care visits for their last birth (%)	40.2	58.4	34.6	27.5	42.0	49.1	84.6	27.1	na
11. Mothers who consumed IFA for 90 days or more when they were pregnant with their last child (%)	11.8	20.2	9.2	5.2	18.0	13.8	38.3	na	na
12. Births assisted by a doctor/nurse/LHV/ANM/other health personnel (%) <sup>2</sup>	37.1	66.4	28.0	24.5	34.2	45.5	88.2	28.9	na
13. Institutional births (%) <sup>2</sup>	29.7	59.9	20.2	17.3	29.3	37.1	79.9	22.0	na
14. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/other health personnel within 2 days of delivery for their last birth (%) <sup>2</sup>	27.9	53.4	20.1	16.8	27.3	34.3	73.1	na	na
<b>Child immunization and vitamin A supplementation<sup>2</sup></b>									
15a. Children 12-23 months fully immunized (BCG, measles, and 3 doses each of polio/DPT) (%)	40.3	68.7	31.5	28.4	(46.5)	48.3	85.6	22.6	na
15b. Children 12-23 months who have received BCG (%)	80.5	91.4	77.0	74.2	(89.4)	84.7	100.0	62.0	na
15c. Children 12-23 months who have received 3 doses of polio vaccine (%)	75.6	87.6	71.8	69.5	(78.7)	81.0	94.4	56.6	na
15d. Children 12-23 months who have received 3 doses of DPT vaccine (%)	49.8	75.8	41.6	37.0	(53.6)	61.2	91.5	35.8	na
15e. Children 12-23 months who have received measles vaccine (%)	61.4	77.4	56.4	53.8	(57.1)	66.4	96.9	34.1	na
16. Children age 12-35 months who received a vitamin A dose in last 6 months (%)	20.1	25.8	18.3	14.7	11.0	27.7	34.1	na	na
<b>Treatment of childhood diseases (children under 3 years)<sup>2</sup></b>									
17. Children with diarrhoea in the last 2 weeks who received ORS (%)	28.6	30.6	27.8	18.8	(15.9)	39.4	(45.5)	29.8	na
18. Children with diarrhoea in the last 2 weeks taken to a health facility (%)	60.1	52.8	63.2	58.2	(50.3)	61.4	(71.0)	60.1	na
19. Children with acute respiratory infection or fever in the last 2 weeks taken to a health facility (%)	68.7	76.4	66.5	66.7	*	69.6	(92.6)	na	na
<b>Child Feeding Practices and Nutritional Status of Children<sup>2,3</sup></b>									
20. Children under 3 years breastfed within one hour of birth (%)	14.9	20.3	13.3	12.4	13.5	15.9	27.6	8.9	na
21. Children age 0-5 months exclusively breastfed (%)	21.6	21.9	21.5	19.3	*	20.2	(31.0)	na	na
22. Children age 6-9 months receiving solid or semi-solid food and breastmilk (%)	51.9	60.5	48.9	49.0	*	60.6	(43.3)	na	na
23. Children under 3 years who are stunted (%)	46.5	42.2	47.8	49.8	42.7	45.4	34.7	55.1	na
24. Children under 3 years who are wasted (%)	39.5	36.4	40.4	40.8	47.5	39.1	25.9	25.2	na
25. Children under 3 years who are underweight (%)	57.9	50.1	60.2	62.1	65.9	55.4	34.1	50.8	na
<b>Nutritional Status of Ever-Married Adults (age 15-49)</b>									
26. Women whose Body Mass Index is below normal (%)	40.1	28.7	44.2	44.5	49.0	35.4	20.0	35.2	na
27. Men whose Body Mass Index is below normal (%)	36.3	22.8	41.1	45.5	40.3	38.9	19.2	na	na
28. Women who are overweight or obese (%)	8.6	22.8	3.5	4.6	8.6	10.3	27.2	6.8	na
29. Men who are overweight or obese (%)	5.4	13.3	2.6	1.9	4.5	2.1	15.0	na	na
<b>Anaemia among Children and Adults</b>									
30. Children age 6-35 months who are anaemic (%)	82.6	75.2	84.9	85.7	79.9	81.7	69.9	71.3	na
31. Ever-married women age 15-49 who are anaemic (%)	57.7	48.3	61.0	61.3	58.0	55.5	42.2	49.3	na
32. Pregnant women age 15-49 who are anaemic (%)	57.9	44.3	62.4	60.9	*	55.7	48.3	49.9	na
33. Ever-married men age 15-49 who are anaemic (%)	24.7	15.7	27.9	30.3	30.9	24.5	14.7	na	na
<b>Knowledge of HIV/AIDS among Ever-Married Adults (age 15-49)</b>									
34. Women who have heard of AIDS (%)	45.3	74.1	35.0	24.2	48.0	72.9	97.3	23.7	na
35. Men who have heard of AIDS (%)	68.3	94.8	58.8	33.5	56.6	82.1	97.4	na	na
36. Women who know that consistent condom use can reduce the chances of getting HIV/AIDS (%)	35.7	63.3	25.7	15.7	37.2	58.9	91.7	na	na
37. Men who know that consistent condom use can reduce the chances of getting HIV/AIDS (%)	61.9	91.3	51.4	24.3	47.1	75.4	96.7	na	na
<b>Women's Empowerment</b>									
38. Currently married women who usually participate in household decisions (%)	29.4	39.9	25.6	28.5	29.2	27.6	38.3	na	na
39. Ever-married women who have ever experienced spousal violence (%)	45.7	43.7	46.4	51.0	56.8	41.1	17.8	na	na

na : Not available

( ) Based on 25-49 unweighted cases

\* Based on fewer than 25 unweighted cases

<sup>1</sup> Excludes pregnant women.

<sup>2</sup> Based on the last 2 births in the 3 years before the survey to ever-married women.

<sup>3</sup> Based on WHO standard.

<sup>4</sup> For children education refers to mother's education. Children with missing information on the mother's education are not included in the education columns.